

Algebraic Geometry III/IV

Problems, set 9.

Exercise 12. Let \mathcal{L} be the set of all projective lines in $\mathbb{P}_{\mathbb{C}}^2$, i.e.,

$$\mathcal{L} = \{C_{uX+vY+wZ} \mid (u, v, w) \neq 0\}.$$

There is a natural identification $\Phi : \mathcal{L} \rightarrow \mathbb{P}_{\mathbb{C}}^2$, given by $C_{uX+vY+wZ} \mapsto [u, v, w]$. This identification leads to the definition of the dual of a projective algebraic curve: Let $C \subset \mathbb{P}_{\mathbb{C}}^2$ is a non-singular algebraic curve and $\mathcal{T}(C)$ be the set of all tangent lines of C . Then the **dual** of C is the set $C^* = \Phi(\mathcal{T}(C)) \subset \mathbb{P}_{\mathbb{C}}^2$. Prove the following facts:

(a) Let

$$A = \begin{pmatrix} a & d & e \\ d & b & f \\ e & f & c \end{pmatrix}.$$

A projective conic $C_F \subset \mathbb{P}_{\mathbb{C}}^2$, given by the equation

$$F(X, Y, Z) = (X \ Y \ Z) A \begin{pmatrix} X \\ Y \\ Z \end{pmatrix},$$

is non-singular if and only if $\det A \neq 0$.

(b) Let $C_F \subset \mathbb{P}_{\mathbb{C}}^2$ be a non-singular conic and F as in (a). Let $P = [\alpha, \beta, \gamma] \in C_F$. Then the tangent line of C_F at P is given by the equation

$$(\alpha \ \beta \ \gamma) A \begin{pmatrix} X \\ Y \\ Z \end{pmatrix} = 0.$$

- (c) Let $C = C_F$ be a non-singular conic and F as in (a). Then the dual $C^* \subset \mathbb{P}_{\mathbb{C}}^2$ of the conic C is again a non-singular conic $C^* = C_G$, given by the equation

$$G(X, Y, Z) = (X \ Y \ Z) A^{-1} \begin{pmatrix} X \\ Y \\ Z \end{pmatrix}.$$

Exercise 13. Find the genus of a non-singular model of the irreducible curve $C_F \subset \mathbb{P}_{\mathbb{C}}^2$ with $F(X, Y, Z) = 3Y^4 + 4Y^3Z + X^4$.

Exercise 14. Find the genus of a non-singular model of the irreducible curve $C_F \subset \mathbb{P}_{\mathbb{C}}^2$ with $F(X, Y, Z) = Y^4 - 2X^2Y^2 + XZ^3$.

Exercise 15. Find the genus of a non-singular model of the irreducible curve $C_F \subset \mathbb{P}_{\mathbb{C}}^2$ with $F(X, Y, Z) = X^5 + 3Y^5 - 5Y^3Z^2$.