

# NVFEM: a Galerkin method for (fully) nonlinear elliptic equations

Omar Lakkis

Mathematics — University of Sussex — Brighton, England

based on joint work  
with

Tristan Pryer (University of Reading, GB)  
and Ellya Kawecki (University of Sussex, GB)

talk given  
15 July 2014



# Outline

- 1 PDE Background: fully nonlinear elliptic PDE's
- 2 History and competing approaches
  - Finite differences
  - Finite elements
- 3 Iterative nonlinear solvers
  - Fixed point
  - Newton
  - Hessian recovery
- 4 A non-variational FEM (NVFEM) solver
- 5 Convergence
- 6 Experiments
- 7 Conclusions
- 8 References

# Fully nonlinear elliptic PDE's

## Definition and notation

Given a real-valued nonlinear function  $F$  of matrices

$$(FNFun) \quad F : \text{Sym}(\mathbb{R}^{d \times d}) \rightarrow \mathbb{R}.$$

# Fully nonlinear elliptic PDE's

## Definition and notation

Given a real-valued nonlinear function  $F$  of matrices

$$(FNFun) \quad F : \text{Sym}(\mathbb{R}^{d \times d}) \rightarrow \mathbb{R}.$$

Consider the equation

$$(FNE) \quad \mathfrak{N}[\mathbf{u}] := F(D^2 \mathbf{u}) - \mathbf{f} = \mathbf{0} \text{ and } \mathbf{u}|_{\partial\Omega} = 0$$

# Fully nonlinear elliptic PDE's

## Definition and notation

Given a real-valued nonlinear function  $F$  of matrices

$$(FNFun) \quad F : \text{Sym}(\mathbb{R}^{d \times d}) \rightarrow \mathbb{R}.$$

Consider the equation

$$(FNE) \quad \mathfrak{N}[u] := F(D^2 u) - f = 0 \text{ and } u|_{\partial\Omega} = 0$$

Conditional ellipticity condition, i.e.,

$$(NL\text{-Ellip}) \quad \lambda(\mathbf{M}) \sup_{|\boldsymbol{\xi}|=1} |\mathbf{N}\boldsymbol{\xi}| \leq F(\mathbf{M} + \mathbf{N}) - F(\mathbf{M}) \leq \Lambda \sup_{|\boldsymbol{\xi}|=1} |\mathbf{N}\boldsymbol{\xi}|$$
$$\forall \mathbf{M} \in \mathfrak{C} \subseteq \text{Sym}(\mathbb{R}^{d \times d}), \mathbf{N} \in \text{Sym}(\mathbb{R}^{d \times d}).$$

for some ellipticity domain  $\mathfrak{C}$  and “constants”  $\lambda(\cdot), \Lambda > 0$ .

# Fully nonlinear elliptic PDE's

The ellipticity fauna

$$\mathfrak{N}[\mathbf{u}] := F(D^2 \mathbf{u}) - f = 0$$

# Fully nonlinear elliptic PDE's

## The ellipticity fauna

$$\mathfrak{N}[\mathbf{u}] := F(D^2 \mathbf{u}) - f = 0$$

- Conditionally elliptic

$$\exists \mathfrak{C} \subseteq \text{Sym}(\mathbb{R}^{d \times d}), \lambda(\cdot), \Lambda > 0 :$$

$$\lambda(\mathbf{M}) \sup_{|\boldsymbol{\xi}|=1} |\mathbf{N}\boldsymbol{\xi}| \leq F(\mathbf{M} + \mathbf{N}) - F(\mathbf{M}) \leq \Lambda \sup_{|\boldsymbol{\xi}|=1} |\mathbf{N}\boldsymbol{\xi}|$$

$$\forall \mathbf{M} \in \mathfrak{C} \subseteq \text{Sym}(\mathbb{R}^{d \times d}), \mathbf{N} \in \text{Sym}(\mathbb{R}^{d \times d}).$$

- Unconditionally elliptic if  $\mathfrak{C} = \text{Sym}(\mathbb{R}^{d \times d})$ .
- Uniformly elliptic  $\inf \lambda > 0$ .

# Characterisation of the ellipticity condition

in the smooth case

Ellipticity condition, i.e.,

$$\begin{aligned} \text{(NL-Ellip)} \quad & \lambda(\mathbf{M}) \sup_{|\boldsymbol{\xi}|=1} |\mathbf{N}\boldsymbol{\xi}| \leq F(\mathbf{M} + \mathbf{N}) - F(\mathbf{M}) \leq \Lambda \sup_{|\boldsymbol{\xi}|=1} |\mathbf{N}\boldsymbol{\xi}| \\ & \forall \mathbf{M} \in \mathfrak{C} \subseteq \text{Sym}(\mathbb{R}^{d \times d}), \mathbf{N} \in \text{Sym}(\mathbb{R}^{d \times d}). \end{aligned}$$

for some ellipticity “constants”  $\lambda(\cdot), \Lambda > 0$ .



# Characterisation of the ellipticity condition

in the smooth case

**Ellipticity condition**, i.e.,

$$\begin{aligned} \text{(NL-Ellip)} \quad & \lambda(\mathbf{M}) \sup_{|\boldsymbol{\xi}|=1} |\mathbf{N}\boldsymbol{\xi}| \leq F(\mathbf{M} + \mathbf{N}) - F(\mathbf{M}) \leq \Lambda \sup_{|\boldsymbol{\xi}|=1} |\mathbf{N}\boldsymbol{\xi}| \\ & \forall \mathbf{M} \in \mathfrak{C} \subseteq \text{Sym}(\mathbb{R}^{d \times d}), \mathbf{N} \in \text{Sym}(\mathbb{R}^{d \times d}). \end{aligned}$$

for some **ellipticity “constants”**  $\lambda(\cdot), \Lambda > 0$ . If  $F$  is differentiable then (NL-Ellip) is satisfied if and only if for each  $\mathbf{M} \in \mathfrak{C}$  there exists  $\mu > 0$  such that

$$(9.1) \quad \boldsymbol{\xi}^T F'(\mathbf{M}) \boldsymbol{\xi} \geq \mu |\boldsymbol{\xi}|^2 \quad \forall \boldsymbol{\xi} \in \mathbb{R}^d.$$

Furthermore  $\mathfrak{C} = \text{Sym}(\mathbb{R}^{d \times d})$  and  $\mu$  is independent of  $\mathbf{M}$  if and only if  $F$  is **uniformly elliptic**.

# The Monge–Ampère–Dirichlet problem

A classical fully nonlinear elliptic PDE

Boundary value problem

$$\begin{aligned} \text{(MAD)} \quad & \det D^2 u = f && \text{in } \Omega \\ & u = 0 && \text{on } \partial\Omega \end{aligned}$$

admits a unique solution in the cone of convex functions when  $f > 0$ .<sup>[Caffarelli and Cabré, 1995]</sup>

# The Monge–Ampère–Dirichlet problem

A classical fully nonlinear elliptic PDE

Boundary value problem

$$\begin{aligned} \text{(MAD)} \quad & \det D^2 u = f && \text{in } \Omega \\ & u = 0 && \text{on } \partial\Omega \end{aligned}$$

admits a unique solution in the cone of convex functions when  $f > 0$ .<sup>[Caffarelli and Cabré, 1995]</sup>

Derivative of nonlinear function  $F(\mathbf{X}) = \det \mathbf{X}$  yields

$$F'(\mathbf{X}) = \text{Cof } \mathbf{X}.$$

# The Monge–Ampère–Dirichlet problem

A classical fully nonlinear elliptic PDE

Boundary value problem

$$\begin{aligned} \text{(MAD)} \quad & \det D^2 u = f && \text{in } \Omega \\ & u = 0 && \text{on } \partial\Omega \end{aligned}$$

admits a unique solution in the cone of convex functions when  $f > 0$ . [Caffarelli and Cabré, 1995]

Derivative of nonlinear function  $F(\mathbf{X}) = \det \mathbf{X}$  yields

$$F'(\mathbf{X}) = \text{Cof } \mathbf{X}.$$

Problem elliptic if and only if

$$\xi^T \text{Cof } D^2 u \xi \geq \lambda |\xi|^2 \quad \forall \xi \in \mathbb{R}^d$$

for some  $\lambda > 0$ .

# The Monge–Ampère–Dirichlet problem

A classical fully nonlinear elliptic PDE

Boundary value problem

$$\begin{aligned} \text{(MAD)} \quad & \det D^2 u = f && \text{in } \Omega \\ & u = 0 && \text{on } \partial\Omega \end{aligned}$$

admits a unique solution in the cone of convex functions when  $f > 0$ . [Caffarelli and Cabré, 1995]

Derivative of nonlinear function  $F(\mathbf{X}) = \det \mathbf{X}$  yields

$$F'(\mathbf{X}) = \text{Cof } \mathbf{X}.$$

Problem elliptic if and only if

$$\xi^T \text{Cof } D^2 u \xi \geq \lambda |\xi|^2 \quad \forall \xi \in \mathbb{R}^d$$

for some  $\lambda > 0$ .

## Conotonic constraint

Restriction on unknown functions  $u$ : they must be **globally either convex or concave (conotonic)**.

# A simple fully nonlinear elliptic PDE

Consider problem

$$\begin{aligned}\mathfrak{N}[u] &:= \sin(\Delta u) + 2\Delta u - f = 0 \text{ in } \Omega, \\ u &= 0 \text{ on } \partial\Omega.\end{aligned}$$

Differentiating, we see that

$$D\mathfrak{N}[v]w = (\cos(\Delta v) + 2) \mathbf{I} : D^2 w = (\cos(\Delta v) + 2) \Delta w.$$

Hence problem uniformly elliptic.

# A Krylov-type cubic elliptic PDE

The problem is for  $d = 2$

$$\begin{aligned} \text{(Krylov)} \quad \mathfrak{N}[u] &:= (\partial_{11}u)^3 + (\partial_{22}u)^3 + \partial_{11}u + \partial_{22}u - f = 0 && \text{in } \Omega \\ &u = 0 && \text{on } \partial\Omega. \end{aligned}$$

Problem is uniformly elliptic since its differentiation gives:

$$F'(\mathbf{X}) = \begin{bmatrix} 3x_{22}^2 + 1 & 0 \\ 0 & 3x_{11}^2 + 1 \end{bmatrix}.$$

# Pucci's equation

Consider  $F : \text{Sym}(\mathbb{R}^{d \times d}) \rightarrow \mathbb{R}$  to be the **extremal function**

$$\text{(Pucci)} \quad F(\mathbf{N}) = \sum_{i=1}^d \alpha_i \lambda_i(\mathbf{N}) \text{ where } \lambda_i(\mathbf{N}) \text{ eigenvalues of } \mathbf{N}$$

for some given  $\alpha_1, \dots, \alpha_d \in \mathbb{R}$ .

Special case when  $d = 2$ ,  $\alpha_1 = \alpha \geq 1$  and  $\alpha_2 = 1$  yields equation

$$\text{(\mathbb{R}^2 Pucci)} \quad 0 = (\alpha + 1) \Delta u + (\alpha - 1) \left( (\Delta u)^2 - 4 \det D^2 u \right)^{1/2}.$$

The problem is unconditionally elliptic.



# Classes of fully nonlinear equations

A rough guide

See **Caffarelli and Cabré, 1995** for a more systematic classification.

- **Isaacs form**:  $\inf_{\beta} \sup_{\alpha} L_{\alpha\beta} \mathbf{u} = 0$ .

# Classes of fully nonlinear equations

## A rough guide

See **Caffarelli and Cabré, 1995** for a more systematic classification.

- **Isaacs form**:  $\inf_{\beta} \sup_{\alpha} L_{\alpha\beta} \mathbf{u} = 0$ .
  - **Bellman type**: Isaacs with only one  $\beta$  ( $\Leftrightarrow$  no inf). Related to Hamilton–Jacobi–Bellman, stochastic control and differential game theory.

# Classes of fully nonlinear equations

## A rough guide

See **Caffarelli and Cabré, 1995** for a more systematic classification.

- **Isaacs form**:  $\inf_{\beta} \sup_{\alpha} L_{\alpha\beta} \mathbf{u} = 0$ .
  - **Bellman type**: Isaacs with only one  $\beta$  ( $\Leftrightarrow$  no inf). Related to Hamilton–Jacobi–Bellman, stochastic control and differential game theory.
    - △ Isaacs form is very general: **“non-algebraic”** and harder to treat numerically. We don't, yet. [**Jensen and Smears, 2012; Lio and Ley, 2010, e.g.**].

# Classes of fully nonlinear equations

## A rough guide

See **Caffarelli and Cabré, 1995** for a more systematic classification.

- **Isaacs form**:  $\inf_{\beta} \sup_{\alpha} L_{\alpha\beta} \mathbf{u} = 0$ .
  - **Bellman type**: Isaacs with only one  $\beta$  ( $\Leftrightarrow$  no inf). Related to Hamilton–Jacobi–Bellman, stochastic control and differential game theory.
    - △ Isaacs form is very general: **“non-algebraic”** and harder to treat numerically. We don't, yet. [Jensen and Smears, 2012; Lio and Ley, 2010, e.g.]
- Hessian invariants (**algebraic**): Monge–Ampère, Pucci, Laplace (!). Subdivided into **unconditionally elliptic** (Pucci, Laplace) and **conditionally elliptic** (Monge–Ampère).

# Classes of fully nonlinear equations

## A rough guide

See **Caffarelli and Cabré, 1995** for a more systematic classification.

- **Isaacs form**:  $\inf_{\beta} \sup_{\alpha} L_{\alpha\beta} \mathbf{u} = 0$ .
  - **Bellman type**: Isaacs with only one  $\beta$  ( $\Leftrightarrow$  no inf). Related to Hamilton–Jacobi–Bellman, stochastic control and differential game theory.
    - △ Isaacs form is very general: **“non-algebraic”** and harder to treat numerically. We don't, yet. [**Jensen and Smears, 2012; Lio and Ley, 2010, e.g.**].
- Hessian invariants (**algebraic**): Monge–Ampère, Pucci, Laplace (!). Subdivided into **unconditionally elliptic** (Pucci, Laplace) and **conditionally elliptic** (Monge–Ampère).
- Other algebraic FNE's (Krylov, algebraic nonlinearities, etc.)

# Classes of fully nonlinear equations

## A rough guide

See **Caffarelli and Cabré, 1995** for a more systematic classification.

- **Isaacs form**:  $\inf_{\beta} \sup_{\alpha} L_{\alpha\beta} \mathbf{u} = 0$ .
  - **Bellman type**: Isaacs with only one  $\beta$  ( $\Leftrightarrow$  no inf). Related to Hamilton–Jacobi–Bellman, stochastic control and differential game theory.
    - △ Isaacs form is very general: **“non-algebraic”** and harder to treat numerically. We don't, yet. <sup>[Jensen and Smears, 2012; Lio and Ley, 2010, e.g.]</sup>
- Hessian invariants (**algebraic**): Monge–Ampère, Pucci, Laplace (!). Subdivided into **unconditionally elliptic** (Pucci, Laplace) and **conditionally elliptic** (Monge–Ampère).
- Other algebraic FNE's (Krylov, algebraic nonlinearities, etc.)
- Aronson equations and infinite-harmonic functions, nicely reviewed in **Barron, Evans, and Jensen, 2008**. (These aren't proper FNE's, as they are quasilinear, nevertheless, Hessian recovery applies well.)

# Monge's mass transportation problem

- consider densities  $f$  and  $g \geq 0$

# Monge's mass transportation problem

- consider densities  $f$  and  $g \geq 0$
- supports  $\text{spt } f =: \Omega$  and  $\text{spt } g =: \Upsilon$  convex



# Monge's mass transportation problem

- consider densities  $f$  and  $g \geq 0$
- supports  $\text{spt } f =: \Omega$  and  $\text{spt } g =: \Upsilon$  convex
- $f, g > 0$  on  $\text{int } \Omega$  and  $\text{int } \Upsilon$ .

# Monge's mass transportation problem

- consider densities  $f$  and  $g \geq 0$
- supports  $\text{spt } f =: \Omega$  and  $\text{spt } g =: \Upsilon$  convex
- $f, g > 0$  on  $\text{int } \Omega$  and  $\text{int } \Upsilon$ .

# Monge's mass transportation problem

- consider densities  $f$  and  $g \geq 0$
- supports  $\text{spt } f =: \Omega$  and  $\text{spt } g =: \Upsilon$  convex
- $f, g > 0$  on  $\text{int } \Omega$  and  $\text{int } \Upsilon$ .

Look for  $\psi : \Omega \rightarrow \Upsilon$  that **transports the mass density**  $f$  into the mass density  $g$ .

# Monge's mass transportation problem

- consider densities  $f$  and  $g \geq 0$
- supports  $\text{spt } f =: \Omega$  and  $\text{spt } g =: \Upsilon$  convex
- $f, g > 0$  on  $\text{int } \Omega$  and  $\text{int } \Upsilon$ .

Look for  $\psi : \Omega \rightarrow \Upsilon$  that transports the mass density  $f$  into the mass density  $g$ .

Mass conservation:

$$(28.1) \quad \int_A f(\mathbf{x}) \, d\mathbf{x} = \int_{\psi(A)} g(\mathbf{y}) \, d\mathbf{y} \quad \forall A \text{ (Borel)} \subseteq \Omega.$$

# Monge's mass transportation problem

- consider densities  $f$  and  $g \geq 0$
- supports  $\text{spt } f =: \Omega$  and  $\text{spt } g =: \Upsilon$  convex
- $f, g > 0$  on  $\text{int } \Omega$  and  $\text{int } \Upsilon$ .

Look for  $\psi : \Omega \rightarrow \Upsilon$  that **transports the mass density**  $f$  into the mass density  $g$ .

Mass conservation:

$$(29.1) \quad \int_A f(\mathbf{x}) \, d\mathbf{x} = \int_{\psi(A)} g(\mathbf{y}) \, d\mathbf{y} \quad \forall A \text{ (Borel)} \subseteq \Omega.$$

Then

$$(29.2) \quad \int_{\psi(A)} g(\mathbf{y}) \, d\mathbf{y} = \int_A g(\psi(\mathbf{x})) |\det D\psi(\mathbf{x})| \, d\mathbf{x}.$$

# Monge's mass transportation problem

- consider densities  $f$  and  $g \geq 0$
- supports  $\text{spt } f =: \Omega$  and  $\text{spt } g =: \Upsilon$  convex
- $f, g > 0$  on  $\text{int } \Omega$  and  $\text{int } \Upsilon$ .

Look for  $\psi : \Omega \rightarrow \Upsilon$  that transports the mass density  $f$  into the mass density  $g$ .

Mass conservation:

$$(30.1) \quad \int_A f(\mathbf{x}) \, d\mathbf{x} = \int_{\psi(A)} g(\mathbf{y}) \, d\mathbf{y} \quad \forall A \text{ (Borel)} \subseteq \Omega.$$

Then

$$(30.2) \quad \int_{\psi(A)} g(\mathbf{y}) \, d\mathbf{y} = \int_A g(\psi(\mathbf{x})) |\det D\psi(\mathbf{x})| \, d\mathbf{x}.$$

Hence

$$(30.3) \quad g(\psi(\mathbf{x})) |\det D\psi(\mathbf{x})| = f(\mathbf{x}) \quad \forall \mathbf{x} \in \Omega.$$

# From Monge to Monge–Ampère

Following Caffarelli, 1990a; Caffarelli, 1990b,c; Caffarelli and Cabré, 1995  
Evans, 2001 Urbas, 1997 under convexity and regularity assumptions, the  
Monge–Ampère equation

$$\det D^2 u(\mathbf{x}) = k(\mathbf{x}, u(\mathbf{x}), \nabla u(\mathbf{x}))$$

coupled to the second boundary condition **second boundary condition**

$$(31.1) \quad \nabla u(\Omega) = \gamma,$$

provides a solution to the Monge problem and the right-hand side

$$(31.2) \quad \frac{f(\mathbf{x})}{g(\nabla u(\mathbf{x}))}$$

# Finite difference approaches

- ① Earliest known provided approximations of the Monge–Ampère (and other equations) by **Oliker and Prussner, 1988**.
- ② **Kuo and Trudinger, 1992** gave mostly theoretical work introduced the concept of **wide stencils** and proving convergence for wide enough stencils.
- ③ **Benamou and Brenier, 2000** proposed an approach based on the **Brenier-solution** concept related to fluid-dynamics and mass-transportation.
- ④ **Oberman, 2008** introduced more practically effective work working out the details, providing a bound on the wide stencil's width. See also **Froese, 2011** and **Benamou, Froese, and Oberman, 2012** for second boundary conditions.



# Galerkin (mainly finite element) methods I

- Dean and Glowinski, 2006 (and earlier work) introduced a FE least square method to solve Monge–Ampère equation.

# Galerkin (mainly finite element) methods II

- **Awanou, 2011** uses a **pseudo time** [sic] approach.
- **Jensen and Smears, 2012** provide and analyze a FEM for a special class of **Hamilton–Jacobi–Bellman** equation. Further work in **Smears and Süli, 2013, 2014** for a DGFEM approach.

# A fixed-point solution

Nonlinear PDE

$$\mathfrak{N}[u] := F(D^2 u) - f = 0$$

can be rewritten as follows

$$\mathfrak{N}[u] = \left[ \int_0^1 F'(t D^2 u) dt \right] : D^2 u + F(0) - f = 0.$$

Define

$$\begin{aligned} \mathbf{N}(D^2 u) &:= \int_0^1 F'(t D^2 u) dt, \\ g &:= f - F(0), \end{aligned}$$

then if  $u$  solves (FNE), it also solves

$$\mathbf{N}(D^2 u) : D^2 u = g.$$

Fixed point iteration: given  $u^0$  find

$$\mathbf{N}(D^2 u^n) : D^2 u^{n+1} = g, \text{ for } n = 1, 2, \dots$$

# Crucial remark

Note that solving

$$\mathbf{N}(D^2 \mathbf{u}^n):D^2 \mathbf{u}^{n+1} = g$$

involves a **linear elliptic equation in non-divergence form**.

## Big fat note

Standard variational FEM's do not apply.

# Newton's method

Given an initial guess  $\mathbf{u}^0$ , let

$$D \mathfrak{N}[\mathbf{u}^n] (\mathbf{u}^{n+1} - \mathbf{u}^n) = -\mathfrak{N}[\mathbf{u}^n], \text{ for } n = 0, 1, 2, \dots,$$

where

$$D \mathfrak{N}[\mathbf{u}] \mathbf{v} = F'(D^2 \mathbf{u}) : D^2 \mathbf{v}.$$

i.e.,

$$F'(D^2 \mathbf{u}^n) : D^2 (\mathbf{u}^{n+1} - \mathbf{u}^n) = f - F(D^2 \mathbf{u}^n).$$

**Big fat note (repeated)**

Equation in nondivergence form, standard FEM's will not apply.

# The need for Hessian recovery

Detailed in [Lakkis and Pryer, 2013](#)

Fixed point iteration

$$\mathbf{N}(D^2 \mathbf{u}^n) : D^2 \mathbf{u}^{n+1} = \mathbf{g}$$

and Newton's iteration

$$F'(D^2 \mathbf{u}^n) : D^2 (\mathbf{u}^{n+1} - \mathbf{u}^n) = \mathbf{f} - F(D^2 \mathbf{u}^n).$$

besides being nonvariational, like fixed-point, **requires the suitable approximation of a Hessian's function.**

## Big fat note (a variation)

Hence the use of the **recovered Hessian** introduced by [Lakkis and Pryer, 2011](#).

Introduce Galerkin **finite element spaces**

$$\mathbb{V}_h := \left\{ \Phi \in H^1(\Omega) : \Phi|_K \in \mathbb{P}^p \forall K \in \mathcal{T} \text{ and } \Phi \in C^0(\Omega) \right\},$$
$$\mathbb{V}_0 := \mathbb{V} \cap H_0^1(\Omega),$$


**Unbalanced mixed problem:**

Find  $(\mathbf{U}, \mathbf{H}) \in \mathbb{V}_0 \times \mathbb{V}^{d \times d}$  satisfying

$$\langle \mathbf{H}, \Phi \rangle + \int_{\Omega} \nabla \mathbf{U} \otimes \nabla \Phi - \int_{\partial\Omega} \nabla \mathbf{U} \otimes \mathbf{n} \Phi = 0$$
$$\langle \mathbf{A}:\mathbf{H}, \Psi \rangle = \langle f, \Psi \rangle \quad \forall (\Phi, \Psi) \in \mathbb{V} \times \mathbb{V}_0.$$

# A (sometimes) commutative diagram

discretization are often possible (e.g., when the nonlinearity is algebraic in the Hessian):





# Convergence analysis

Available for the **linear nondivergence case** so far

A priori estimates for the error

$$\left\| \mathbf{A} : (\mathbf{D}^2 \mathbf{u} - \mathbf{H}[\mathbf{u}_h]) \right\|_{\mathbf{H}^{-1}(\Omega)}.$$

A posteriori error estimate for the error

$$\|\mathbf{u} - \mathbf{u}_h\|_{L_2(\Omega)}^2 \leq \sum_{K \in \mathfrak{T}} \left( h_K^2 \|\mathbf{f} - \mathbf{A} : \mathbf{D}^2 \mathbf{u}\|_{L_2(K)}^2 + h_K \|\mathbf{A} : \llbracket \nabla \mathbf{u} \otimes \rrbracket\|_{L_2(\partial K)}^2 \right)$$

where the **tensor jump** of a field  $\mathbf{v}$  across an edge  $E = \bar{K} \cap \bar{K}'$  is given by



$$\llbracket \mathbf{v} \otimes \rrbracket_E := \lim_{\epsilon \rightarrow 0} (\mathbf{v}(\mathbf{x} + \epsilon \mathbf{n}_K) \otimes \mathbf{n}_K + \mathbf{v}(\mathbf{x} - \epsilon \mathbf{n}_{K'}) \otimes \mathbf{n}_{K'})$$

# A nonlinear function of $\Delta u$

$$\mathfrak{N}[u] := \sin(\Delta u) + 2\Delta u - f = 0 \text{ in } \Omega,$$



$$u = 0 \text{ on } \partial\Omega.$$

P1 elements (left) and P2 elements (right)



# Krylov's equation



## P1 elements (left) and P2 elements (right)



# Pucci's equation

$$0 = (\alpha + 1) \Delta u + (\alpha - 1) \left( (\Delta u)^2 - 4 \det D^2 u \right)^{1/2}.$$

$\mathbb{P}^2, \alpha = 2$  (left) and  $\mathbb{P}^2, \alpha = 5$  (right)





# Some MAD stuff

reminder: MAD = Monge–Ampère–Dirichlet

FE-convexity check inspired from [Aguilera and Morin, 2009](#).

Exact solution and EOC's for  $\mathbb{P}^2$  elements (suboptimal for  $\mathbb{P}^1$ )




# Some MAD stuff

reminder: MAD = Monge–Ampère–Dirichlet

FE-convexity check inspired from [Aguilera and Morin, 2009](#).



principal minor and determinant instances



# Nonclassical solutions

Viscosity or Alexandrov



Singular solution  $u(\mathbf{x}) = |\mathbf{x}|^{2\alpha}$



# Nonclassical solutions

Viscosity or Alexandrov

More singular,  $\alpha = 0.6$ ,  $\alpha = 0.55, \dots$







# Adaptive approximation of nonclassical solutions

Viscosity or Alexandrov

Singular solution  $u(\mathbf{x}) = |\mathbf{x}|^{1.1}$  (empirical ZZ-estimators)



# Conclusions and outlook

- Obtained and tested a practical and “easy” Netwon scheme based on **nonvariational FEM (NVFEM) via Hessian recovery**.

# Conclusions and outlook

- Obtained and tested a practical and “easy” Newton scheme based on **nonvariational FEM (NVFEM) via Hessian recovery**.
- Convergence rates optimal in all examples.

# Conclusions and outlook

- Obtained and tested a practical and “easy” Netwon scheme based on **nonvariational FEM (NVFEM) via Hessian recovery**.
- Convergence rates optimal in all examples.
- A posteriori error estimates for very weak norms in the linear problem, provide an elementary way to do **adaptivity**.

# Conclusions and outlook

- Obtained and tested a practical and “easy” Newton scheme based on **nonvariational FEM (NVFEM) via Hessian recovery**.
- Convergence rates optimal in all examples.
- A posteriori error estimates for very weak norms in the linear problem, provide an elementary way to do **adaptivity**.
- **In progress**: prove a priori convergence for stronger norms in linear problems.

# Conclusions and outlook

- Obtained and tested a practical and “easy” Netwon scheme based on **nonvariational FEM (NVFEM) via Hessian recovery**.
- Convergence rates optimal in all examples.
- A posteriori error estimates for very weak norms in the linear problem, provide an elementary way to do **adaptivity**.
- **In progress**: prove a priori convergence for stronger norms in linear problems.
- **In progress**: embed **second boundary condition** ( $\nabla u(\Omega) = \Upsilon$  with prescribed  $\Upsilon$ ). (This was achieved for wide-stencils but on **structured grids** by **Benamou, Froese, and Oberman, 2012**.)

# Conclusions and outlook

- Obtained and tested a practical and “easy” Netwon scheme based on **nonvariational FEM (NVFEM) via Hessian recovery**.
- Convergence rates optimal in all examples.
- A posteriori error estimates for very weak norms in the linear problem, provide an elementary way to do **adaptivity**.
- **In progress**: prove a priori convergence for stronger norms in linear problems.
- **In progress**: embed **second boundary condition** ( $\nabla u(\Omega) = \Upsilon$  with prescribed  $\Upsilon$ ). (This was achieved for wide-stencils but on **structured grids** by **Benamou, Froese, and Oberman, 2012**.)
- **Open problem**: prove conservation of **conotonicity** for MAD/MAS.

# Conclusions and outlook

- Obtained and tested a practical and “easy” Netwon scheme based on **nonvariational FEM (NVFEM) via Hessian recovery**.
- Convergence rates optimal in all examples.
- A posteriori error estimates for very weak norms in the linear problem, provide an elementary way to do **adaptivity**.
- **In progress**: prove a priori convergence for stronger norms in linear problems.
- **In progress**: embed **second boundary condition** ( $\nabla u(\Omega) = \Upsilon$  with prescribed  $\Upsilon$ ). (This was achieved for wide-stencils but on **structured grids** by Benamou, Froese, and Oberman, 2012.)
- **Open problem**: prove conservation of **conotonicity** for MAD/MAS.
- **Open problem**: **apriori and aposteriori analysis** for nonlinear problem.



# References I

- Aguilera, Néstor E. and Pedro Morin (2009). “On convex functions and the finite element method”. In: **SIAM J. Numer. Anal.** 47.4, pp. 3139–3157. ISSN: 0036-1429. DOI: 10.1137/080720917. URL: <http://dx.doi.org/10.1137/080720917>.
- Awanou, Gerard (2011). **Pseudo time continuation and time marching methods for Monge–Ampère type equations**. online preprint. URL: <http://www.math.niu.edu/~awanou/MongePseudo05.pdf>.
- Barron, E. N., L. C. Evans, and R. Jensen (2008). “The infinity Laplacian, Aronsson’s equation and their generalizations”. In: **Trans. Amer. Math. Soc.** 360.1, pp. 77–101. ISSN: 0002-9947. DOI: 10.1090/S0002-9947-07-04338-3. URL: <http://dx.doi.org/10.1090/S0002-9947-07-04338-3>.

## References II

- Benamou, Jean-David and Yann Brenier (2000). “A computational fluid mechanics solution to the Monge-Kantorovich mass transfer problem”. In: **Numer. Math.** 84.3, pp. 375–393. ISSN: 0029-599X. DOI: 10.1007/s002110050002. URL: <http://dx.doi.org/10.1007/s002110050002>.
- Benamou, Jean-David, Brittany D. Froese, and Adam M. Oberman (Aug. 2012). **A viscosity solution approach to the Monge-Ampere formulation of the Optimal Transportation Problem**. Tech. rep. eprint: 1208.4873. URL: <http://arxiv.org/abs/1208.4873>.
- Böhmer, Klaus (2010). **Numerical methods for nonlinear elliptic differential equations**. Numerical Mathematics and Scientific Computation. A synopsis. Oxford: Oxford University Press, pp. xxviii–746. ISBN: 978-0-19-957704-0.
- Brenner, Susanne C. et al. (2011). “ $C^1$  penalty methods for the fully nonlinear Monge-Ampère equation”. In: **Math. Comp.** 80, pp. 1979–1995. DOI: 10.1090/S0025-5718-2011-02487-7.

## References III

- Caffarelli, L. A. (1990a). “A localization property of viscosity solutions to the Monge-Ampère equation and their strict convexity”. In: **Ann. of Math. (2)** 131.1, pp. 129–134. ISSN: 0003-486X. DOI: 10.2307/1971509. URL: <http://dx.doi.org/10.2307/1971509>.
- Caffarelli, Luis A. (1990b). “Interior regularity of solutions to Monge-Ampère equations”. In: **Harmonic analysis and partial differential equations (Boca Raton, FL, 1988)**. Vol. 107. Contemp. Math. Amer. Math. Soc., Providence, RI, pp. 13–17. DOI: 10.1090/conm/107/1066467. URL: <http://dx.doi.org/10.1090/conm/107/1066467>.
- (1990c). “Interior  $W^{2,p}$  estimates for solutions of the Monge-Ampère equation”. In: **Ann. of Math. (2)** 131.1, pp. 135–150. ISSN: 0003-486X. DOI: 10.2307/1971510. URL: <http://dx.doi.org/10.2307/1971510>.

## References IV

- Caffarelli, Luis A. and Xavier Cabré (1995). **Fully nonlinear elliptic equations**. Vol. 43. American Mathematical Society Colloquium Publications. Providence, RI: American Mathematical Society, pp. vi+104. ISBN: 0-8218-0437-5.
- Davydov, Oleg and Abid Saeed (2013). “Numerical solution of fully nonlinear elliptic equations by Böhmer’s method”. In: **J. Comput. Appl. Math.** 254, pp. 43–54. ISSN: 0377-0427. DOI: [10.1016/j.cam.2013.03.009](https://doi.org/10.1016/j.cam.2013.03.009). URL: <http://dx.doi.org/10.1016/j.cam.2013.03.009>.
- Dean, Edward J. and Roland Glowinski (2006). “Numerical methods for fully nonlinear elliptic equations of the Monge-Ampère type”. In: **Comput. Methods Appl. Mech. Engrg.** 195.13-16, pp. 1344–1386. ISSN: 0045-7825. DOI: [10.1016/j.cma.2005.05.023](https://doi.org/10.1016/j.cma.2005.05.023). URL: <http://dx.doi.org/10.1016/j.cma.2005.05.023>.

- Evans, Lawrence C. (2001). **Partial Differential Equations and Monge–Kantorovich Mass Transfer**. online lecture notes. University of California, Berkley, CA USA. URL: <http://math.berkeley.edu/~evans/Monge-Kantorovich.survey.pdf>.
- Feng, Xiaobing and Michael Neilan (2009). “Vanishing moment method and moment solutions for fully nonlinear second order partial differential equations”. In: **J. Sci. Comput.** 38.1, pp. 74–98. ISSN: 0885-7474. DOI: 10.1007/s10915-008-9221-9. URL: <http://dx.doi.org/10.1007/s10915-008-9221-9>.
- Froese, Brittany D. (Jan. 2011). **A numerical method for the elliptic Monge-Ampère equation with transport boundary conditions**. Tech. rep. eprint: 1101.4981v1. URL: <http://arxiv.org/abs/1101.4981v1>.
- Jensen, Max and Iain Smears (Jan. 2012). **Finite Element Methods with Artificial Diffusion for Hamilton-Jacobi-Bellman Equations**. Tech. rep. eprint: 1201.3581v2. URL: <http://arxiv.org/abs/1201.3581v2>.

## References VI

- Kuo, Hung Ju and Neil S. Trudinger (1992). “Discrete methods for fully nonlinear elliptic equations”. In: **SIAM J. Numer. Anal.** 29.1, pp. 123–135. ISSN: 0036-1429. DOI: 10.1137/0729008. URL: <http://dx.doi.org/10.1137/0729008>.
- Lakkis, O. and T. Pryer (2013). “A finite element method for nonlinear elliptic problems”. In: **SIAM Journal on Scientific Computing** 35.4, A2025–A2045. DOI: 10.1137/120887655. eprint: <http://epubs.siam.org/doi/pdf/10.1137/120887655>. URL: <http://epubs.siam.org/doi/abs/10.1137/120887655>.
- Lakkis, Omar and Tristan Pryer (2011). “A finite element method for second order nonvariational elliptic problems”. In: **SIAM J. Sci. Comput.** 33.2, pp. 786–801. ISSN: 1064-8275. DOI: 10.1137/100787672. URL: <http://dx.doi.org/10.1137/100787672>.

## References VII

- Lio, Francesca Da and Olivier Ley (Feb. 2010). **Uniqueness Results for Second Order Bellman-Isaacs Equations under Quadratic Growth Assumptions and Applications**. Tech. rep. eprint: 1002.2373v1. URL: <http://arxiv.org/abs/1002.2373v1>.
- Neilan, Michael J. (2012). **Finite element methods for fully nonlinear second order PDEs based on the discrete Hessian**. online preprint. University of Pittsburgh. URL: [https://dl.dropbox.com/u/48847074/Publications/MA\\_Mixed.pdf](https://dl.dropbox.com/u/48847074/Publications/MA_Mixed.pdf).
- Oberman, Adam M. (2008). "Wide stencil finite difference schemes for the elliptic Monge-Ampère equation and functions of the eigenvalues of the Hessian". In: **Discrete Contin. Dyn. Syst. Ser. B** 10.1, pp. 221–238. ISSN: 1531-3492. DOI: 10.3934/dcdsb.2008.10.221. URL: <http://dx.doi.org/10.3934/dcdsb.2008.10.221>.
- Oliker, V. I. and L. D. Prussner (1988). "On the numerical solution of the equation  $\partial_x^2 z \partial_y^2 z - \partial^2 z_{x,y}$  and its discretizations, I". In: **Numerische Mathematik** 54.3, pp. 271–293.

## References VIII

- Smears, Iain and Endre Süli (2013). “Discontinuous Galerkin finite element approximation of nondivergence form elliptic equations with Cordès coefficients”. In: **SIAM J. Numer. Anal.** 51.4, pp. 2088–2106. ISSN: 0036-1429. DOI: 10.1137/120899613. URL: <http://dx.doi.org/10.1137/120899613>.
- (2014). “Discontinuous Galerkin finite element approximation of Hamilton-Jacobi-Bellman equations with Cordes coefficients”. In: **SIAM J. Numer. Anal.** 52.2, pp. 993–1016. ISSN: 0036-1429. DOI: 10.1137/130909536. URL: <http://dx.doi.org/10.1137/130909536>.
- Urbas, John (1997). “On the second boundary value problem for equations of Monge-Ampère type”. In: **J. Reine Angew. Math.** 487, pp. 115–124. ISSN: 0075-4102. DOI: 10.1515/crll.1997.487.115. URL: <http://dx.doi.org/10.1515/crll.1997.487.115>.