




Models of hyperbolic geometry

model	Poincaré disk	Upper half-plane	Klein disk	two-sheet hyperboloid
\mathbb{H}^2	$\{z \in \mathbb{C} \mid z < 1\}$	$\{z \in \mathbb{C} \mid \text{Im}z > 0\}$	$\{z \in \mathbb{C} \mid z < 1\}$	$\{v \in \mathbb{R}^{2,1} \mid \langle v, v \rangle = -1, v_3 > 0\}$ where $\langle v, u \rangle = v_1u_1 + v_2u_2 - v_3u_3$
$\partial\mathbb{H}^2$ (absolute)	$\{z \in \mathbb{C} \mid z = 1\}$	$\{z \in \mathbb{C} \mid \text{Im}z = 0\}$	$\{z \in \mathbb{C} \mid z = 1\}$	$\{v \in \mathbb{R}^{2,1} \mid \langle v, v \rangle = 0, v_3 > 0\}$ $v \sim \lambda v$
lines				$\{v \mid \langle v, a \rangle = 0\}$ where $\langle a, a \rangle > 0$
distance	$d(A, B) = \ln[A, B, X, Y] $ X, Y = the “endpoints” of the line AB		$d(A, B) = \frac{1}{2} \ln[A, B, X, Y] $	$d(A, B) = \frac{1}{2} \ln[A, B, X, Y] $ cross-ratio of four lines*
formula	$\cosh d(u, v) = 1 + \frac{ u-v ^2}{2\text{Im}(u)\text{Im}(v)}$			$Q = \left \frac{\langle u, v \rangle^2}{\langle u, u \rangle \langle v, v \rangle} \right $ if $\langle u, u \rangle < 0, \langle v, v \rangle < 0$ $Q = \cosh^2 d(pt, pt)$ if $\langle u, u \rangle < 0, \langle v, v \rangle > 0$ $Q = \sinh^2 d(pt, line)$ if $\langle u, u \rangle > 0, \langle v, v \rangle > 0$ $Q < 1$, intersecting lines $Q = \cos^2 \alpha$ $Q = 1$, parallel lines $Q > 1$, ultraparallel lines $Q = \cosh^2 d(line, line)$
isometries**	Möbius transformations		Projective tr	Linear transformations of $\mathbb{R}^{2,1}$
orientation-preserving isometries		$\frac{az+b}{cz+d}$ $a, b, c, d \in \mathbb{R}, ad - bc = 1$		
orientation-reversing isometries		$\frac{a\bar{z}+b}{c\bar{z}+d}$ $a, b, c, d \in \mathbb{R}, ad - bc = -1$		
reflections	Euclidean inversions or reflections			$r_a(v) = v - 2\frac{\langle v, a \rangle}{\langle a, a \rangle}a$
circles	Euclidean circles		ellipses	plane sections of the hyperboloid
angles	angles=Euclidean angles		distorted angles good for right angles***	

* Cross-ratio of four lines lying in one plane and passing through one point is the cross-ratio of four points at which these lines are intersected by an arbitrary line l (it does not depend on l !).

**We only list the type of the transformations not specifying that they preserve the model.

*** See the next page.

*****Right angles in the Klein model.**

Let l be a hyperbolic line.

Let \bar{l} be a Euclidean line containing the segment which represents l in the Klein model.

Let $X_1(l)$ and $X_2(l)$ be the endpoints of l (intersections of \bar{l} with the unit circle).

Let $t_1(l)$ and $t_2(l)$ be tangent lines to the unit circle at the points $X_1(l)$ and $X_2(l)$.

Let $T(l) = t_1(l) \cap t_2(l)$ (if $t_1 \parallel t_2$, i.e. l is represented by a diameter, then $T(l)$ is a point at infinity).

Thm. l' is orthogonal to l if and only if $T(l) \in l'$.

In particular, if l is represented by a **diameter**, then $l' \perp l$ if and only if $\bar{l}' \perp \bar{l}$ (in Euclidean sense).

