

HOMOLOGY OF MODULI SPACES OF LINKAGES IN HIGH-DIMENSIONAL EUCLIDEAN SPACE

DIRK SCHÜTZ

ABSTRACT. We study the topology of moduli spaces of closed linkages in \mathbb{R}^d depending on a length vector $\ell \in \mathbb{R}^n$. In particular, we use equivariant Morse theory to obtain information on the homology groups of these spaces, which works best for odd d . In the case $d = 5$ we calculate the Poincaré polynomial in terms of combinatorial information encoded in the length vector.

1. INTRODUCTION

In this paper, we consider polygons, or linkages, with fixed side lengths in Euclidean space \mathbb{R}^d . The topology of the corresponding moduli spaces $\mathcal{M}_d(\ell)$, see Section 2 for precise definitions, has been studied extensively in the cases $d = 2$ and 3. In particular, a lot of information on the homology and cohomology of these spaces have been obtained, see for example [6, 9, 15, 18, 23]. Furthermore, cohomology can be used to show that the topological type of $\mathcal{M}_d(\ell)$ for $d = 2$ and 3 is determined by the length vector $\ell \in \mathbb{R}^n$, [4, 22].

A lot less is known for $d > 3$. Kamiyama [13] has obtained a formula for the Euler characteristic of $\mathcal{M}_4(\ell)$ in the equilateral case, that is, when $\ell = (1, 1, \dots, 1) \in \mathbb{R}^n$. Schoenberg [21] shows that $\mathcal{M}_d(\ell)$ is homeomorphic to a disc if $d \geq n$, which implies that it is homeomorphic to a sphere for $d = n - 1$.

Another class of examples which has been extensively studied is given by $\ell = (1, \dots, 1, n - 2) \in \mathbb{R}^n$, as in this case $\mathcal{M}_d(\ell)$ coincides with the shape space Σ_{d-1}^{n-1} , which is thoroughly examined in [17]. In particular, the homology groups of these spaces are completely known [17]. It follows from their calculations that for $4 \leq d < n - 1$ the space Σ_{d-1}^{n-1} is not a manifold.

Our method to study $\mathcal{M}_d(\ell)$ is through equivariant Morse theory. We use the fact that $\mathcal{M}_d(\ell) = \mathcal{C}_d(\ell)/SO(d - 1)$, where $\mathcal{C}_d(\ell)$ is the so-called chain space, and construct a $SO(d - 1)$ -invariant Morse-Bott function on $\mathcal{C}_d(\ell)$. By analyzing the critical manifolds we obtain information on the homology of $\mathcal{M}_d(\ell)$. Our method works best for odd d and rational coefficients, as the spectral sequence arising from the filtration given by the Morse-Bott function collapses. For even d we can still obtain useful information on the topology of $\mathcal{M}_d(\ell)$.

Theorem 1.1. *Let $\ell \in \mathbb{R}^n$ be a generic length vector such that $\mathcal{M}_d(\ell) \neq \emptyset$ for $d \geq 2$.*

- (1) *For $d \geq 3$ the space $\mathcal{M}_d(\ell)$ is $((d - 1)(d - 2)/2 + d - 3)$ -connected.*
- (2) *For $n \geq 3$, $\mathcal{M}_{n-1}(\ell)$ is homeomorphic to the sphere of dimension $n(n - 3)/2$.*
- (3) *For $4 \leq d \leq n - 2$, $\mathcal{M}_d(\ell)$ is not homotopy equivalent to a closed manifold.*

Our Morse-theoretic methods imply homotopy equivalence in (2), but using the result of Schoenberg [21] the homeomorphism can be obtained directly.

Part (3) is obtained by showing that $\mathcal{M}_d(\ell)$ does not satisfy Poincaré duality. Note that for $d = 2, 3$ the space $\mathcal{M}_d(\ell)$ is a smooth closed manifold.

As we have mentioned before, our homology calculations work best for odd d . To obtain simple formulas for the Poincaré polynomial, we stick to the case $d = 5$. Define

$$\begin{aligned} Q_{2m}(t) &= \frac{(t^{m+1} - 1)^2}{(t - 1)^2} \\ Q_{2m+1}(t) &= \frac{(t^{m+2} - 1)(t^{m+1} - 1)}{(t - 1)^2} \end{aligned}$$

for all $m \geq 0$.

Theorem 1.2. *Let $\ell \in \mathbb{R}^n$ be a generic length vector. Let $m \geq 3$ be such that $n = 2m - 1$ or $n = 2m$. Then there exist non-negative integers $c_i(\ell)$ depending only on ℓ such that the Poincaré polynomial of $\mathcal{M}_5(\ell)$ is*

$$P_5^\ell(t) = 1 + t^9 \cdot \sum_{i=0}^{m-2} (c_i(\ell) (Q_{n-6-i}(t^4) - Q_{i-4}(t^4))).$$

The exact form of $c_i(\ell)$ can be seen in Theorem 10.4.

The case $d = 5$ also contains interesting geometry closely related to the case $d = 3$. Indeed, $\mathcal{M}_3(\ell)$ carries extra symplectic and Kähler structures, which have been studied in detail in [9, 16, 18]. In particular, in [16] a complex analytic equivalence between $\mathcal{M}_3(\ell)$ and a weighted quotient of $(S^2)^n$ by $PSL(2, \mathbb{C})$ is established. Foth and Lozano [7] obtain an analogous statement for $\mathcal{M}_5(\ell)$ and a weighted quotient of $(S^4)^n$ by $PSL(2, \mathbb{H})$. They also generalize the Gel'fand-MacPherson correspondence to the quaternion context and realize $\mathcal{M}_5(\ell)$ as a quotient of a subspace in a quaternion Grassmannian.

It can be easily read off Theorem 1.2 that the reduced rational homology of $\mathcal{M}_5(\ell)$ starts in degree 9 and is limited to odd degrees. In particular the rational cohomology ring structure is trivial. Given that the cohomology ring structure is instrumental in distinguishing topological types for $d = 2$ and 3, one would hope for more algebraic information also in the cases $d \geq 4$. A suitable setting for this appears to be intersection homology, which we plan to examine in a future project.

The paper is organized as follows. Section 2 collects some basic properties of linkage spaces and Section 3 introduces the equivariant Morse-Bott function. While our interest is mainly for large values of d , we also obtain results on the existence of perfect Morse functions in the case $d = 2$ (Section 4) and $d = 3$ (Section 5). A cell decomposition for $\mathcal{M}_d(\ell)$ based on the Morse-Bott function is obtained in Sections 6 and 7, which is used to prove Theorem 1.1. Local homology calculations are done in Sections 8 and 9, which culminate in the proof of Theorem 1.2 in Section 10. We also obtain some Euler characteristic results for even d in Section 11. There are two appendices, one showing the equivalence of the shape space with a certain linkage space, and one deals with basic properties of the polynomials $Q_n(t)$.

2. BASIC DEFINITIONS AND PROPERTIES OF LINKAGE SPACES

Let d, n be positive integers and $\ell = (\ell_1, \dots, \ell_n)$ satisfy $0 < \ell_i$ for all $i = 1, \dots, n$. We call ℓ a *length vector*. The *moduli space of ℓ* in \mathbb{R}^d consists of all closed linkages with lengths ℓ up to rotations and translations. We can describe this space as

$$\mathcal{M}_d(\ell) = \left\{ (z_1, \dots, z_n) \in (S^{d-1})^n \mid \sum_{i=1}^n \ell_i z_i = 0 \right\} / SO(d)$$

where $SO(d)$ acts diagonally on $(S^{d-1})^n$. We also denote the *space of chains* of ℓ as

$$\mathcal{C}_d(\ell) = \left\{ (z_1, \dots, z_n) \in (S^{d-1})^{n-1} \mid \sum_{i=1}^{n-1} \ell_i z_i = (-\ell_n, 0, \dots, 0) \right\}$$

If we let $SO(d-1)$ act on S^{d-1} by fixing the first coordinate, we get an $SO(d-1)$ -action on $\mathcal{C}_d(\ell)$ such that

$$\mathcal{M}_d(\ell) = \mathcal{C}_d(\ell) / SO(d-1).$$

It is clear that permuting the coordinates of ℓ does not change the homeomorphism type of $\mathcal{M}_d(\ell)$. However this is not true for $\mathcal{C}_d(\ell)$. In the cases $d = 1, 2, 4, 8$ one can use the multiplication structure of S^{d-1} to construct a homeomorphism $\mathcal{C}_d(\ell) \cong \mathcal{C}_d(\sigma\ell)$ for any permutation σ , but for other values of d these spaces are usually not homeomorphic, compare [5, Rm.2.2].

Definition 2.1. Let ℓ be a length vector. A subset $J \subset \{1, \dots, n\}$ is called *ℓ -short*, if

$$\sum_{j \in J} \ell_j < \sum_{i \notin J} \ell_i.$$

It is called *ℓ -long*, if the complement is ℓ -short, and *ℓ -median*, if it is neither ℓ -short nor ℓ -long. The length vector is called *generic*, if there are no ℓ -median subsets.

We also write

$$\ell_J = \sum_{j \in J} \ell_j.$$

For $m \in \{1, \dots, n\}$ the length vector is called *m -dominated*, if $\ell_m \geq \ell_i$ for all $i = 1, \dots, n$.

If the length vector is generic, there do not exist collinear configurations, that is, points $[z_1, \dots, z_n] \in \mathcal{M}_d(\ell)$ for which all $z_i \in \{\pm x\}$ for some $x \in S^{d-1}$. Notice that generic is equivalent to $\mathcal{M}_1(\ell) = \emptyset$.

In the case that ℓ is generic, it is easy to see that $\mathcal{C}_d(\ell)$ is a closed manifold of dimension $(n-2)(d-1) - 1$. In the case that $d = 2$ or $d = 3$, we then get that $SO(d-1)$ acts freely on $\mathcal{C}_d(\ell)$, and $\mathcal{M}_d(\ell)$ is also a closed manifold of dimension $(d-1)(n-3)$. For $d \geq 4$, the action is no longer free, and we will see that generally $\mathcal{M}_d(\ell)$ is not a manifold. we denote the dimension of $\mathcal{M}_d(\ell)$ by \mathbf{d}_d^n . If $d \leq n-1$, there exists a free

Definition 2.2. Let $\ell \in \mathbb{R}^n$ be an m -dominated generic length vector. For $k \in \{0, \dots, n-3\}$ we write

$$\mathcal{S}_k^m(\ell) = \{J \subset \{1, \dots, n\} \mid m \in J, |J| = k+1, J \text{ is } \ell\text{-short}\}.$$

and

$$a_k(\ell) = |\mathcal{S}_k^m(\ell)|.$$

So the union $\mathcal{S}_*^m(\ell)$ of $\mathcal{S}_k^m(\ell)$ over all $k = 0, \dots, n-3$ contains all short subsets $J \subset \{1, \dots, n\}$ which include m . It is worth pointing out that $\mathcal{S}_*^m(\ell)$ is an abstract simplicial complex with 0-simplices given by $\mathcal{S}_1^m(\ell)$.

Note that a length vector can be m -dominated by more than one $m \in \{1, \dots, n\}$. The numbers $a_k(\ell)$ however do not depend on this. We have $a_0(\ell) \leq 1$, and for a generic length vector it is easy to see that $\mathcal{M}_d(\ell) \neq \emptyset$ for $d \geq 2$ if and only if $a_0(\ell) = 1$.

If $J \subset \{1, \dots, n\}$, we define the hyperplane

$$H_J = \left\{ (x_1, \dots, x_n) \in \mathbb{R}^n \mid \sum_{j \in J} x_j = \sum_{j \notin J} x_j \right\}$$

and let

$$\mathcal{H} = \mathbb{R}_{>0}^n - \bigcup_{J \subset \{1, \dots, n\}} H_J,$$

where $\mathbb{R}_{>0}^n = \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_i > 0\}$. Then \mathcal{H} has finitely many components, which we call *chambers*. It is clear that a length vector ℓ is generic if and only if $\ell \in \mathcal{H}$.

It is shown in [8] that if ℓ and ℓ' are in the same chamber, then $\mathcal{C}_d(\ell)$ and $\mathcal{C}_d(\ell')$ are $O(d-1)$ -equivariantly diffeomorphic. In particular, $\mathcal{M}_d(\ell)$ and $\mathcal{M}_d(\ell')$ are homeomorphic.

It is easy to see that two m -dominated generic length vectors ℓ, ℓ' are in the same chamber if and only if $\mathcal{S}_*^m(\ell) = \mathcal{S}_*^m(\ell')$.

Definition 2.3. Let $z = (z_1, \dots, z_{n-1}) \in \mathcal{C}_d(\ell)$. The *rank* of z is the maximal number of linearly independent vectors $z_1, \dots, z_{n-1} \in \mathbb{R}^d$. Note that the rank remains the same under the $SO(d-1)$ action, and we can define the *rank* of $z = [z_1, \dots, z_n] \in \mathcal{M}_d(\ell)$ also as the maximal number of linearly independent vectors $z_1, \dots, z_n \in \mathbb{R}^d$.

The natural inclusion $i : \mathcal{C}_{d-1}(\ell) \rightarrow \mathcal{C}_d(\ell)$ induces a natural map

$$\varphi : \mathcal{M}_{d-1}(\ell) \rightarrow \mathcal{M}_d(\ell).$$

This map need not be injective, in fact, if $\ell = (1, 1, 1)$, it is clear that $\mathcal{M}_2(\ell) = S^0$ and $\mathcal{M}_d(\ell) = \{*\}$ for $d \geq 3$.

Lemma 2.4. *Let $\ell \in \mathbb{R}^n$ be a length vector.*

- (1) *Let $n \leq d$. Then $\varphi : \mathcal{M}_{d-1}(\ell) \rightarrow \mathcal{M}_d(\ell)$ is surjective.*
- (2) *Let $n \leq d-1$. Then $\varphi : \mathcal{M}_{d-1}(\ell) \rightarrow \mathcal{M}_d(\ell)$ is a homeomorphism.*
- (3) *Let $z = (z_1, \dots, z_{n-1}) \in \mathcal{C}_d(\ell)$ satisfy $\text{rank } z \geq d-1$. Then z is only fixed by the identity element of $SO(d-1)$.*

Proof. Let $n \leq d$ and $z = [z_1, \dots, z_n] \in \mathcal{M}_d(\ell)$. Since $\sum \ell_i z_i = 0$, the rank of z is at most $n-1 < d$. Thus there exists a $A \in SO(d)$ with $Az_i \in \mathbb{R}^{d-1} \times \{0\} \subset \mathbb{R}^d$ for all $i = 1, \dots, n$. But clearly $[Az_1, \dots, Az_n]$ is in the image of φ .

If $n \leq d - 1$ and $\varphi(z) = \varphi(z')$, we get $\text{rank } z = \text{rank } z' \leq d - 2$. After using rotations in \mathbb{R}^{d-1} we can therefore assume that all $z_i, z'_i \in \mathbb{R}^{\text{rank } z} \subset \mathbb{R}^{d-2} \subset \mathbb{R}^d$. By assumption there is $A \in SO(d)$ with $Az_i = z'_i$ for all $i = 1, \dots, n$, which therefore fixes $\mathbb{R}^{\text{rank } z} \subset \mathbb{R}^{d-2}$. We can now extend $A|_{\mathbb{R}^{\text{rank } z}} \in O(\text{rank } z)$ to $B \in SO(d-1)$ with $Bz_i = z'_i$ for all $i = 1, \dots, n$. But this means $z = z' \in \mathcal{M}_{d-1}(\ell)$, and φ is bijective, hence a homeomorphism by compactness.

Finally, if z has rank at least $d - 1$ and $Az = z$ with $A \in SO(d)$, choose a basis of \mathbb{R}^d where the first $d - 1$ elements are taken from the coordinates of z . Then A fixes at least $d - 1$ elements of a basis of \mathbb{R}^d and is therefore the identity. \square

We remark that if $z \in \mathcal{C}_d(\ell)$ satisfies $\text{rank } z \geq d - 1$, then $n \geq d$. One checks that for $n \geq d$ we can also find $z \in \mathcal{C}_d(\ell)$ which have $\text{rank } z \geq d - 1$. If we denote the dimension of $\mathcal{M}_d(\ell)$ by \mathbf{d}_d^n , we thus get for $n \geq d$ that

$$\mathbf{d}_d^n = (n-3)(d-1) - \frac{(d-2)(d-3)}{2}.$$

3. A MORSE-BOTT FUNCTION ON THE SPACE OF CHAINS

In this section we will assume that $\ell \in \mathbb{R}^n$ is n -dominated.

Define the map $F : \mathcal{C}_d(\ell) \rightarrow \mathbb{R}$ by

$$F(z_1, \dots, z_{n-1}) = \ell_{n-1} p_1(z_{n-1}) + \ell_n$$

where $p_1 : \mathbb{R}^d \rightarrow \mathbb{R}$ is projection to the first coordinate. Notice that F is $SO(d-1)$ invariant and

$$F(z_1, \dots, z_{n-1}) = -p_1 \left(\sum_{i=1}^{n-2} \ell_i z_i \right).$$

We have obvious maxima and minima for points with $z_{n-1} = \pm e_1$. This leads to embeddings of $\mathcal{C}_d(\ell^+)$ and $\mathcal{C}_d(\ell^-)$ into $\mathcal{C}_d(\ell)$, where

$$\begin{aligned} \ell^+ &= (\ell_1, \dots, \ell_{n-2}, \ell_n + \ell_{n-1}) \\ \ell^- &= (\ell_1, \dots, \ell_{n-2}, \ell_n - \ell_{n-1}). \end{aligned}$$

Note that for generic ℓ we can assume that $\ell_n > \ell_{n-1}$, but ℓ^- need not be $n-1$ -dominated.

Let $J \subset \{1, \dots, n-2\}$ be such that $J \cup \{n\}$ is ℓ -short, and $J \cup \{n-1, n\}$ is ℓ -long. Then with $\bar{J} = \{1, \dots, n-2\} - J$ we get

$$\ell_n - \ell_{n-1} < \ell_{\bar{J}} - \ell_J < \ell_n + \ell_{n-1}$$

and there exists a unique $x \in S^1 = \{(x_1, x_2, 0, \dots, 0) \in S^{d-1}\}$ with $x_1 > 0$, $x_2 < 0$ such that $(z_1, \dots, z_{n-1}) \in \mathcal{C}_d(\ell)$ with $z_j = x$ for all $j \in J$ and $z_j = -x$ for $j \in \bar{J}$, compare Figure 1.

The orbit under the $SO(d-1)$ action is a sphere of dimension $d-2$ which we denote by

$$S_J \subset \mathcal{C}_d(\ell).$$

Lemma 3.1. *The critical points of F are given by $\mathcal{C}_d(\ell^\pm)$, and S_J for all $J \subset \{1, \dots, n-2\}$ for which $J \cup \{n\}$ is ℓ -short and $J \cup \{n-1, n\}$ is ℓ -long.*




FIGURE 1.

Proof. We use Lagrange multipliers. Let $f : (\mathbb{R}^d)^{n-1} \times \mathbb{R}^{n-1} \times \mathbb{R}^d \rightarrow \mathbb{R}$ be given by

$$\begin{aligned} f(z_1, \dots, z_{n-1}, \lambda, \mu) &= \ell_{n-1} z_{n-1,1} + \ell_n + \lambda_1 \left(\sum_{j=1}^d z_{1,j}^2 - 1 \right) + \dots + \\ &\quad \lambda_{n-1} \left(\sum_{j=1}^d z_{n-1,j}^2 - 1 \right) + \mu_1 \left(\sum_{i=1}^{n-1} \ell_i z_{i,1} + \ell_n \right) + \\ &\quad \mu_2 \left(\sum_{i=1}^{n-1} \ell_i z_{i,2} \right) + \dots + \mu_d \left(\sum_{i=1}^{n-1} \ell_i z_{i,d} \right) \end{aligned}$$

Taking partial derivatives with respect to $z_{k,j}$ and setting them zero leads to equations

$$\begin{aligned} \mu_j &= -\frac{2\lambda_k z_{k,j}}{\ell_k} \quad \text{for } (j, k) \neq (1, n-1) \\ \mu_1 &= -\frac{\ell_{n-1} + 2\lambda_{n-1} z_{n-1,1}}{\ell_{n-1}}. \end{aligned}$$

For $i = 1, \dots, n-1$, let $\nu_i = \lambda_i / \ell_i$. This implies that

$$z_i \nu_i = z_1 \nu_1$$

for all $i = 1, \dots, n-2$. Since the $z_i \in S^{d-1}$ we either get that $\nu_i = 0$ for all $i = 1, \dots, n-2$ or that $z_1, \dots, z_{n-2} \in \{\pm x\}$ for some $x \in S^{d-1}$.

The case that $\nu_i = 0$ for $i = 1, \dots, n-2$ easily leads to the case where $z_{n-1} = \pm e_1$, which means that $(z_1, \dots, z_{n-1}) \in \mathcal{C}_d(\ell^\pm)$. These points are clearly critical points of F as they are the maxima and minima.

If the ν_i are such that $z_1, \dots, z_{n-2} \in \{\pm x\}$ for some $x \in S^{d-1}$, the condition that $(z_1, \dots, z_{n-1}) \in \mathcal{C}_d(\ell)$ ensures that $(z_1, \dots, z_{n-1}) \in S_J$ for some $J \subset \{1, \dots, n-2\}$. Conversely, let $(z_1, \dots, z_{n-1}) \in S_J$. It is straightforward to check that μ_j and λ_i can be chosen so that all partial derivatives of f vanish. \square

We want to show that F is Morse-Bott. To do this consider the projection $P : \mathcal{C}_d(\ell) \rightarrow S^{d-1}$ given by $P(z) = z_{n-1}$.

Lemma 3.2. *The critical points of $P : \mathcal{C}_d(\ell) \rightarrow S^{d-1}$ are those points for which the first $n-2$ points are collinear.*

Proof. Let $G : (S^{d-1})^{n-2} \rightarrow \mathbb{R}^d$ be given by

$$G(z_1, \dots, z_{n-2}) = \frac{1}{\ell_{n-1}} \left(\ell_n e_1 + \sum_{i=1}^{n-2} \ell_i z_i \right)$$

The critical points of G are clearly the collinear points. We have $\mathcal{C}_d(\ell) = G^{-1}(S^{d-1})$, and since ℓ is generic, we get that G intersects S^{d-1} transversally. Furthermore, P is just the restriction of G to $\mathcal{C}_d(\ell)$, so if $z \in \mathcal{C}_d(\ell)$ is a regular point for G , then z is a regular point for P . Also, if $z \in \mathcal{C}_d(\ell)$ is collinear, the rank of G_* is $d-1$, and since the intersection with S^{d-1} is transversal, the rank of P_* at z is $d-2$. \square

Lemma 3.3. *For generic ℓ , the critical submanifolds $\mathcal{C}_d(\ell^\pm)$ are Morse-Bott with respect to F . Furthermore, the normal bundle is trivial.*

Proof. let $D^{d-1} \subset S^{d-1}$ be a small disc around $\pm e_1$. By Lemma 3.2, we have $P^{-1}(D^{d-1}) \cong D^{d-1} \times \mathcal{C}_d(\ell^\pm)$. The map $F : \mathcal{C}_d(\ell) | P^{-1}(D^{d-1})$ is just a scaling and translation of the standard projection $p_1 : S^{d-1} \rightarrow \mathbb{R}$ to the first coordinate, restricted to D^{d-1} . Since this map is a Morse function with critical points $\pm e_1$, the lemma follows. \square

We remark that ℓ in the following proposition need not be generic, as the S_J stay away from non-manifold points of $\mathcal{C}_d(\ell)$.

Proposition 3.4. *Let ℓ be a length vector and $J \subset \{1, \dots, n-2\}$ such that $J \cup \{n\}$ is ℓ -short and $J \cup \{n-1, n\}$ is ℓ -long. Then S_J is a Morse-Bott critical submanifold of F with index $(n-3-|J|)(d-1)$.*

Proof. Let

$$\mathcal{K}_d(\ell) = \{(z_1, \dots, z_{n-1}) \in \mathcal{C}_d(\ell) \mid z_{n-1} \in S^1 \times \{0\} \subset S^{d-1}\}.$$

This has codimension $d-2$ in $\mathcal{C}_d(\ell)$, and $S_J \cap \mathcal{K}_d(\ell) = S^0$ consists of two points. We claim that $f|_{\mathcal{K}_d(\ell)}$ has Morse singularities near $S_J \cap \mathcal{K}_d(\ell)$, and the proposition follows easily from that.

Let $x \in S^1 \times \{0\} \subset S^{d-1}$ so that $z_J = (\pm x, \dots, \pm x, z_{n-1}) \in \mathcal{K}_d(\ell) \cap S_J$, where we assume that the sign of $\pm x$ is positive of the coordinate is in J , and negative otherwise. Write $x = (\cos \varphi, \sin \varphi)$ and assume $\varphi \in (-\pi/2, 0)$, so that Figure 1 applies.

Note that we can write $F|_{\mathcal{K}_d(\ell)}$ as a composition $\mathcal{K}_d(\ell) \xrightarrow{\tilde{F}} S^1 \xrightarrow{p_1} \mathbb{R}$ with \tilde{F} being projection to $S^1 \times \{0\}$. If we replace $p_1 : S^1 \rightarrow \mathbb{R}$ by $p : S^1 \rightarrow \mathbb{R}$ given by $p(z) = z \cdot x$, that is, scalar product with x , it is clear that near $\tilde{F}(z_J)$ we can write $p_1 = h \circ p$ where h is an orientation preserving diffeomorphism of open intervals of \mathbb{R} .

So to calculate the index of F at z_J we can look at the map $\bar{F} : \mathcal{K}_d(\ell) \rightarrow \mathbb{R}$ given by

$$\bar{F}(z_1, \dots, z_{n-1}) = z_{n-1} \cdot x$$

and calculate its index at the singularity z_J .

Note that we have an inclusion $\mathcal{K}_d(\ell) \subset (S^{d-1})^{n-2}$ as those points for which $\sum_{i=1}^{n-2} \ell_i z_i$ sits inside $\mathbb{R}^2 \times \{0\} \subset \mathbb{R}^d$ with distance ℓ_{n-1} from $(-\ell_n, 0, \dots, 0)$. Using

the Implicit Function Theorem, we can parametrize $\mathcal{K}_d(\ell)$ near z_J as

$$\begin{aligned} (S^{d-1})^{n-3} &\longrightarrow (S^{d-1})^{n-2} \\ (u_1, \dots, u_{n-3}) &\mapsto (u_1, \dots, u_{n-3}, g(u_1, \dots, u_{n-3})) \end{aligned}$$

where near z_J the u_i are close to $\pm x$.

In this parametrization, the map \tilde{F} is given by

$$\tilde{F}(u_1, \dots, u_{n-3}) = \frac{-1}{\ell_{n-1}} \left(\ell_n e_1 + \sum_{i=1}^{n-3} \ell_i u_i + \ell_{n-2} g(u) \right) \cdot x.$$

Use standard polar coordinates for the u_i , that is, we write

$$\begin{aligned} u_i &= (\sin \theta_{d-1 i} \cdots \sin \theta_{2 i} \cos \theta_{1 i}, \sin \theta_{d-1 i} \cdots \sin \theta_{2 i} \sin \theta_{1 i}, \\ &\quad \sin \theta_{d-1 i} \cdots \sin \theta_{3 i} \cos \theta_{2 i}, \dots, \sin \theta_{d-1 i} \cos \theta_{d-2 i}, \cos \theta_{d-1 i}) \end{aligned}$$

for $i = 1, \dots, n-3$, and $\theta_{1 i}$ near φ or $\varphi + \pi$, depending on whether $i \in J$, and $\theta_{j i}$ near $\pi/2$ for $j = 2, \dots, d-1$. The $(n-2)$ -th coordinate can also be written in angles g_j which depend smoothly on the $\theta_{j i}$ for all $i = 1, \dots, n-3$ and $j = 1, \dots, d-1$. After ignoring the scaling factor¹ and the translation through $e_1 \cdot x$, we have to consider the function

$$\begin{aligned} \tilde{F} &= \sum_{i=1}^{n-3} \ell_i (\sin \theta_{d-1 i} \cdots \sin \theta_{2 i} \cos \theta_{1 i} \cos \varphi + \sin \theta_{d-1 i} \cdots \sin \theta_{2 i} \sin \theta_{1 i} \sin \varphi) \\ &\quad + \ell_{n-2} (\sin g_{d-1} \cdots \sin g_2 \cos g_1 \cos \varphi + \sin g_{d-1} \cdots \sin g_2 \sin g_1 \sin \varphi) \\ &= \sum_{i=1}^{n-3} \ell_i \sin \theta_{d-1 i} \cdots \sin \theta_{2 i} (\cos(\theta_{1 i} - \varphi)) + \\ &\quad + \ell_{n-2} \sin g_{d-1} \cdots \sin g_2 (\cos(g_1 - \varphi)) \end{aligned}$$

Writing $\tilde{\theta}_{1 i} = \theta_{1 i} - \varphi + \pi/2$ and $\tilde{g}_1 = g_1 - \varphi + \pi/2$ (and rewriting as $\theta_{1 i}$ and g_1), we get that

$$\begin{aligned} \frac{\partial \tilde{F}}{\partial \theta_{j i}} &= \ell_i \sin \theta_{d-1 i} \cdots \cos \theta_{j i} \cdots \sin \theta_{2 i} \sin \theta_{1 i} \\ &\quad + \sum_{k=1}^{d-1} \ell_{n-2} \sin g_{d-1} \cdots \cos g_k \cdots \sin g_2 \sin g_1 \frac{\partial g_k}{\partial g_{j i}} \end{aligned}$$

Note that the point z_J now corresponds to all angles being $\pi/2$ or $3\pi/2$, so that the cosine terms always vanish. At the point z_J , we therefore get

$$\frac{\partial^2 \tilde{F}}{\partial \theta_{j i} \partial \theta_{j i}} = -\ell_i \sin \theta_{1 i} - \sum_{k=1}^{d-1} \ell_{n-2} \sin g_1 \left(\frac{\partial g_k}{\partial \theta_{j i}} \right)^2$$

and for $(j, i) \neq (l, m)$ we get

$$\frac{\partial^2 \tilde{F}}{\partial \theta_{j i} \partial \theta_{l m}} = -\sum_{k=1}^{d-1} \ell_{n-2} \sin g_1 \frac{\partial g_k}{\partial \theta_{j i}} \frac{\partial g_k}{\partial \theta_{l m}}$$

¹Since we also ignore the -1 , we have to flip the index at the end.

where $\sin \theta_{1i} = 1$ for $i \in J$, -1 for $i \notin J$ and $\sin g_1 = 1$ for $n-2 \in J$ and -1 for $n-2 \notin J$. If we write $\delta_i = +1$ for $i \in J$ and $\delta_i = -1$ for $i \notin J$ ($i \leq n-2$), it follows from Lemma 3.5 that

$$\begin{aligned} \frac{\partial^2 \tilde{F}}{\partial \theta_{ji} \partial \theta_{ji}} &= -\ell_i \delta_i - \delta_{n-2} \frac{\ell_i^2}{\ell_{n-2}} \\ \frac{\partial^2 \tilde{F}}{\partial \theta_{ji} \partial \theta_{lm}} &= 0 \quad \text{for } j \neq l \\ \frac{\partial^2 \tilde{F}}{\partial \theta_{ji} \partial \theta_{jm}} &= \begin{cases} -\delta_{n-2} \delta_i \delta_m \frac{\ell_i \ell_m}{\ell_{n-2}} & j = 1 \\ -\delta_{n-2} \frac{\ell_i \ell_m}{\ell_{n-2}} & j \geq 2 \end{cases} \end{aligned}$$

The matrix $\left(\frac{\partial^2 \tilde{F}}{\partial \theta_{ji} \partial \theta_{lm}} \right)$ is a $(n-3)(d-1) \times (n-3)(d-1)$ matrix, which we consider as a $(d-1) \times (d-1)$ matrix with entries $(n-3) \times (n-3)$ matrices $\left(\frac{\partial^2 \tilde{F}}{\partial \theta_{ji} \partial \theta_{lm}} \right)_{i,m}$ for fixed j, l . The off-diagonal entries are then 0, while the diagonal entries are matrices $\left(\frac{\partial^2 \tilde{F}}{\partial \theta_{ji} \partial \theta_{jm}} \right)_{i,m}$ for $j = 1, \dots, d-1$. These matrices are of the form

$$\begin{aligned} \left(\frac{\partial^2 \tilde{F}}{\partial \theta_{1i} \partial \theta_{1m}} \right)_{i,m} &= -\Delta(\ell_1 \delta_1, \dots, \ell_{n-3} \delta_{n-3}) - \frac{\delta_{n-2}}{\ell_{n-2}} (\delta_i \ell_i \delta_m \ell_m)_{i,m} \\ \left(\frac{\partial^2 \tilde{F}}{\partial \theta_{ji} \partial \theta_{jm}} \right)_{i,m} &= -\Delta(\ell_1 \delta_1, \dots, \ell_{n-3} \delta_{n-3}) - \frac{\delta_{n-2}}{\ell_{n-2}} (\ell_i \ell_m)_{i,m} \end{aligned}$$

for $j = 2, \dots, d-1$. Here Δ is a diagonal matrix with the given entries.

Since $|J| \leq n-3$, we can assume (after possibly rearranging the order of the links) that $n-2 \notin J$, that is, $\delta_{n-2} = -1$. It follows that these matrices are congruent to

$$M = -\frac{1}{\ell_{n-2}} \left(\Delta \left(\frac{\ell_{n-2} \delta_1}{\ell_1}, \dots, \frac{\ell_{n-2} \delta_{n-3}}{\ell_{n-3}} \right) - E \right)$$

where E has every entry equal to 1. Ignoring the factor $-1/\ell_{n-2}$, which leads to the second flipping of the index, compare Footnote 1, we need to calculate the index of $\Delta \left(\frac{\ell_{n-2} \delta_1}{\ell_1}, \dots, \frac{\ell_{n-2} \delta_{n-3}}{\ell_{n-3}} \right) - E$. A calculation as in [3, Lm.1.4] shows that the index is $n-3 - |J|$. Since we have $d-1$ such matrices, the result follows. \square

Lemma 3.5. *With notation as in Proposition 3.4, we have*

$$\begin{aligned} \frac{\partial g_k}{\partial \theta_{ji}} &= 0 \quad \text{for } k \neq j \\ \frac{\partial g_k}{\partial \theta_{ki}} &= \begin{cases} \delta_i \frac{\ell_i}{\ell_{n-2}} & k = 1 \\ \frac{\ell_i}{\ell_{n-2}} & k \geq 2 \end{cases} \end{aligned}$$

Proof. Define $G : (S^{d-1})^{n-2} \rightarrow \mathbb{R}^{d-1}$

$$G(u_1, \dots, u_{n-2}) = \begin{pmatrix} \left| \sum_{i=1}^{n-2} \ell_i u_i + \ell_n e_1 \right|^2 \\ p_3 \left(\sum_{i=1}^{n-2} \ell_i u_i \right) \\ \vdots \\ p_d \left(\sum_{i=1}^{n-2} \ell_i u_i \right) \end{pmatrix}$$

Then $\mathcal{K}_d(\ell) = G^{-1}(\ell_{n-1}, 0, \dots, 0)$. In polar coordinates a calculation shows that

$$\begin{aligned} G_1 &= \ell_n^2 + \sum_{i=1}^{n-2} \ell_i^2 + 2\ell_n \sum_{i=1}^{n-2} \ell_i \sin \theta_{d-1 i} \cdots \sin \theta_{2 i} \cos \theta_{1 i} \\ &\quad + 2 \sum_{i < j} \ell_i \ell_j (\sin \theta_{d-1 i} \cdots \sin \theta_{2 i} \sin_{d-1 j} \cdots \sin \theta_{2 j} (\cos(\theta_{1 i} - \theta_{1 j}) \\ &\quad \quad \quad + \sin \theta_{d-1 i} \cdots \sin \theta_{3 i} \cos \theta_{2 i} \sin \theta_{d-1 j} \cdots \sin \theta_{3 j} \cos \theta_{2 j} \\ &\quad \quad \quad + \cdots + \cos \theta_{d-1 i} \cos \theta_{d-1 j})). \end{aligned}$$

Similarly, for $k \geq 2$ we have

$$G_k = \sum_{i=1}^{n-2} \ell_i \sin \theta_{d-1 i} \cdots \sin \theta_{k+1 i} \cos \theta_{k i}.$$

Using the fact that z_J has $\theta_{j i} = \pi/2$ for $j > 1$, it is easy to see that

$$\begin{aligned} \frac{\partial G_1}{\partial \theta_{1 i}}(z_J) &= \begin{cases} 2\ell_n \ell_i \sin \varphi & i \in J \\ -2\ell_n \ell_i \sin \varphi & i \notin J \end{cases} \\ \frac{\partial G_k}{\partial \theta_{j i}}(z_J) &= 0 \quad \text{for } j \neq k \\ \frac{\partial G_k}{\partial \theta_{k i}}(z_J) &= -\ell_i \quad \text{for } k \geq 2 \end{aligned}$$

So for fixed $i \leq n-2$, each $\left(\frac{\partial G_k}{\partial \theta_{j i}}(z_J)\right)$ is an invertible diagonal matrix. In particular, by the Implicit Function Theorem we get for $i \leq n-3$

$$\begin{aligned} \frac{\partial g_k}{\partial \theta_{j i}} &= 0 \quad \text{for } k \neq j \\ \frac{\partial g_k}{\partial \theta_{k i}} &= \begin{cases} \delta_i \frac{\ell_i}{\ell_{n-2}} & k = 1 \\ \frac{\ell_i}{\ell_{n-2}} & k \geq 2 \end{cases} \end{aligned}$$

since the g_k are obtained by applying the Implicit Function Theorem to G . \square

4. PERFECT MORSE FUNCTIONS FOR PLANAR POLYGON SPACES

We can apply the results from the previous section to construct perfect Morse functions on the manifolds $\mathcal{M}_2(\ell) \cong \mathcal{C}_2(\ell)$ for generic ℓ . In this case, each S_J consists of 2 points, and the index is $n-3-|J|$. The maxima and minima give of course rise to Morse-Bott singularities, which are not Morse in general.

Recall that up to scaling and translation, $F : \mathcal{C}_d(\ell) \rightarrow \mathbb{R}$ factors as $p_1 \circ P$ with $P : \mathcal{C}_d(\ell) \rightarrow S^{d-1}$ projection to the $(n-1)$ -th coordinate. In the generic case and if $d = 2$ the critical points are exactly the 0-spheres S_J for $J \subset \{1, \dots, n-2\}$ such that $J \cup \{n\}$ is ℓ -short and $J \cup \{n-1, n\}$ is ℓ -long. In fact, we can consider the resulting map $P : \mathcal{C}_2(\ell) \rightarrow S^1$ a Morse function to the circle. To calculate the index of the appropriate critical points, note that S^1 can be given the orientation in the counterclockwise direction. Projection to the x -axis is then locally orientation reversing, if $z \in S^1$ has positive imaginary part, and orientation preserving if it has negative imaginary part. Each 0-sphere S_J has a point which is mapped to the upper half of S^1 and one which is mapped to the lower half of S^1 . Therefore, one

of these critical points has index $n - 3 - |J|$ and the other has index $|J|$. We have thus proved the following.

Proposition 4.1. *Let $\ell \in \mathbb{R}^n$ be an n -dominated generic length vector. Then the projection $P : \mathcal{C}_2(\ell) \rightarrow S^1$ to the $(n - 1)$ -th link is a Morse function whose critical points are in one-to-one correspondence to 0-spheres S_J to subsets $J \subset \{1, \dots, n - 2\}$ such that $J \cup \{n\}$ is ℓ -short and $J \cup \{n - 1, n\}$ is ℓ -long. Furthermore, the indices of the pair S_J are $n - 3 - |J|$ and $|J|$.*

Circle-valued Morse functions need not have maxima. In fact, in order to get a maximum, we would need a J with $|J| = 0$ or $|J| = n - 3$. If $|J| = n - 3$, then there is $i \leq n - 2$ such that $\{i, n - 1\}$ is ℓ -long. Since ℓ is n -dominated, this implies that $\mathcal{C}_2(\ell) = T^{n-3} \times S^0$, and there is only one such chamber up to permutations.

If $|J| = 0$, then $J = \emptyset$, and $\{n - 1, n\}$ is long. This actually means that the map $P : \mathcal{C}_2(\ell) \rightarrow S^1$ is not surjective, and we can interpret this map as a map $\tilde{P} : \mathcal{C}_2(\ell) \rightarrow \mathbb{R}$.

Proposition 4.2. *If $\{n - 1, n\}$ is ℓ -long, then the Morse function $\tilde{P} : \mathcal{C}_2(\ell) \rightarrow \mathbb{R}$ is perfect.*

Proof. The homology of $\mathcal{C}_2(\ell)$ is free abelian, and every $J \subset \{1, \dots, n - 1\}$ with $J \cup \{n\}$ short contributes two elements towards a basis, compare [6]. Since $\{n - 1, n\}$ is long, such J already has to satisfy $J \subset \{1, \dots, n - 2\}$. Furthermore, $J \cup \{n - 1, n\}$ has to be long. Therefore, any such J gives rise to two critical points, and these give all the critical points of \tilde{P} . Hence the map is perfect. \square

For the general case we have to make a modification to construct a perfect Morse function.

Theorem 4.3. *Let ℓ be a generic length vector. Then there exists a perfect Morse function $f : \mathcal{M}_2(\ell) \rightarrow \mathbb{R}$.*

Proof. The proof is by induction on n . If $n \leq 5$, the statement is obvious because there are only very few cases which can be checked by hand. For $n > 5$, let $I \subset S^1$ be a small closed interval neighborhood of $1 \in S^1$ which does not contain any critical values of $P : \mathcal{M}_2(\ell) \cong \mathcal{C}_2(\ell) \rightarrow S^1$.

Since there are no critical values in I , we get for $z \in \partial I$ that $P^{-1}(z) \cong P^{-1}(1) \cong \mathcal{M}_2(\ell^+)$.

Let $h : S^1 \rightarrow \mathbb{R}$ be a perfect Morse function whose two critical points are ∂I . The composition $h \circ P : \mathcal{M}_2(\ell) \rightarrow \mathbb{R}$ is then a Morse-Bott function with Morse critical points for every $J \subset \{1, \dots, n - 2\}$ with $J \cup \{n\}$ ℓ -short and $J \cup \{n - 1, n\}$ ℓ -long, a singular manifold $\mathcal{M}_2(\ell^+)$ of index 0 and another $\mathcal{M}_2(\ell^+)$ of index 1.

By induction, there exists a perfect Morse function $g : \mathcal{M}_2(\ell^+) \rightarrow \mathbb{R}$, which can be glued into $h \circ P$ to give a Morse function $f : \mathcal{M}_2(\ell) \rightarrow \mathbb{R}$. Since g is perfect, every $J \subset \{1, \dots, n - 2\}$ such that $J \cup \{n - 1\}$ is ℓ^+ -short gives rise to two critical points of g , whose indices are $n - 4 - |J|$ and $|J|$.

The critical points of f are therefore as follows:

- Every $J \subset \{1, \dots, n - 2\}$ with $J \cup \{n\}$ ℓ -short and $J \cup \{n - 1, n\}$ ℓ -long gives rise to two critical points of index $|J|$ and $n - 3 - |J|$.

- Every $J \subset \{1, \dots, n-2\}$ with $J \cup \{n-1\}$ ℓ^+ -short gives rise to two pairs of critical points of index $|J|, n-4-|J|$, and $|J|+1, n-3-|J|$.

We claim that f is perfect. Every $J \subset \{1, \dots, n-1\}$ with $J \cup \{n\}$ ℓ -long produces to \mathbb{Z} -summands in $H_*(\mathcal{M}_2(\ell))$, and therefore has to correspond uniquely to a pair of critical points listed above.

Let $J \subset \{1, \dots, n-2\}$ with $J \cup \{n\}$ ℓ -short and $J \cup \{n-1, n\}$ ℓ -long. Then $J \cup \{n-1\}$ is ℓ^+ -long, so does not produce critical points coming from g .

If $J \subset \{1, \dots, n-2\}$ has $J \cup \{n-1, n\}$ ℓ -short, then $J \cup \{n-1\}$ is ℓ^+ -short, so we get four critical points. Two of them, with index $|J|$ and $n-3-|J|$ correspond to the set J , which satisfies $J \cup \{n\}$ ℓ -short. The other two, of index $|J|+1$ and $n-4-|J|$ correspond to $J \cup \{n-1\}$, which satisfies $J \cup \{n-1\} \cup \{n\}$ is ℓ -short. Therefore the critical points of f do correspond to a basis of the homology, which means that f is perfect. \square

Perfect Morse functions for some special cases of ℓ were constructed in [14]. It is worth pointing out that the signed area of a linkage induces a Morse function on $\mathcal{M}_2(\ell)$, but this Morse function need not be perfect: see [11], where it is shown that for $\ell = (1, 1, 1, 1, 1)$ the signed area has two maxima and two minima.

Let us return to the circle-valued map $P : \mathcal{C}_2(\ell) \rightarrow S^1$. One can ask for perfectness of this function in the sense of Novikov homology, compare [2, 20]. In fact, if $\{n-1, n\}$ is ℓ -short, then there do not exist maxima and minima, and it is easy to see that P induces a surjection on fundamental group. Let $\tilde{\mathcal{M}}_2(\ell)$ be the regular \mathbb{Z} -covering space corresponding to $\ker P_\#$. Then P lifts to a smooth map $\tilde{P} : \tilde{\mathcal{M}}_2(\ell) \rightarrow \mathbb{R}$ whose critical points are liftings of the critical points of P and who are Morse with associated indices.

Assume that $\mathcal{M}_2(\ell)$ is triangulated. The triangulation can be lifted to $\tilde{\mathcal{M}}_2(\ell)$, and the corresponding chain complex $C_*(\tilde{\mathcal{M}}_2(\ell))$ is a finitely generated free chain complex over the ring $\mathbb{Z}[t, t^{-1}]$. Let $\mathbb{Z}((t))$ be the ring whose elements are Laurent power series of the form $\sum_{i=-r}^{\infty} a_i t^i$ for some $r \in \mathbb{Z}$ and $a_i \in \mathbb{Z}$. The homology

$$H_*(\mathcal{M}_2(\ell); \mathbb{Z}((t))) = H_*(\mathbb{Z}((t)) \otimes_{\mathbb{Z}[t, t^{-1}]} C_*(\tilde{\mathcal{M}}_2(\ell)))$$

is called the *Novikov homology* of $\mathcal{M}_2(\ell)$ with respect to P .

The ring $\mathbb{Z}((t))$ is a Euclidean ring, see [2, 20], so these homology groups have a finitely generated free part and a torsion part.

Theorem 4.4. *Let $\ell \in \mathbb{R}^n$ be an n -dominated and generic length vector such that $\{n-1, n\}$ is ℓ -short, and let $P : \mathcal{M}_2(\ell) \rightarrow S^1$ be projection to the $(n-1)$ -th link. Then*

$$H_i(\mathcal{M}_2(\ell); \mathbb{Z}((t))) = \mathbb{Z}((t))^{a_i + a_{n-3-i}}$$

where a_i is the number of subsets $J \subset \{1, \dots, n-2\}$ with $J \cup \{n\}$ ℓ -short and $J \cup \{n-1, n\}$ ℓ -long.

In other words, P is perfect with respect to Novikov homology.

Proof. Just as in ordinary Morse theory, Novikov homology can be calculated from the circle-valued Morse function by using chain complex over $\mathbb{Z}((t))$ generated by the critical points and counting trajectories between critical points of adjacent

index, see [19]. We claim that the boundary is always the zero map. To see this, let $\tilde{p}, \tilde{q} \in \tilde{\mathcal{M}}_2(\ell)$ be liftings of two critical points $p, q \in \mathcal{M}_2(\ell)$ of P of adjacent index. We need to show that the incidence numbers between \tilde{p} and $t^k \tilde{q}$ are zero for all $k \in \mathbb{Z}$. Here $t : \tilde{\mathcal{M}}_2(\ell) \rightarrow \tilde{\mathcal{M}}_2(\ell)$ is the covering translation satisfying $\tilde{P}(z) - \tilde{P}(t(z)) = 1$. To calculate the incidence number, we only have to look at a compact cobordism $W = \tilde{P}^{-1}([a, b])$, where $[a, b]$ is a compact interval large enough so that \tilde{p} and $t^k \tilde{q}$ are in W (for fixed k). That the incidence number is zero follows from [6, Thm.4], once we produce an involution on W which fixes exactly the critical points. Note that [6, Thm.4] applies to manifolds whose boundary is the maximum, but the same argument works for cobordisms with the homology being relative with respect to the negative boundary.

The involution can be constructed on $\mathcal{M}_2(\ell)$ and then lifted to W . Let $z \in \mathcal{M}_2(\ell) \cong \mathcal{C}_2(\ell)$. Since $\ell_{n-1} < \ell_n$, which we can assume because ℓ is generic, we get that

$$\sum_{i=1}^{n-2} \ell_i z_i \neq 0.$$

If we let $T_z : S^1 \rightarrow S^1$ be the standard reflection which fixes the line through this point and 0, then $(T_z(z_1), \dots, T_z(z_{n-2}), z_{n-1}) \in \mathcal{C}_2(\ell)$, compare Figure 2.




FIGURE 2. Three links being reflected in the dotted line.

Define $T : \mathcal{C}_2(\ell) \rightarrow \mathcal{C}_2(\ell)$ by $T(z_1, \dots, z_{n-1}) = (T_z(z_1), \dots, T_z(z_{n-2}), z_{n-1})$. Since

$$\sum_{i=1}^{n-2} \ell_i z_i = \sum_{i=1}^{n-2} \ell_i T_z(z_i),$$

this function is an involution and it preserves P . Therefore it lifts to an involution on $\tilde{\mathcal{M}}_2(\ell)$ and W . Furthermore, the critical points of P are exactly the points which are fixed by T , since T_z is then the reflection in the line through x , that is, x and $-x$ are fixed. Hence the boundary in the Novikov complex is just 0. \square

Nonvanishing Novikov homology is an obstruction for P to be homotopic to a fibre bundle map over S^1 . On the other hand, if P has no critical points then $\mathcal{M}_2(\ell)$ fibres over S^1 along P . Let us analyze this situation in more detail.

If ℓ is a generic length vector, then there exists a $\delta > 0$ such that

$$\left| \sum_{j \in J} \ell_j - \sum_{i \notin J} \ell_i \right| > \delta$$

for all subsets $J \subset \{1, \dots, n\}$. For $0 < \varepsilon < \delta$ it is easy to see that the length vector $\ell_\varepsilon = (\varepsilon, \ell_1, \dots, \ell_n)$ satisfies

$$\mathcal{M}_2(\ell_\varepsilon) \cong S^1 \times \mathcal{M}_2(\ell)$$

and projection to the first coordinate gives the fibration over S^1 .

Assume P has no critical points. Then every $J \subset \{1, \dots, n-2\}$ with $J \subset \{n\}$ short also satisfies $J \cup \{n-1, n\}$ short. Let

$$\ell_t = (\ell_1, \dots, \ell_{n-2}, \ell_{n-1} - t, \ell_n + t)$$

for $0 \leq t < \ell_{n-1}$. Since P has no critical points, it is easy to see that every $J \subset \{1, \dots, n\}$ is ℓ -short if and only if J is ℓ_t -short for any $0 \leq t < \ell_{n-1}$, so ℓ_t is in the same chamber as ℓ . But for t close to ℓ_{n-1} we get $\mathcal{M}_2(\ell) \cong S^1 \times \mathcal{M}_2(\ell')$, where $\ell' = (\ell_1, \dots, \ell_{n-2}, \ell_{n-1} + \ell_n)$. So if P has no critical points, the $(n-1)$ -th link is already so small that $\mathcal{M}_2(\ell)$ fibres over S^1 for obvious reasons.

It is therefore a natural question to ask whether $\mathcal{M}_2(\ell)$ fibres over the circle exactly when there is a short link in the length vector. This is true by the above if we only look at maps $\mathcal{M}_2(\ell) \rightarrow S^1$ given by projections to a coordinate, but it is not clear if we look at homotopy classes of arbitrary maps to S^1 .

5. HOMOLOGY FOR THE 3-DIMENSIONAL CASE

In this section we show how the Betti numbers for $\mathcal{M}_3(\ell)$ can be obtained from the Morse-Bott function above. We will only sketch the argument, as these results have already been obtained in [18]. Information on the cohomology is contained in [9].

For $d = 3$ and generic ℓ , the action of $SO(2)$ is free on $\mathcal{C}_3(\ell)$, and $\mathcal{M}_3(\ell)$ is a closed manifold. Furthermore, the $SO(2)$ -invariant function F induces a Morse-Bott function $f : \mathcal{M}_3(\ell) \rightarrow \mathbb{R}$, which has $\mathcal{M}_3(\ell^-)$ as minimum, $\mathcal{M}_3(\ell^+)$ as maximum (with index 2), and for each $J \subset \{1, \dots, n-2\}$ with $J \cup \{n\}$ short and $J \cup \{n-1, n\}$ long a critical point p_J of index $2(n-3-|J|)$.

A simple induction argument using the Morse-Bott spectral sequence shows that the homology of $\mathcal{M}_3(\ell)$ is free abelian and concentrated in even degrees. If we write $P_\ell(t)$ for the Poincaré polynomial of $\mathcal{M}_3(\ell)$, we get the following recursive formula.

Proposition 5.1. *Let $\ell \in \mathbb{R}^n$ be a generic length vector. Then the Poincaré polynomial of $\mathcal{M}_3(\ell)$ satisfies*

$$P_\ell(t) = P_{\ell^-}(t) + t^2 P_{\ell^+}(t) + \sum_{J \subset \mathcal{T}(\ell)} t^{2|J|}$$

where $\mathcal{T}(\ell) = \{J \subset \{1, \dots, n-2\} \mid J \cup \{n\} \text{ short, } J \cup \{n-1, n\} \text{ long}\}$.

Remark 5.2. A similar recursive formula is obtained in [18, Cor.2.2.2] by using different methods. In fact, we can get that formula by looking at $-f$ instead of f .

Klyachko goes on to give the following explicit formula for the Poincaré polynomial, see [18, Thm.2.2.4].

$$P_\ell(t) = \frac{1}{t^2(t^2-1)} \left((1+t^2)^{n-1} - \sum_{J \in \mathcal{S}(\ell)} t^{2|J|} \right),$$

where $\mathcal{S}(\ell) = \{J \subset \{1, \dots, n\} \mid J \text{ short}\}$.

Hausmann and Knutson [9, Cor.4.3] derive another formula, given by

$$P_\ell(t) = \frac{1}{1-t^2} \sum_{J \in \mathcal{S}_*^n(\ell)} (t^{2|J|} - t^{2(n-2-|J|)}),$$

where we assume that ℓ is n -dominated.

Corollary 5.3. *Let $\ell \in \mathbb{R}^n$ be a generic length vector. Then there exists a perfect Morse function $f_3 : \mathcal{M}_3(\ell) \rightarrow \mathbb{R}$, all of whose critical points are of even index.*

Proof. The proof is by induction on n . For $n = 4$, we get $\mathcal{M}_3(\ell) \cong S^2$ and the statement is clear. For $n > 4$ look at the Morse-Bott function $f : \mathcal{M}_3 \rightarrow \mathbb{R}$ with minimum attained at $\mathcal{M}_3(\ell^-)$ and maximum attained at $\mathcal{M}_3(\ell^+)$. Using standard techniques to replace the Morse-Bott critical manifolds by a Morse function on these manifolds, which can be assumed to be perfect by induction, we get a Morse function on $\mathcal{M}_3(\ell)$. Notice that since the index of the Morse-Bott manifolds are even, the indices of the critical points are also even. It follows that the Morse function is perfect. \square

Let us give a formula for the number of critical points of a given index. For this let $\mu_k(\ell)$ be the number of critical points of f_3 having index $2k$.

Proposition 5.4. *Let $\ell \in \mathbb{R}^n$ be a generic length vector, and let $m \in \mathbb{Z}$ be such that $n = 2m - 1$ or $n = 2m$. Then*

$$\mu_k(\ell) = \sum_{i=0}^k a_i(\ell) - a_{n-2-i}(\ell)$$

for all $k = 0, \dots, m-2$, and

$$\mu_k(\ell) = \mu_{n-3-k}(\ell)$$

for all $k = m-1, \dots, n-3$.

Proof. The second equation just follows from Poincaré duality, so let us assume that $k \leq m-2$.

We use the formula from [9]. In terms of the coefficients $a_k(\ell)$ this is

$$P_\ell(t) = \frac{1}{1-t^2} \sum_{i=0}^{n-3} a_i(\ell) (t^{2i} - t^{2(n-2-i)}).$$

Since

$$(1-t^2)(t^{2i} + t^{2(i+1)} + \dots + t^{2(n-3-i)}) = t^{2i} - t^{2(n-2-i)}$$

we can write the Poincaré polynomial as

$$(1) \quad P_\ell(t) = \sum_{i=0}^{m-2} (a_i(\ell) - a_{n-2-i}(\ell)) (t^{2i} + t^{2(i+1)} + \dots + t^{2(n-3-i)}).$$

Comparing coefficients then gives the result. \square

6. AN EQUIVARIANT CELL DECOMPOSITION FOR $\mathcal{C}_d(\ell)$

We want to derive an equivariant cell decomposition for $\mathcal{C}_d(\ell)$ using the Morse-Bott function F in order to get a cell decomposition for $\mathcal{M}_d(\ell)$ for $d \geq 3$.

To do this we first want to understand the equivariant handle structure near a critical manifold S_J in the sense of [24]. So if $J \subset \{1, \dots, n-2\}$ has the property that S_J is a critical sphere, let $x \in S^1 \times \{0\}$ be such that $p_J = (\pm x, \dots, \pm x, z_{n-1}) \in S_J$ and the minus signs correspond to coordinates from J . We may assume that $n-2 \in J$. Let $D^{d-1} \subset S^{d-1}$ be a small disc with center at $-x$, and define

$$\begin{aligned} P : (D^{d-1})^{n-3-|J|} &\longrightarrow \mathcal{K}_d(\ell) \\ (u_1, \dots, u_{n-3-|J|}) &\longmapsto (v_1, \dots, v_{n-3}, g(u_1, \dots, u_{n-3-|J|})) \end{aligned}$$

where $v_i = x$ if $i \notin J$, $v_i = u_{k_i}$ for $i \in J = \{k_1, \dots, k_{n-3-|J|}\}$. That is, we use the parametrization of $\mathcal{K}_d(\ell)$ from the proof of Proposition 3.4, but we keep the coordinates away from J fixed.

By the same argument as in the proof of Proposition 3.4, $F \circ P$ has a nondegenerate maximal point at $(-x, \dots, -x)$, which is the center of $(D^{d-1})^{n-3-|J|}$.

For simplicity, let us center D^{d-1} at 0, and we think of P as an inclusion $i : (D^{d-1})^{n-3-|J|} \rightarrow \mathcal{C}_d(\ell)$. If we let $SO(d-2)$ act diagonally on $(D^{d-1})^{n-3-|J|}$, with $SO(d-2)$ acting in a standard way on $D^{d-1} \subset \mathbb{R}^{d-1}$ by fixing the first coordinate, we get that i is $SO(d-2)$ equivariant.

The image of i is in $\mathcal{K}_d(\ell)$, and by using the action of $SO(d-1)$ on the image, we get the negative normal bundle of S_J in the sense of equivariant Morse theory, compare [24]. We thus write

$$N^-(S_J) = \left\{ Ai(x) \in \mathcal{C}_d(\ell) \mid A \in SO(d-1), x \in (D^{d-1})^{n-3-|J|} \right\}.$$

The map $N^-(S_J) \rightarrow SO(d-1)/SO(d-2) \cong S_J$ given by $Ai(x) \mapsto A \cdot SO(d-2)$ is then a disc bundle map with fibre $(D^{d-1})^{n-3-|J|}$.

We want to have an equivariant Morse-Bott function $\tilde{F} : \mathcal{C}_d(\ell) \rightarrow \mathbb{R}$ such that all critical manifolds are spheres $SO(d-1)/SO(d-2)$ with negative normal bundle as the $N^-(S_J)$. The idea is to use the argument in the proof of Corollary 5.3, but equivariantly. This can be done, as there are neighborhoods of $\mathcal{C}_d(\ell^\pm)$ in $\mathcal{C}_d(\ell)$ which are equivariantly diffeomorphic to $\mathcal{C}_d(\ell) \times D^{d-1}$, compare Lemma 3.3 and use the fact that for ℓ and ℓ' in the same chamber, the chain spaces are equivariantly diffeomorphic [8]. Notice that the critical manifolds do not depend on d . We thus get the following result.

Proposition 6.1. *Let $\ell \in \mathbb{R}^n$ be a generic length vector. For all $d \geq 3$ there is an $SO(d-1)$ invariant Morse-Bott function $\tilde{F} : \mathcal{C}_d(\ell) \rightarrow \mathbb{R}$ such that all critical manifolds are of the form $SO(d-1)/SO(d-2)$, and their indices are of the form $k(d-1)$ for some $k = 0, \dots, n-3$. The negative normal bundle $N^-(S)$ to each critical manifold S is of the form $(D^{d-1})^k \rightarrow N^-(S) \rightarrow SO(d-1)/SO(d-2)$, where $SO(d-2)$ acts on $(D^{d-1})^k$ diagonally, and by fixing the first coordinate of D^{d-1} .*

Furthermore, the critical manifolds S of index $k(d-1)$ are in one-to-one correspondence to the critical points of index $2k$ of the perfect Morse function $f_3 : \mathcal{M}_3(\ell) \rightarrow \mathbb{R}$ from Corollary 5.3. \square

Denote by $\partial N^-(S)$ the sphere bundle corresponding to $N^-(S)$. In order to understand the homotopy type of $\mathcal{M}_d(\ell)$ we want to understand a relative $SO(d-1)$ -equivariant cell structure on $(N^-(S), \partial N^-(S))$. Since $N^-(S)$ is the $SO(d-1)$ orbit of the image of $(D^{d-1})^k$, we have to find a relative $SO(d-2)$ -equivariant cell structure of $((D^{d-1})^k, \partial(D^{d-1})^k)$.

Let us begin with some elementary observations. If $k = 1$, the set $D^1 \times \{0\} \subset D^{d-1}$ is the fixed set of the $SO(d-2)$ action. It therefore defines a 1-cell with $SO(d-2)$ as the stabilizer group. If $x \in D^{d-1} - D^1 \times \{0\}$, we can find an $A \in SO(d-2)$ such that $Ax = (a, b, 0, \dots, 0) \in D^{d-1}$, with $b \neq 0$. If $d > 3$, we can furthermore assume that $b > 0$. In particular, every other element of D^{d-1} will be in the orbit of an element of $D_+^2 = \{(a, b, 0, \dots, 0) \in D^{d-1} \mid b \geq 0, a^2 + b^2 \leq 1\}$.

In particular, we only need two cells. If we denote $X = D^{d-1}/SO(d-2)$ and $\partial X = \partial D^{d-1}/SO(d-2)$, we get a relative CW-structure of $(X, \partial X)$ with X being obtained from ∂X by an elementary expansion in the sense of [1, §4]. If $d = 3$, note that $SO(d-2)$ is the trivial group. We either have to use two 2-cells (one for $b > 0$ and one for $b < 0$), or we do not use the 1-cell, and just use the 2-cell D^2 .

We can ignore the case $d = 3$, in which case we only need one cell for $((D^2)^k, \partial(D^2)^k)$ of dimension $2k$. So assume $d \geq 4$ now. Let $(x_1, \dots, x_k) \in (D^{d-1})^k$. After applying an element of $SO(d-2)$ we can assume $x_1 \in D_+^2$. If we actually have $x_1 \in D^1$, we apply another element of $SO(d-2)$ to get $x_2 \in D_+^2$. We repeat this until we get an element $x_i \in D_+^2 - D^1$. If we do not get such an element, the original element (x_1, \dots, x_k) is in $(D^1)^k$. So assume $x_i \in D_+^2 - D^1$ and $x_j \in D^1$ for $j < i$. Applying an element of $SO(d-3)$ does not affect the first i elements, and can move x_{i+1} into D^3 , in fact D_+^3 if $d > 4$. We can continue this and achieve that up to elements of $SO(d-2)$, the element (x_1, \dots, x_k) is in a product of an increasing sequence of discs.

To make this more precise, write

$$D^i = \{(x_1, \dots, x_i, 0, \dots, 0) \in D^{d-1} \mid x_1^2 + \dots, x_i^2 \leq 1\}$$

for $i = 1, \dots, d-2$, and also write

$$D_+^i = \{(x_1, \dots, x_i, 0, \dots, 0) \in D^{d-1} \mid x_1^2 + \dots, x_i^2 \leq 1, x_i \geq 0\}$$

for $i = 2, \dots, d-2$.

Up to an element of $SO(d-2)$, any $(x_1, \dots, x_k) \in (D^{d-1})^k$ sits in

$$(D^1)^{k_1} \times (D^2)^{k_2} \times \dots \times (D^{d-2})^{k_{d-2}} \times (D^{d-1})^{k_{d-1}}$$

where all $k_i \geq 0$ and add up to k . Furthermore, if $k_i = 0$ for $i \geq 2$, then all $k_j = 0$ with $j \geq i$, and if $k_i \neq 0$ for $i \in \{2, \dots, d-2\}$, we can replace $(D^i)^{k_i}$ by $D_+^i \times (D^i)^{k_i-1}$.

In order to organise the cells we are going to introduce symbolic matrices. For $n, m \geq 1$ let $\mathbf{S}(m, n)$ be the set of upper semi-diagonal $n \times m$ matrices whose entries are from the set $\{0, +, *\}$, which have a $+$ sign for the first non-zero entry in each of the first $n-1$, with all entries to the right of the $+$ as $*$, and the last row contains only 0 and $*$, with no 0 to the right of any $*$.

Typical examples are

$$(2) \quad \left(\begin{array}{cccccccc} 0 & 0 & + & * & * & * & * \\ 0 & 0 & 0 & + & * & * & * \\ 0 & 0 & 0 & 0 & 0 & * & * \end{array} \right), \left(\begin{array}{cccc} + & * & * & * \\ 0 & 0 & + & * \\ 0 & 0 & 0 & 0 \end{array} \right).$$

Each matrix stands for a product of discs, with columns referring to each disc. Here the zero column stands for D^1 , a column containing a $+$ stands for D_+^k and a column with only $*$ and 0 stands for D^k , where $k - 1$ is the number of non-zero entries in the column.

Remark 6.2. Using such symbolic matrices was already done in [17] to get a cell decomposition of the shape spaces Σ_d^m , and our homology calculations below are indeed quite similar to the calculations in [17].

So if $A \in \mathbf{S}(d - 2, k)$, we denote by $D_A \subset (D^{d-1})^k$ the corresponding product of discs. Also, let $SO(A) \subset SO(d - 2)$ be the stabilizer group of D_A . Then $SO(A) = SO(d - 2 - i)$, where i is the maximal number of non-zero elements in the columns of A , and $SO(d - 2 - i)$ acts on D^{d-1} by fixing the first $i + 1$ coordinates. In particular $SO(0)$ is the trivial group. We denote the image of D_A under the $SO(d - 2)$ -action by σ_A , and call this the cell corresponding to A .

Lemma 6.3. *Every interior point $x \in (D^{d-1})^k$ is contained in the interior of a cell corresponding to a symbolic matrix $A \in \mathbf{S}(d - 2, k)$. If the stabilizer of x is non-trivial, this cell is unique.*

Proof. The proof is by induction on k . If $k = 1$, there is only the zero matrix and a matrix with one non-zero entry. It is easy to see that the result holds in this case. Now let $x = (x_1, \dots, x_k) \in (D^{d-1})^k$ with $k > 1$. If $x_1 \in D^1$, we can use induction on $x' = (x_2, \dots, x_k)$ to get a matrix $A' \in \mathbf{S}(d - 2, k - 1)$ so that $x' \in \sigma_{A'}$, and which is unique if the stabilizer of x' is non-trivial. Then $x \in \sigma_A$, where A is the matrix obtained from A' by adding a zero column to the left of A' . Note that the stabilizer of x is the stabilizer of x' , and the uniqueness applies if it is non-trivial. If $x_1 \notin D^1$, we can find an $A \in SO(d - 2)$ such that $Ax_1 \in D_+^2$. Now let $p : D^{d-1} \rightarrow D^{d-2}$ be projection to the last $d - 2$ coordinates and consider the point $x' = (p(Ax_2), \dots, p(Ax_k))$. By induction, we can find a symbolic matrix $A' \in \mathbf{S}(d - 3, k - 1)$ with $x' \in \sigma_{A'}$, and the matrix is unique if the stabilizer of x' is non-trivial (which implies that the stabilizer of x is non-trivial). Then x is in the cell σ_A , where

$$A = \begin{pmatrix} + & * \\ 0 & A' \end{pmatrix},$$

and the cell is unique if the stabilizer of x is non-trivial. \square

Lemma 6.4. *Let $x \in (D^{d-1})^k$ be an interior point with trivial stabilizer. Then x is contained in a cell σ_A where the last two rows of A are of the form*

$$(3) \quad \left(\begin{array}{cccccccc} 0 & \cdots & 0 & + & * & \cdots & * \\ 0 & \cdots & 0 & 0 & * & \cdots & * \end{array} \right).$$

Furthermore, no two such matrices have interior points in common.

Proof. We know from the previous lemma that x is contained in some matrix, and since the stabilizer of x is trivial, the second but last row has to be non-zero. In particular, there has to be a $+$ in that row. Since the $*$ symbolize any possible entry, including 0, x will be in a cell corresponding to such a matrix.

To see that no two such matrices have interior points in common, note that in the column which has a $+$ in second but last row, interior points $y \in D_+^{d-2}$ satisfy $y_{d-2} > 0$, and this is the first column, for which this occurs. \square

Notice that the matrices in (2) are not in the form of Lemma 6.4.

Define

$$\mathbf{S}_c(d-2, k) = \{A \in \mathbf{S}(d-2, k) \mid \text{The last two rows are of the form (3) or 0}\}$$

An equivariant relative cell decomposition of $((D^{d-1})^k, \partial(D^{d-1})^k)$ is therefore given by the cells σ_A , where $A \in \mathbf{S}_c(d-2, k)$.

7. THE BOUNDARY OPERATOR FOR THE CELL DECOMPOSITION

The equivariant cell decomposition described in the previous section gives a relative CW-structure on $(X_d^k, \partial X_d^k)$, where $X_d^k = (D^{d-1})^k/SO(d-2)$ and $\partial X_d^k = \partial(D^{d-1})^k/SO(d-2)$. The cells are simply of the form D_A for $A \in \mathbf{S}_c(d-2, k)$, where each D_A is a product of discs or halfdiscs.

Notice that the boundary of each factor D^i is attached to ∂X_d^k , and each factor D_+^i is attached to ∂X_d^k and to the same cell with the factor replaced by D^{i-1} . So the boundary of a cell D_A is contained in ∂X_d^k together with cells $D_{A'}$, where the A' are obtained from A by replacing a $+$ by a 0.

This needs to be made slightly more precise. If a matrix A contains a submatrix $\begin{pmatrix} + & * \\ 0 & + \end{pmatrix}$, replacing the $+$ in the upper left corner by 0 leads to a matrix with submatrix $\begin{pmatrix} 0 & * \\ 0 & + \end{pmatrix}$, which is not an element of \mathbf{S}_c . However, up to elements of $SO(d-2)$ we get that the corresponding boundary points are in the cell containing the submatrix $\begin{pmatrix} 0 & + \\ 0 & 0 \end{pmatrix}$. The dimension of this cell is the dimension of the original cell -2 . In particular, it will not occur in the boundary operator.

If the matrix A contains a submatrix $\begin{pmatrix} + & * \\ 0 & 0 \end{pmatrix}$, replacing the $+$ in the upper left corner by 0 leads to the matrix with submatrix $\begin{pmatrix} 0 & + \\ 0 & 0 \end{pmatrix}$, but the change from $*$ to $+$ in the right upper corner means that the attaching is done twice, so the coefficient in the boundary operator is 0 or 2, depending on orientation considerations.

Finally, if the last non-zero row of the matrix A is of the form $(0 \ \cdots \ 0 \ +)$, replacing this row by the zero row gives a matrix $A' \in \mathbf{S}_c$, and the corresponding coefficient in the boundary operator is ± 1 .

For $i = 1, \dots, d-3$, define $\mathbf{S}^{(i)}(d-2, k)$ to consist of those matrices $A \in \mathbf{S}_c(d-2, k)$ for which the i -th row is $(0 \ \cdots \ 0)$ or $(0 \ \cdots \ 0 \ +)$.

Let $(X_{(i),d}^k, \partial X_d^k)$ be the relative CW-complex consisting of the cells corresponding to $\mathbf{S}^{(i)}(d-2, k)$.

Lemma 7.1. *The relative CW-complex $(X_{(i),d}^k, \partial X_d^k)$ collapses to $(\partial X_d^k, \partial X_d^k)$ for all $i = 1, \dots, d-3$.*

Proof. The proof is by induction on i . For $i = 1$, we only have two cells, corresponding to the zero matrix and the matrix whose only non-zero entry is a $+$. By the discussion above, the two cells form an elementary collapse in the sense of [1, §4], and the result follows.

For $i > 1$ we show that $(X_{(i),d}^k, \partial X_d^k)$ collapses to $(X_{(i-1),d}^k, \partial X_d^k)$. Note that if $i > k$, then $X_{(i),d}^k = X_{(i-1),d}^k$ and there is nothing to show. So assume $i \leq k$ and let $A \in \mathbf{S}^{(i)}(d-2, k) - \mathbf{S}^{(i-1)}(d-2, k)$. Then the $(i-1)$ -th row of A is non-zero, and different from $(0 \ \cdots \ 0 \ +)$. The i -th row is either $(0 \ \cdots \ 0)$ or $(0 \ \cdots \ 0 \ +)$, and the two possibilities form an elementary collapse. By collapsing these pairs in the order of decreasing dimension, we see that $(X_{(i),d}^k, \partial X_d^k)$ collapses to $(X_{(i-1),d}^k, \partial X_d^k)$. The result follows. \square

Corollary 7.2. *For $k < d-2$, the pair $(X_d^k, \partial X_d^k)$ is m -connected for all $m \geq 0$.*

Proof. The cells to form the relative CW-complex are in one-to-one correspondence with $\mathbf{S}_c(d-2, k)$, but since $k < d-2$, we get $\mathbf{S}_c(d-2, k) = \mathbf{S}^{(d-3)}(d-2, k)$. The result thus follows from Lemma 7.1. \square

In the next result the condition $a_0(\ell) = 1$ is needed to avoid the case $\mathcal{M}_d(\ell) = \emptyset$.

Proposition 7.3. *Let $\ell \in \mathbb{R}^n$ be a generic length vector with $a_0(\ell) = 1$, and $d \geq n \geq 3$. Then $\mathcal{M}_d(\ell)$ is contractible.*

Proof. Let $\tilde{F} : \mathcal{C}_d(\ell) \rightarrow \mathbb{R}$ the $SO(d-1)$ -invariant Morse-Bott function from Proposition 6.1, $F : \mathcal{M}_d(\ell) \rightarrow \mathbb{R}$ the induced function and let

$$\emptyset = \mathcal{M}^0 \subset \mathcal{M}^1 \subset \cdots \subset \mathcal{M}^m = \mathcal{M}_d(\ell)$$

be a filtration such that $\mathcal{M}^m = F^{-1}((-\infty, a_m])$ for some sequence of regular values of \tilde{F} such that $\mathcal{M}^m - \mathcal{M}^{m-1}$ contains exactly one critical point.

By Morse-Bott theory, \mathcal{M}^m is homotopy equivalent to $\mathcal{M}^{m-1} \cup X_d^k$, where X_d^k is attached to \mathcal{M}^{m-1} along ∂X_d^k , and k is such that $k(d-1)$ is the index of the critical point in $\mathcal{M}^m - \mathcal{M}^{m-1}$. Since $k \leq n-3$, we get $k < d-2$, and \mathcal{M}^m has the same homotopy type as \mathcal{M}^{m-1} , provided $k \geq 1$. As there is a unique minimum for F by the perfectness of the map F_3 in Proposition 6.1, we get that \mathcal{M}^1 has the homotopy type of a point, and all other critical points have index bigger than 0. \square

Of course, by [21] these spaces are homeomorphic to a disc.

If $k \geq d-2$, then $\mathbf{S}^{(d-3)}(d-2, k) \neq \mathbf{S}_c(d-2, k)$. A matrix $A \in \mathbf{S}_c(d-2, k) - \mathbf{S}^{(d-3)}(d-2, k)$ has to have at least one $*$ in its last row, and therefore it has $(d-1)(d-2)/2$ non-zero entries. It follows that D_A has at least dimension $(d-1)(d-2)/2 + k$.

Lemma 7.4. *Let $k \geq d-2 \geq 2$, then $(X_d^k, \partial X_d^k)$ is $((d-1)(d-2)/2 + k - 1)$ -connected, but not $((d-1)(d-2)/2 + k)$ -connected.*

Proof. We look at the connectedness of the pair $(X_d^k, X_{(d-3),d}^k)$, which is obtained by attaching cells corresponding to $A \in \mathbf{S}_c(d-2, k) - \mathbf{S}^{(d-3)}(d-2, k)$. There is only one cell D_A which has dimension at most $(d-1)(d-2)/2 + k$, namely the one corresponding to

$$A = \begin{pmatrix} 0 & \cdots & 0 & + & * & \cdots & * \\ \vdots & & & \ddots & \ddots & \ddots & \vdots \\ \vdots & & & & \ddots & + & * \\ 0 & \cdots & \cdots & & 0 & * \end{pmatrix},$$

and if $k > d-2$, there is only one cell with dimension $(d-1)(d-2)/2 + k + 1$, namely the one corresponding to

$$A' = \begin{pmatrix} 0 & \cdots & 0 & + & * & * & \cdots & \cdots & * \\ \vdots & & & 0 & 0 & + & \ddots & & \vdots \\ \vdots & & & & & \ddots & \ddots & \ddots & \vdots \\ \vdots & & & & & & \ddots & + & * \\ 0 & \cdots & \cdots & & & & 0 & * \end{pmatrix}.$$

With the discussion on boundaries given above, we get $H_m(X_d^k, X_{(d-3),d}^k; \mathbb{Z}/2\mathbb{Z}) = \mathbb{Z}/2\mathbb{Z}$ for $m = (d-1)(d-2)/2 + k$. Using Corollary 7.1, the result follows. \square

Theorem 7.5. *Let $\ell \in \mathbb{R}^n$ be a generic length vector with $a_0(\ell) = 1$, and $d \geq 3$. Then $\mathcal{M}_d(\ell)$ is $((d-1)(d-2)/2 + d-3)$ -connected. Furthermore $\mathcal{M}_{n-1}(\ell)$ is homotopy equivalent to the sphere of dimension $n(n-3)/2$.*

Proof. The proof begins in the same way as the proof of Proposition 7.3, with the filtration

$$\emptyset = \mathcal{M}^0 \subset \mathcal{M}^1 \subset \cdots \subset \mathcal{M}^m = \mathcal{M}_d(\ell).$$

As long as the index of the critical point is $k(d-1)$ with $k < d-2$, no new homology occurs, but if $k \geq d-2$, something can happen, but by Lemma 7.4 the new \mathcal{M}^{i+1} is still $((d-1)(d-2)/2 + d-3)$ -connected.

If $d = n-1$, the case $k \geq d-2 = n-3$ only appears once, with the absolute maximum of the function. In that case only one cell of dimension $(d-1)(d-2)/2 + d-2$ is attached to a contractible space. Hence, up to homotopy, we get a sphere of dimension $(n-2)(n-3)/2 + n-3 = n(n-3)/2$. \square

As mentioned in the introduction, the last result can be improved to a homeomorphism between $\mathcal{M}_{n-1}(\ell)$ and the sphere. To see this, note that the closure of the space Ω_{n-1} of [21] can be identified with $\mathcal{M}_n(\ell)$ for $\ell \in \mathbb{R}^n$ by sending a linkage configuration to the distances between the points. By [21, Thm.1], this space is homeomorphic to a disc of dimension $n(n-3)/2$, and the boundary points correspond to those points $x \in \mathcal{M}_n(\ell)$ whose rank is at most $n-2$.

The space $\mathcal{M}_{n-1}(\ell)$ is now obtained by doubling $\mathcal{M}_n(\ell)$ along the boundary, compare Lemma 2.4 and also the proof of [12, Thm.C].

8. BASIC HOMOLOGICAL PROPERTIES OF $(X_d^k, \partial X_d^k)$

Let us denote the cellular chain complex for the pair $(X_d^k, \partial X_d^k)$ by C_* , that is, the chain complex freely generated by the matrices of $\mathbf{S}_c(d-2, k)$. This contains the subcomplex for the pair $(X_{(d-3),d}^k, \partial X_d^k)$, which we denote by C_*^0 , and which is freely generated by the matrices of $\mathbf{S}^{(d-3)}(d-2, k)$. By Lemma 7.1 we get that $H_*(C_*^0) = 0$, and hence

$$H_*(X_d^k, \partial X_d^k) = H_*(D_*),$$

where $D_* = C_*/C_*^0$ is freely generated by matrices whose last two rows are of the form (3), and where the last row is non-zero.

Let us assume that $k \geq d-2$, so that $D_* \neq 0$.

Notice that we can write D_* as a direct sum of chain complexes

$$(4) \quad D_* = \bigoplus_{j=1}^{k-d+3} D_*^k(j),$$

where $D_*^k(j)$ is generated by those matrices which have $(k-d+4) - j$ columns² containing just *. In particular, $D_*^k(1)$ has only one generator, corresponding to the matrix

$$A = \begin{pmatrix} + & * & \cdots & * & \cdots & * \\ 0 & \ddots & \ddots & \vdots & & \vdots \\ \vdots & \ddots & + & * & \cdots & * \\ 0 & \cdots & 0 & * & \cdots & * \end{pmatrix}$$

The dimension of the cell D_A is therefore

$$k + (k-d+3)(d-2) + \frac{(d-2)(d-3)}{2} = k(d-1) - \frac{(d-2)(d-3)}{2}.$$

The top-dimensional cell in $D_*^k(j)$ corresponds to a matrix of the form

$$\begin{pmatrix} + & * & \cdots & * & \cdots & \cdots & * & * & \cdots & * \\ 0 & \ddots & \ddots & \vdots & & & \vdots & \vdots & & \vdots \\ \vdots & \ddots & + & * & \cdots & \cdots & * & \vdots & & \vdots \\ 0 & \cdots & 0 & 0 & \cdots & 0 & + & * & \cdots & * \\ 0 & \cdots & & & & \cdots & 0 & * & \cdots & * \end{pmatrix}$$

while the minimal-dimensional cell corresponds to a matrix of the form

$$\begin{pmatrix} 0 & \cdots & 0 & + & * & \cdots & * & \cdots & * \\ \vdots & & \vdots & 0 & \ddots & \ddots & \vdots & & \vdots \\ \vdots & & \vdots & \vdots & \ddots & + & * & \cdots & * \\ 0 & \cdots & 0 & 0 & \cdots & 0 & * & \cdots & * \end{pmatrix}$$

So all the cells in $D_*^k(j)$ have dimension between

$$k(d-1) - \frac{(d-2)(d-3)}{2} - (j-1)(d-2) \text{ and } k(d-1) - \frac{(d-2)(d-3)}{2} - 2(j-1).$$

²The number is chosen so that $D_*^k(1)$ has exactly one generator, while $D_*^k(k-d+3)$ has the most generators.

If we consider the complexes $D_*^k(j)$ with coefficients in $\mathbb{Z}/2\mathbb{Z}$, we get that every boundary is zero. This follows from the discussion at the beginning of Section 7, as there always is a column containing only $*$.

Even with coefficients in \mathbb{Z} we can obtain some basic results on the homology of $\mathcal{M}_d(\ell)$.

Proposition 8.1. *Let $\ell \in \mathbb{R}^n$ be a generic length vector with $a_0(\ell) = 1$, let $d \geq 4$ and let $n \geq d + 1$. Then*

$$H_{\mathbf{d}_d^n}(\mathcal{M}_d(\ell); \mathbb{Z}) = \mathbb{Z} \quad \text{and} \quad H_{\mathbf{d}_d^n - 1}(\mathcal{M}_d(\ell); \mathbb{Z}) = 0$$

Recall that \mathbf{d}_d^n denotes the dimension of $\mathcal{M}_d(\ell)$.

Proof. Let $\tilde{F} : \mathcal{C}_d(\ell) \rightarrow \mathbb{R}$ be the $SO(d-1)$ -invariant Morse function from Proposition 6.1, $F : \mathcal{M}_d(\ell) \rightarrow \mathbb{R}$ the induced function and let

$$\emptyset = \mathcal{M}^0 \subset \mathcal{M}^1 \subset \dots \subset \mathcal{M}^m = \mathcal{M}_d(\ell)$$

be a filtration such that $\mathcal{M}^m = F^{-1}((-\infty, a_m])$ for some sequence of regular values of \tilde{F} such that $\mathcal{M}^m - \mathcal{M}^{m-1}$ contains exactly one critical point.

Notice that \tilde{F} has only one critical manifold of index $(n-3)(d-1)$, which is the absolute maximum. Since \mathcal{M}^{m-1} has the homotopy type of a CW-complex with lower dimensional cells, we get $H_q(\mathcal{M}^{m-1}; \mathbb{Z}) = 0$ for $q \geq \mathbf{d}_d^n - 1$. Now $\mathcal{M}_d(\ell)$ is, up to homotopy, obtained from \mathcal{M}^{m-1} by attaching the cells from $((X_d^{n-3}, \partial X_d^{n-3}))$. Only one cell has dimension $\geq \mathbf{d}_d^n - 1$, and this cell has dimension \mathbf{d}_d^n . The result follows. \square

Theorem 8.2. *Let $\ell \in \mathbb{R}^n$ be a generic length vector with $a_0(\ell) = 1$, let $d \geq 4$ and let $n \geq d + 2$. Then $\mathcal{M}_d(\ell)$ does not satisfy Poincaré duality with coefficients in $\mathbb{Z}/2\mathbb{Z}$. In particular, $\mathcal{M}_d(\ell)$ is not a topological manifold, with or without boundary.*

Proof. First notice that $\mathcal{M}_d(\ell)$ cannot be a manifold with non-empty boundary, as by Proposition 8.1 $H_{\mathbf{d}_d^n}(\mathcal{M}_d(\ell)) \neq 0$.

Let us use the same filtration as in the previous proof.

We will distinguish the cases $d = 4$ and $d \geq 5$. Let us first assume that $d \geq 5$. Then \mathcal{M}^{m-1} has the homotopy type of a CW-complex with cells of dimension at most $(n-4)(d-1) - \frac{(d-2)(d-3)}{2}$.

As $n \geq d+2$, we get that $H_{\mathbf{d}_d^n - 2}(D^{n-3}(2); \mathbb{Z}/2\mathbb{Z}) = \mathbb{Z}/2\mathbb{Z}$, which corresponds to the maximal cell for $D^{n-3}(2)$. As $d \geq 5$, we get $\mathbf{d}_d^n - 2 - ((n-4)(d-1) - \frac{(d-2)(d-3)}{2}) \geq 2$, so

$$H_{\mathbf{d}_d^n - 2}(\mathcal{M}_d(\ell); \mathbb{Z}/2\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}.$$

But $H_2(\mathcal{M}_d(\ell); \mathbb{Z}/2\mathbb{Z}) = 0$ by Theorem 7.5, so Poincaré duality cannot hold.

Now consider the case $d = 4$. Since $\mathbf{d}_d^n - 2 - ((n-4)(d-1) - \frac{(d-2)(d-3)}{2}) = 1$ now, it is not clear whether $H_{\mathbf{d}_d^n - 2}(\mathcal{M}_d(\ell); \mathbb{Z}/2\mathbb{Z}) \neq 0$.

But let c be the number of critical manifolds of index $3(n-4)$. By Proposition 5.4, we get $c = 1 + a_1(\ell) - a_{n-3}$. Now $a_{n-3}(\ell) \leq 1$, and if $a_{n-3}(\ell) = 1$, then $a_1(\ell) = n-3$. As $n \geq 6$, we get $c \geq 2$, unless $a_1(\ell) = 0$ in which case $c = 1$.

Let us first consider the case $c \geq 2$. In that case the top-dimensional non-zero homology group of \mathcal{M}^{m-1} is in degree $3(n-4) - 1$, and the rank of this homology group is c . Attaching one cell of dimension $3(n-4)$ cannot kill this homology group, therefore $H_{3(n-4)-1}(\mathcal{M}_4(\ell); \mathbb{Z}/2\mathbb{Z}) \neq 0$. But by Theorem 7.5 we have $H_3(\mathcal{M}_4(\ell); \mathbb{Z}/2\mathbb{Z}) = 0$, so Poincaré duality cannot hold.

It remains to consider the case $c = 1$. In that case $\mathcal{S}_1^m(\ell) = \emptyset$ (where m is chosen so that ℓ is m -dominated), which uniquely determines the chamber of ℓ . In fact, we can assume that

$$\ell = (1, \dots, 1, n-2).$$

By Proposition A.1 we get that $\mathcal{M}_4(\ell) \approx \Sigma_3^{n-1}$, the shape space defined in the appendix. But this space is known to not satisfy Poincaré duality, see [17, §4, §5]. In fact, the homology calculations in [17] give the same contradiction as above. \square

9. HOMOLOGY OF $(X_d^k, \partial X_d^k)$

In this section we want to improve on the homology calculations of $H_*(X_d^k, \partial X_d^k)$. Let us begin with the case $d = 4$. In that case

$$D_*^k(j) = (\mathbb{Z}, 3k - 1 - 2(j-1)),$$

where we use the notation (G, n) for the graded group whose only non-zero degree is $n \in \mathbb{Z}$, in which case the entry is the abelian group G .

It follows that for $k \geq 1$ we get

$$H_*(X_4^k, \partial X_4^k) = \bigoplus_{j=1}^{k-1} (\mathbb{Z}, 3k - 1 - 2(j-1)).$$

In the case $d \geq 5$ we have to analyze the boundary operator more carefully. This is done by following the methods of [17, §4]. Let us take a closer look at $d = 5$. The matrices appearing for the generators of $D_*^k(j)$ are of the form

$$\begin{pmatrix} 0 & \cdots & 0 & + & * & \cdots & * & * & * & \cdots & * \\ 0 & \cdots & 0 & 0 & \cdots & \cdots & 0 & + & * & \cdots & * \\ 0 & & & \cdots & & & 0 & 0 & * & \cdots & * \end{pmatrix}$$

A typical boundary is of the form

$$\partial \begin{pmatrix} + & * & * & * \\ 0 & 0 & + & * \\ 0 & 0 & 0 & * \end{pmatrix} = \begin{pmatrix} 0 & + & * & * \\ 0 & 0 & + & * \\ 0 & 0 & 0 & * \end{pmatrix} + \begin{pmatrix} 0 & - & * & * \\ 0 & 0 & + & * \\ 0 & 0 & 0 & * \end{pmatrix}$$

The second matrix on the right-hand side comes from the fact that we write $D^2 = D_+^2 \cup D_-^2$, but our symbolic matrices require a $+$ and not a $-$. Let $A \in SO(3) \leq SO(4)$ be the diagonal matrix which has -1 in the second and fourth entry, and 1 in the first and third entry. Then $A(D_-^2) = D_+^2$. This means we get the same matrix on the right side twice. To work out the exact coefficients, we need to take a closer look at orientations.

Recall that the matrices stand for products of discs D^i or D_+^i , and every non-zero entry corresponds to one dimension. To choose an orientation, we choose the standard orientation of the discs D^i . We can actually think of every non-zero entry in the matrix coming with a basis vector into that dimension, and by picking an order of the entries in the matrix we get the orientation.

Let us go back to the example above. The matrix $A \in SO(3)$ used to turn D_-^2 into D_+^2 changes the orientation of D_+^2 . It also changes the orientation of the next factor, which is a D_+^3 . But for the final factor D^4 , two basis elements are changed, so there is no impact on the orientation. Since we had two changes of orientations, we see that

$$\partial \begin{pmatrix} + & * & * & * \\ 0 & 0 & + & * \\ 0 & 0 & 0 & * \end{pmatrix} = \pm 2 \begin{pmatrix} 0 & + & * & * \\ 0 & 0 & + & * \\ 0 & 0 & 0 & * \end{pmatrix}$$

Notice that adding extra factors of D^4 two the right has no impact on the signs. Adding extra factors of D^2 however does change the signs. So to work out the homology of $D_*^k(j)$, we can ignore the last two rows of the matrix.

The remaining $d-2$ rows all have to start with a $+$. Let us describe the remaining matrices using sequences of decreasing numbers, compare [17, §4]. We can encode the matrix by a sequence of numbers (k_1, \dots, k_m) with

$$k_1 > k_2 > \dots > k_m \geq 1$$

where each number k_i stands for the number of non-zero entries in the i -th row.

Let $E_*(m, j)$ be the chain complex freely generated by such sequences (k_1, \dots, k_m) where $k_1 \leq m + j - 1$, and we say that the sequence (k_1, \dots, k_m) has degree $k_1 + \dots + k_m - m(m+1)/2$. The boundary is given by

$$\partial(k_1, \dots, k_m) = \sum_{j=1}^m (-1)^{k_1 + \dots + k_{j-1}} (1 + (-1)^{k_j}) (k_1, \dots, k_j - 1, \dots, k_m).$$

where a sequence (k'_1, \dots, k'_m) is interpreted as 0 if $k'_i = k'_{i+1}$ for some $i \in 1, \dots, m-1$ or if $k'_m = 0$.

Remark 9.1. The sign $(-1)^{k_1 + \dots + k_{j-1}}$ comes from the following: Each non-zero entry in the symbolic matrix spans a dimension, but only the entries with a $+$ have a non-zero boundary. If we order the basis for the orientation by starting with the first row on the left, the $+$ is at the 0-th position. Similarly, the $+$ in the second row is in the k_1 -th position, and so on.

It follows that, up to possibly a sign which has no impact on the homology,

$$(5) \quad D_*^k(j) = E_{*-u}(d-4, j),$$

where $u = (d-1)k - (d-2)(d-3)/2 - (d-2)(j-1)$.

Let us take a look at the case $d = 5$. Then $E_*(1, j)$ is generated by (k) , where $k \leq j$, the boundary maps are alternating between 2 and 0, and $\partial(2) = 2(1)$. We thus get

Lemma 9.2. *Let $j \geq 1$. Then*

$$H_q(E_*(1, j)) = \begin{cases} \mathbb{Z} & j \text{ odd, } q = j - 1 \\ 0 & q \text{ odd, or } q \geq j \\ \mathbb{Z}/2\mathbb{Z} & q \text{ even} \end{cases}.$$

To understand the homology of $E_*(m, j)$ for $m \geq 2$, notice that we can think of this complex as the total complex of a double complex E_{**} , where the horizontal grading measures the first row, and the vertical grading the remaining rows. We can

therefore think of $E_*(m, j)$ as the total complex of the sequence of chain complexes with chain maps

$$(6) \quad E_*(m-1, 1) \xrightarrow{1+(-1)^{m-1}} E_*(m-1, 2) \xrightarrow{1+(-1)^{m-2}} \dots \xrightarrow{1+(-1)^{m-j+1}} E_*(m-1, j).$$

Notice that every second map is 0, so that the total complex is just a direct sum of sequences

$$E_*(m-1, i) \xleftarrow{2} E_*(m-1, i+1).$$

Using this and the particular form of the boundary in $E_*(m, j)$ one can show that the homology of $E_*(m, j)$ only contains direct summands of \mathbb{Z} and $\mathbb{Z}/2\mathbb{Z}$. One should compare this with the results in [17, §4, §5], where closed formulas for the number of such summands in the homology of similar chain complexes are given. As these closed formulas are not that enlightening, and since we need to enter the homology of $(X_d^k, \partial X_{d-1}^k)$ into another spectral sequence coming from the filtration $(\mathcal{M}^k)_{k \geq 0}$, we will abandon torsion and look instead at homology with coefficients in \mathbb{Q} .

Lemma 9.2 then reduces to

$$H_*(E_*(1, j); \mathbb{Q}) = \begin{cases} 0 & j \text{ even} \\ (\mathbb{Q}, j-1) & j \text{ odd} \end{cases}.$$

To describe the rational homology of $E_*(m, j)$ for $m \geq 2$ we want to give concrete generators, and then show that they span the homology. Let us begin with m even, that is, $m = 2n$ for some $n \geq 1$.

Let $j + m - 1 \geq k_1 > \dots > k_n \geq 2$ be a sequence of even numbers. Then

$$(k_1, k_1 - 1, k_2, k_2 - 1, \dots, k_n, k_n - 1)$$

is easily seen to be a cycle in $E(2n, j)$. Furthermore, no non-zero integer multiple can be a boundary, as only sequences which have a term $(k_i + 1, k_i - 1)$ in them could have this sequence in their boundary. But since $k_i + 1$ is odd, the boundary formula has a factor $1 + (-1)^{k_i+1} = 0$. In particular, such cycles span a \mathbb{Z} factor in $H_*(E_*(2n, j))$. It is also easy to see that the degree of this cycle is a multiple of 4.

For $m = 2n + 1$ we can look at the sequence

$$(k_0, k_1, k_1 - 1, k_2, k_2 - 1, \dots, k_n, k_n - 1)$$

where the k_i are as before for $i \geq 1$, and $j + m - 1 \geq k_0 > k_1$. For this to be a cycle, we need k_0 to be odd. But if $k_0 + 1 \leq j + m - 1$, we get this to be a (rational) boundary. To obtain a \mathbb{Z} factor in $H_*(E_*(2n + 1, j))$, we therefore need $k_0 = j + m - 1$. As m is odd, this is only possible if j is odd. In this case, notice that the degree of this cycle is $j - 1 + 4i$ for some $i \geq 0$.

Proposition 9.3. *Let $n \geq 1$ and $j \geq 1$. Then $H_*(E_*(2n, j); \mathbb{Q})$ has a basis given by elements*

$$(k_1, k_1 - 1, k_2, k_2 - 1, \dots, k_m, k_m - 1)$$

where $k_i = 2(n + 1 - i) + j_i$ for $i = 1, \dots, n$, where $j_1 \geq j_2 \geq \dots \geq j_n \geq 0$ is a sequence of even numbers with $j_1 + 2n \leq j + 2n - 1$. The degree of $(k_1, k_1 - 1, k_2, k_2 - 1, \dots, k_n, k_n - 1)$ is $2(j_1 + \dots + j_n)$.

Furthermore, $H_*(E_*(2n+1, j); \mathbb{Q}) = 0$ for j even, and $H_*(E_*(2n+1, j); \mathbb{Q})$ for j odd has a basis given by elements

$$(k_0, k_1, k_1 - 1, k_2, k_2 - 1, \dots, k_n, k_n - 1)$$

with the k_i as above, and $k_0 = j + 2n$. The degree of $(k_0, k_1, k_1 - 1, k_2, k_2 - 1, \dots, k_n, k_n - 1)$ is $(j - 1) + 2(j_1 + \dots + j_n)$.

Proof. The proof is by induction. Let us first show that the statement for $2n - 1$ implies the statement for $2n$. We get that the chain complex $E_*(2n, j)$ is the total complex of the sequence (6). Now all the chain complexes $E_*(2n - 1, i)$ with i even have 0 as their homology, so the homology of $E_*(2n, j)$ is the direct sum of the homologies of $E_*(2m - 1, i)$ with i odd. The basis elements for $H_*(E_*(2m - 1, i); \mathbb{Q})$ are then of the form

$$(i + 2n - 2, k_1, k_1 - 1, \dots, k_{n-1}, k_{n-1} - 1)$$

with $i \leq j$ odd by the induction assumption. The way we think of $E_*(2n, j)$ as a double complex means these generators correspond to

$$(i + 2n - 1, i + 2n - 2, k_1, k_1 - 1, \dots, k_{n-1}, k_{n-1} - 1).$$

But this gives exactly the statement for the rational homology of $E_*(2n, j)$. Notice that this also works for $n = 1$.

It remains to show that the statement for $2n$ implies the statement for $2n + 1$.

Again we use the sequence (6). The condition that $k_1 \leq i + 2n - 1$, implies that for i odd the homologies of $E_*(2n, i)$ and $E_*(2n, i + 1)$ have the same basis. Furthermore, in (6) we get for i odd terms of the form

$$E_*(2n, i) \xleftarrow{2} E_*(2n, i + 1)$$

which induce isomorphisms on rational homology. In particular, for j even all homology vanishes. For j odd we are left with

$$H_*(E_*(2n + 1, j); \mathbb{Q}) \cong H_{*-j}(E_*(2n, j); \mathbb{Q}),$$

and because of the way the double complex structure of $E_*(2n + 1, j)$ is formed, we see that the basis is represented by elements

$$(j + 2n, k_1, k_1 - 1, k_2, k_2 - 1, \dots, k_n, k_n - 1).$$

The statement about the degrees of these basis elements is easy to see. \square

Definition 9.4. Let $m \geq 1$. Then define

$$\nabla_m = \{(j_1, \dots, j_m) \in \mathbb{Z}^m \mid j_1 \geq j_2 \geq \dots \geq j_m \geq 0\}.$$

Also, if $(j_1, \dots, j_m) \in \nabla_m$, we define

$$|(j_1, \dots, j_m)| = j_1 + j_2 + \dots + j_m$$

and

$$\|(j_1, \dots, j_m)\| = 2j_1 + 1.$$

Also, for $m = 0$ we let $\nabla_0 = \{()\}$, where we think of $()$ as a point with $|()| = 0$ and $\|()\| = 1$.

We write elements of ∇_m as $\mathbf{j} = (j_1, \dots, j_m)$. Notice that every $\mathbf{j} \in \nabla_m$ produces a generator in the homology of $E(2m, j)$, provided that $j \geq \|\mathbf{j}\|$, whose degree is $4\|\mathbf{j}\|$, and a generator in the homology of $E(2m+1, 2i+1)$, provided that $2i+1 \geq \|\mathbf{j}\|$, whose degree is $2i+4\|\mathbf{j}\|$.

For $j \geq 1$ and $m \geq 0$ let

$$\nabla_m(j) = \{\mathbf{j} \in \nabla_m \mid \|\mathbf{j}\| \leq j\}.$$

We denote the Poincaré polynomial of the pair $(X_d^k, \partial X_d^k)$ by $P_d^k(t)$, that is,

$$P_d^k(t) = b_0(X_d^k, \partial X_d^k) + b_1(X_d^k, \partial X_d^k)t + \dots + b_n(X_d^k, \partial X_d^k)t^n$$

where n is the dimension of X_d^k and $b_j(X_d^k, \partial X_d^k)$ is the \mathbb{Z} -rank of $H_j(X_d^k, \partial X_d^k)$ for all $j = 0, \dots, n$.

Theorem 9.5. *Let $m \geq 0$ and $k \geq 1$. Then*

$$P_{2m+4}^k(t) = t^{k+(m+1)(2m+3)} \sum_{\mathbf{j} \in \nabla_m(k-2m-1)} t^{4\|\mathbf{j}\|} \frac{t^{(2m+2)(k-2m-\|\mathbf{j}\|)} - 1}{t^{2m+2} - 1}$$

and

$$P_{2m+5}^k(t) = t^{u(k,m)} \sum_{\mathbf{j} \in \nabla_m(k-2m-2)} t^{4\|\mathbf{j}\|} \frac{t^{4(m+1)\lfloor \frac{k-2m-\|\mathbf{j}\|}{2} \rfloor} - 1}{t^{4(m+1)} - 1}$$

where

$$u(k, m) = (2m+4)k - (2m+3)(m+1) - 4(m+1)\lfloor \frac{k-2m-3}{2} \rfloor.$$

Here $\lfloor x \rfloor = \max\{n \in \mathbb{Z} \mid n \leq x\}$.

Proof. The proof is now merely an organisation of our previous results, using $d = 2m+4$ or $2m+5$. By (4) the homology of $(X_d^k, \partial X_d^k)$ splits into summands, which by (5) come from $E_*(2m, j)$ or $E_*(2m+1, j)$ shifted by

$$v(d, k, j) = (d-1)k - (d-2)(d-3)/2 - (d-2)(j-1),$$

and where $j = 1, \dots, k-d+3$.

Let $d = 2m+4$. Using Proposition 9.3 we see that each $\mathbf{j} \in \nabla_m(k-2m-1)$ produces a homology generator, and in fact for each $j = 1, \dots, k-2m-1$ with $j \geq \|\mathbf{j}\|$. The degree of such a generator is $4\|\mathbf{j}\| + v(2m+4, k, j)$, so the degrees vary from

$$4\|\mathbf{j}\| + (2m+3)k - (m+1)(2m+1) - 2(m+1)(\|\mathbf{j}\| - 1)$$

down to

$$4\|\mathbf{j}\| + v(2m+4, k, k-2m-1) = 4\|\mathbf{j}\| + k + (m+1)(2m+3)$$

in steps of $2(m+1)$. Using

$$1 + t^{2m+2} + \dots + t^{(2m+2)(k-2m-1-\|\mathbf{j}\|)} = \frac{t^{(2m+2)(k-2m-\|\mathbf{j}\|)} - 1}{t^{2m+2} - 1},$$

we get the result.

The case $d = 2m+5$ is very similar, each $\mathbf{j} \in \nabla_m(k-2m-1)$ produces a homology generator, but only for each odd $j = 1, \dots, k-2m-2$ with $j \geq \|\mathbf{j}\|$, and with degree $(j-1) + 4\|\mathbf{j}\| + v(2m+5, k, j)$. A similar argument as in the even case gives the stated result. Note that $2\lfloor \frac{n-1}{2} \rfloor + 1$ is the largest odd number not bigger than

n , and the degree increase for each \mathbf{j} is $4(m+1)$ because we only consider odd numbers between $\|\mathbf{j}\|$ and $k-2m-2$. \square

For small values of m the sets ∇_m have a very simple form, so we collect the Poincaré polynomials in these special cases in the next corollary.

Corollary 9.6. *For $k \geq 2$ we have*

$$P_4^k(t) = t^{k+3} \sum_{i=0}^{k-2} t^{2i}.$$

For $k \geq 3$ we have

$$P_5^k(t) = t^{4k-4\lfloor \frac{k-3}{2} \rfloor - 3} \sum_{i=0}^{\lfloor \frac{k-3}{2} \rfloor} t^{4i}.$$

For $k \geq 4$ we have

$$P_6^k(t) = t^{k+10} Q_{k-4}(t^4).$$

For $k \geq 5$ we have

$$P_7^k(t) = t^{6k-8\lfloor \frac{k-5}{2} \rfloor - 10} (Q_{\lfloor \frac{k-5}{2} \rfloor}(t^8) + t^4 Q_{\lfloor \frac{k-5}{2} \rfloor - 1}(t^8)).$$

Proof. The cases with $m=0$ are easy to see, as ∇_0 only consists of one element.

To determine $P_6^k(t)$ and $P_7^k(t)$, note that

$$\nabla_1(n) = \left\{ (i) \mid 0 \leq i \leq \lfloor \frac{n-1}{2} \rfloor \right\}.$$

Therefore

$$\begin{aligned} P_6^k(t) &= t^{k+10} \sum_{i=0}^{\lfloor \frac{k-4}{2} \rfloor} t^{4i} \frac{t^{4(k-3-2i)} - 1}{t^4 - 1} \\ &= t^{k+10} Q_{k-4}(t^4) \end{aligned}$$

by Lemma B.3. Similarly,

$$\begin{aligned} P_7^k(t) &= t^{u(k,1)} \sum_{i=0}^{\lfloor \frac{k-5}{2} \rfloor} t^{4i} \frac{t^{8(\lfloor \frac{k-5}{2} \rfloor + 1 - i)} - 1}{t^8 - 1} \\ &= t^{6k-8\lfloor \frac{k-5}{2} \rfloor - 10} (Q_{\lfloor \frac{k-5}{2} \rfloor}(t^8) + t^4 Q_{\lfloor \frac{k-5}{2} \rfloor - 1}(t^8)) \end{aligned}$$

by Lemma B.4. \square

10. POINCARÉ POLYNOMIALS FOR LINKAGE SPACES IN ODD DIMENSIONAL EUCLIDEAN SPACES

In order to calculate the Poincaré polynomial of $\mathcal{M}_d(\ell)$ for $d \geq 4$, we want to take the filtration

$$\emptyset = \mathcal{M}^0 \subset \mathcal{M}^1 \subset \cdots \subset \mathcal{M}^m = \mathcal{M}_d(\ell)$$

which arises from the $SO(d-1)$ -invariant Morse-Bott function from Proposition 6.1, so that

$$\mathcal{M}^{s+1} \simeq \mathcal{M}^s \cup_{\partial X_d^k} X_d^k$$

for all $s = 0, \dots, m-1$ and appropriate k depending on s . The long exact sequence of the pair $(\mathcal{M}^{s+1}, \mathcal{M}^s)$ takes on the form

$$\dots \longrightarrow H_{t+1}(X_d^k, \partial X_d^k) \longrightarrow H_t(\mathcal{M}^s) \longrightarrow H_t(\mathcal{M}^{s+1}) \longrightarrow H_t(X_d^k, \partial X_d^k) \longrightarrow \dots$$

If we look at $u(k, m)$ in Theorem 9.5, we see that for fixed m this number is always odd (for m even) or even (for m odd) for all $k \geq 2m+3$. It follows that

$$(7) \quad H_*(\mathcal{M}^{s+1}; \mathbb{Q}) = H_*(\mathcal{M}^s; \mathbb{Q}) \oplus H_*(X_d^k, \partial X_d^k; \mathbb{Q})$$

for odd $d \geq 5$.

This is not true for $d \geq 4$ even, as the following example shows.

Example 10.1. Let $\ell^6 = (1, 1, 1, 1, 1, 4)$. Then $a_0(\ell^6) = 1$ and $a_i(\ell^6) = 0$ for all $i \geq 1$. Therefore the Morse numbers $\mu_i(\ell)$ of the Morse function $f_3 : \mathcal{M}_3(\ell^6) \rightarrow \mathbb{R}$ of Corollary 5.3 are all 1. Since the function is constructed by induction over ℓ' with similar Morse functions, we can assume that the indices in the filtration are increasing. We thus have 4 critical points of index 0, 2, 4 and 6, respectively, so the respective values for k are 0, 1, 2 and 3.

If we look at the analogous function for $d = 4$, the filtration satisfies

$$\mathcal{M}^1 \simeq *, \quad \mathcal{M}^2 \simeq S^5, \quad \mathcal{M}^3 = \mathcal{M}_4(\ell^6) \simeq \mathcal{M}^2 \cup e^6 \cup e^8$$

which means that up to homotopy $\mathcal{M}_4(\ell^6)$ is obtained from S^5 by adding a 6-cell and an 8-cell. By Proposition A.1, we have $\mathcal{M}_4(\ell^6) = \Sigma_3^5$, and the \mathbb{Z} -homology of this space has been calculated in [17, Table 5.3] as

$$H_*(\Sigma_3^5) = \begin{cases} \mathbb{Z} & * = 8 \\ \mathbb{Z}/2\mathbb{Z} & * = 5 \\ 0 & \text{else} \end{cases}.$$

This shows that there is a non-trivial interaction between the critical points of index 2 and 3, which persists when looking at $\ell^n = (1, \dots, 1, n-2) \in \mathbb{R}^n$, as [17, Table 5.3] shows.

One would expect similar interactions when looking at more general ℓ , but we leave that for a future project.

Definition 10.2. Let $\ell \in \mathbb{R}^n$ be a generic length vector, and $d \geq 2$. We denote the Poincaré polynomial of $\mathcal{M}_d(\ell)$ with \mathbb{Z} coefficients by $P_d^\ell(t)$.

The next proposition follows by a simple induction on (7), using Proposition 5.4.

Proposition 10.3. *Let $\ell \in \mathbb{R}^n$ be a generic length vector, and $d = 2m + 5$ with $m \geq 0$. Then*

$$P_d^\ell(t) = a_0(\ell) + \sum_{k=2m+3}^{n-3} \mu_k(\ell) P_d^k(t),$$

where $\mu_k(\ell)$ are as in Proposition 5.4.

We can express the μ_k in terms of a_k , and the $P_d^k(t)$ are given by Theorem 9.5. Using Corollary 9.6, we can make the dependence on the a_k more explicit.

Theorem 10.4. *Let $\ell \in \mathbb{R}^n$ be a generic length vector with $a_0(\ell) = 1$. Let $m \geq 3$ be such that $n = 2m - 1$ or $n = 2m$. Then*

$$P_5^\ell(t) = 1 + t^9 \cdot \sum_{i=0}^{m-2} (a_i - a_{n-2-i}) (Q_{n-6-i}(t^4) - Q_{i-4}(t^4)),$$

where $a_i = a_i(\ell)$ for $i = 1, \dots, n-3$, and $a_{n-2} = 0 = Q_j$ for $j < 0$.

Proof. We know from Proposition 5.4 that $a_0(\ell)$ contributes to each $\mu_k(\ell)$ for $k = 0, \dots, n-3$. Similarly, $a_1(\ell) - a_{n-3}(\ell)$ contributes to $\mu_1(\ell), \dots, \mu_{n-4}(\ell)$, and $a_{m-2} - a_{n-m}$ contributes to μ_{m-2} and μ_{n-m-1} .

Notice that μ_1 and μ_2 have no impact on the homology of $\mathcal{M}_5(\ell)$.

According to Corollary 9.6 the contribution of $a_0(\ell) = 1$ to the Poincaré polynomial is therefore

$$\begin{aligned} 1 + \sum_{k=3}^{n-3} P_5^k(t) &= 1 + \sum_{k=3}^{n-3} \left(t^{4k-4 \lfloor \frac{k-3}{2} \rfloor - 3} \sum_{i=0}^{\lfloor \frac{k-3}{2} \rfloor} t^{4i} \right) \\ &= 1 + t^9 \cdot \sum_{k=0}^{n-6} t^{4 \lfloor \frac{k+1}{2} \rfloor} \frac{t^{4 \lfloor \frac{k+2}{2} \rfloor} - 1}{t^4 - 1} \\ &= 1 + t^9 Q_{n-6}(t^4), \end{aligned}$$

where we use Lemma B.3 in the last line.

Similarly, the contribution of $a_1(\ell) - a_{n-3}(\ell)$ is $t^9 Q_{n-7}(t^4)$, and so on. But notice that for $j \geq 4$ we get for the contribution of $a_j(\ell) - a_{n-2-j}(\ell)$ the formula

$$t^9 \sum_{k=j-3}^{n-6-j} t^{4 \lfloor \frac{k+1}{2} \rfloor} \frac{t^{4 \lfloor \frac{k+2}{2} \rfloor} - 1}{t^4 - 1} = t^9 (Q_{n-6-j}(t^4) - Q_{j-4}(t^4)).$$

Since we set $Q_k(t) = 0$ for negative k , this also holds for all $j \geq 0$. Adding all terms together gives the result. \square

Remark 10.5. If we write

$$R_k(t) = 1 + t + \dots + t^k,$$

and $R_k(t) = 0$ for $k < 0$, we can describe the Poincaré polynomials of $\mathcal{M}_3(\ell)$ as

$$P_3^\ell(t) = 1 + t^2 \cdot \sum_{i=0}^{m-2} (a_i - a_{n-2-i}) (R_{n-4-i}(t^2) - R_{i-2}(t^2)),$$

as follows easily from (1). Furthermore, we have $Q_{2m}(t) = R_m(t)R_m(t)$ and $Q_{2m+1}(t) = R_m(t)R_{m+1}(t)$. It is therefore natural to ask what the correct formula for $P_{2m+5}^\ell(t)$ is and whether it fits into a similar pattern. However, by looking at Corollary 9.6 in the case $d = 7$, we see that the Poincaré polynomial of $\mathcal{M}_7(\ell)$ will have non-zero coefficients in even degrees between 26 and \mathbf{d}_7^n for $n \geq 9$.

Example 10.6. There exist 135 chambers for $n = 7$ up to permutations [10], and the Poincaré polynomial for $\ell \in \mathbb{R}^7$ is

$$P_5^\ell(t) = a_0(\ell)(1 + t^9 + t^{13}) + (a_1(\ell) - a_4(\ell))t^9.$$

Also, $a_4(\ell) = 0$, unless $\ell = (1, 1, 1, 1, 5, 5, 5)$, in which case $a_1(\ell) - a_4(\ell) = 3$. If we also assume that ℓ is different from $(1, 1, 1, 1, 1, 1, 7)$, the Poincaré polynomial is

$$P_5^\ell(t) = 1 + (a_1(\ell) + 1)t^9 + t^{13}.$$

Since $a_1(\ell) \in \{0, \dots, 6\}$ there are not a lot of variations among the Poincaré polynomials. Also notice that $\mathcal{M}_5(\ell)$ up to homotopy is obtained from a wedge of $(a_1(\ell) + 1)$ 9-spheres by attaching a cell of dimension 10, 11 and 13. As these three cells correspond to $a_0(\ell)$ it seems unlikely to expect too many different homotopy types between the chambers for $n = 7$.

11. THE EULER CHARACTERISTIC FOR EVEN DIMENSIONAL LINKAGE SPACES

Let us begin with $\chi(X_{2m+4}^k, \partial X_{2m+4}^k)$. This is obtained by evaluating $P_{2m+4}^k(-1)$ in Theorem 9.5. Reorganising this term gives the following proposition.

Proposition 11.1. *Let $m \geq 0$ and $k \geq m + 1$. Then*

$$\chi(X_{2m+4}^{2k}, \partial X_{2m+4}^{2k}) = (-1)^{m+1} \sum_{j_m=0}^{k-m-1} \sum_{j_{m-1}=j_m}^{k-m-1} \cdots \sum_{j_1=j_2}^{k-m-1} (2k - 2m - 2j_1 - 1),$$

and

$$\chi(X_{2m+4}^{2k+1}, \partial X_{2m+4}^{2k+1}) = (-1)^m \sum_{j_m=0}^{k-m-1} \sum_{j_{m-1}=j_m}^{k-m-1} \cdots \sum_{j_1=j_2}^{k-m-1} (2k - 2m - 2j_1).$$

The simplest cases $m = 0$ and $m = 1$ are easily seen to give the following.

Corollary 11.2. *Let $m \geq 0$ and $k \geq m + 1$. Then*

$$\chi(X_{2m+4}^{2k}, \partial X_{2m+4}^{2k}) + \chi(X_{2m+4}^{2k+1}, \partial X_{2m+4}^{2k+1}) = (-1)^m |\nabla_m(2k - 2m - 1)|.$$

Furthermore, for $k \geq 0$ we get

$$\chi(X_4^k, \partial X_4^k) = (-1)^{k+1}(k - 1)$$

and

$$\begin{aligned} \chi(X_6^{2k}, \partial X_6^{2k}) &= (k - 1)^2 \\ \chi(X_6^{2k+1}, \partial X_6^{2k+1}) &= -k(k - 1). \end{aligned}$$

It is worth pointing out that X_d^0 is a point with empty boundary, so the Euler characteristic is just 1.

Corollary 11.3. *Let $k \geq 3$ and $\ell \in \mathbb{R}^n$ be a generic length vector, and $n = 2k$ or $n = 2k - 1$. Then for $n = 2k$ we get*

$$\chi(\mathcal{M}_4(\ell)) = \sum_{i=0}^{k-2} (-1)^i (a_i(\ell) - a_{2k-2-i}(\ell)) (k - 1 - i)$$

and for $n = 2k - 1$ we get

$$\chi(\mathcal{M}_4(\ell)) = -(k - 3) \sum_{i=0}^{k-2} (-1)^i (a_i(\ell) - a_{2k-3-i}(\ell)).$$

Proof. Assume that $n = 2k - 1$. Write $\chi_i = \chi(X_4^i, \partial X_4^i) = (-1)^{i+1}(i - 1)$. Then $a_0(\ell)$ contributes $\chi_0 + \chi_1 + \dots + \chi_{2k-4}$ to the Euler characteristic of $\mathcal{M}_4(\ell)$. Since $\chi_{2i} + \chi_{2i+1} = 1$, this gives a contribution of

$$\begin{aligned} \sum_{i=0}^{k-3} (\chi_{2i} + \chi_{2i+1}) + \chi_{2k-4} &= (k-2) - (2k-5) \\ &= -(k-3). \end{aligned}$$

Similarly, the contribution of $(a_j(\ell) - a_{2k-3-j}(\ell))$ is $(-1)^{i+1}(k-3)$, where for odd j one should note that $\chi_j + \chi_{j+1} = j - 1$. Summing the contributions with the appropriate factor gives the result.

The result for $n = 2k$ is using the same type of argument. \square

Example 11.4. Let $\ell = (1, \dots, 1) \in \mathbb{R}^{2m+1}$. A subset $J \subset \{1, \dots, 2m+1\}$ is ℓ -short if and only if has at most m elements. It follows that $a_i = \binom{2m}{i}$ for $i = 0, \dots, m-1$ and $a_i = 0$ for $i \geq m$. Hence

$$\begin{aligned} \chi(\mathcal{M}_4(\ell)) &= -(m-2) \sum_{i=0}^{m-1} (-1)^i \binom{2m}{i} \\ &= (-1)^m (m-2) \binom{2m-1}{m-1} \end{aligned}$$

where we used $(-1)^{m-1} \binom{2m-1}{m-1} = \sum_{i=0}^{m-1} (-1)^i \binom{2m}{i}$ which follows from the binomial formula. This formula has been obtained by Kamiyama in [13, Thm.A].

Corollary 11.5. *Let $k \geq 4$ and $\ell \in \mathbb{R}^n$ be a generic length vector, and $n = 2k$ or $n = 2k - 1$. Then for $n = 2k$ we get*

$$\chi(\mathcal{M}_6(\ell)) = \frac{1}{2} \sum_{i=0}^{k-2} (-1)^{i+1} c_i ((k-3 - \lfloor \frac{i}{2} \rfloor)(k-2 - \lfloor \frac{i}{2} \rfloor) - (\lfloor \frac{i-3}{2} \rfloor \cdot \lfloor \frac{i-1}{2} \rfloor))$$

and for $n = 2k - 1$ we get

$$\chi(\mathcal{M}_6(\ell)) = \frac{1}{2} \sum_{i=0}^{k-2} (-1)^i c_i ((k-3 - \lfloor \frac{i+1}{2} \rfloor)(k-2 - \lfloor \frac{i+1}{2} \rfloor) + (\lfloor \frac{i-3}{2} \rfloor \cdot \lfloor \frac{i-1}{2} \rfloor)),$$

where $c_i = a_i(\ell) - a_{n-2-i}(\ell)$.

Proof. The proof is along the same lines as the proof of Corollary 11.3. If $n = 2k - 1$, the contribution of a_{2j} is

$$\begin{aligned} \sum_{i=2j}^{2k-4-2j} \chi_i &= \sum_{i=j}^{k-3-j} (\chi_{2i} + \chi_{2i+1}) + \chi_{2k-4-2j} \\ &= - \sum_{i=j-1}^{k-4-j} i + (k-3-j)^2 \\ &= \frac{1}{2} ((k-3-j)(k-2-j) + (j-2)(j-1)), \end{aligned}$$

and a_{2j-1} contributes this term with a negative sign. The case $n = 2k$ is similar. \square

Example 11.6. Let $\tilde{\ell} = (1, \dots, 1, 7) \in \mathbb{R}^8$, so that $\mathcal{M}_d(\tilde{\ell}) \cong \Sigma_{d-1}^7$ by Proposition A.1. In particular, $a_0(\tilde{\ell}) = 1$ and $a_i(\tilde{\ell}) = 0$ for $i \geq 1$. Then

$$\chi(\mathcal{M}_4(\tilde{\ell})) = 3 \quad \text{and} \quad \chi(\mathcal{M}_6(\tilde{\ell})) = 0$$

as can be readily seen from Corollaries 11.3 and 11.5.

If we let $\ell = (1, 1, 1, 1, 1, 3, 3, 6)$, we see that $a_0(\ell) = 1$, $a_1(\ell) = 5$, $a_2(\ell) = 10$ and $a_i(\ell) = 0$ for $i \geq 3$. This implies

$$\chi(\mathcal{M}_4(\ell)) = \chi(\mathcal{M}_4(\tilde{\ell})),$$

while

$$\chi(\mathcal{M}_6(\ell)) = 5 \neq \chi(\mathcal{M}_6(\tilde{\ell})).$$

We can still show that $\mathcal{M}_4(\ell)$ does not have the same homotopy type as $\mathcal{M}_4(\tilde{\ell})$. To see this, note that in the filtration (\mathcal{M}^j) the relative complex $(X_4^4, \partial X_4^4)$ is attached 6 = $a_0(\ell) + a_1(\ell)$ times, so that before attaching the final $(X_4^5, \partial X_4^5)$, we have $H_{11}(\mathcal{M}^{m-1}; \mathbb{Q}) \cong \mathbb{Q}^6$. Since $H_{12}(X_4^5, \partial X_4^5; \mathbb{Q}) \cong \mathbb{Q}$, we get that the 11-th Betti number of $\mathcal{M}_4(\ell)$ is at least 5. As $H_{11}(\mathcal{M}_4(\tilde{\ell})) \cong \mathbb{Z}/2\mathbb{Z}$ by [17, Table 5.3], these spaces have different homology.

This last argument can be generalized to obtain some kind of Morse inequalities; we only give a few special cases.

Proposition 11.7. *Let $\ell \in \mathbb{R}^n$ be a generic length vector. For $n \geq 6$ we have*

$$b_{3(n-1)-10}(\mathcal{M}_4(\ell)) - b_{3(n-1)-9}(\mathcal{M}_4(\ell)) = a_1(\ell) - a_{n-3}(\ell) - a_0(\ell),$$

and for $n \geq 7$ we have

$$c_2(\ell) + c_1(\ell) + 2c_0(\ell) \geq b_{3(n-2)-10}(\mathcal{M}_4(\ell)) \geq c_2(\ell) - c_1(\ell),$$

where $c_i(\ell) = a_i(\ell) - a_{n-2-i}(\ell)$ for $i = 0, 1, 2$.

Also, for $n \geq 9$ we have

$$b_{5(n-1)-21}(\mathcal{M}_6(\ell)) - b_{5(n-1)-20}(\mathcal{M}_6(\ell)) = a_1(\ell) - a_{n-3}(\ell) - 2a_0(\ell),$$

and for $n \geq 10$ we have

$$c_2(\ell) + c_1(\ell) + c_0(\ell) \geq b_{5(n-2)-21}(\mathcal{M}_6(\ell)) \geq c_2(\ell) - c_1(\ell) - c_0(\ell).$$

Proof. The dimension of X_4^{n-4} is $3(n-1) - 10$, and in the filtration arising from the standard Morse-Bott function we get $c_1 + c_0$ -many of those. Therefore the $(3n-1) - 10$ -th Betti number is at most $c_1 + c_0$. Furthermore, only one X_4^{n-3} can occur in the filtration, and only at the very end, so the $(3n-1) - 9$ -th Betti number can be at most $a_0(\ell)$. When obtaining the homology of $\mathcal{M}_4(\ell)$ from the filtration, this generator in degree $(3n-1) - 9$ may or may not cancel with a generator in degree $(3n-1) - 10$. In either case, the difference of the Betti numbers is as claimed.

To determine $b_{3(n-2)-10}(\mathcal{M}_4(\ell))$, note that only X_4^{n-5} 's and the final X_4^{n-3} can contribute to this number, and each X_4^{n-4} may cancel a generator. As $(X_4^{n-4}, \partial X_4^{n-4})$ has homology in degrees $3(n-2) - 11$ and $3(n-2) - 9$, we could get two cancellations. As there are $(c_2 + c_1 + 2c_0)$ -many X_4^{n-5} and X_4^{n-3} , and $(c_1 + c_0)$ -many of X_4^{n-4} in the filtration, the result follows.

The result for $d = 6$ is analogous, but we use a different Poincaré polynomial for $(X_6^{n-4}, \partial X_6^{n-4})$, see Corollary 9.6. Note that the conditions $n \geq 9$ and $n \geq 10$ ensure that $Q_2(t^4)$ is used, which means that the coefficient of t^4 is 2. If $Q_1(t^4)$ is used, the formulas slightly improve. \square

APPENDIX A. SHAPE SPACES

In this appendix, we define the shape spaces occurring in statistical shape theory. Roughly speaking, a shape is a collection of points in some \mathbb{R}^d up to rotations and translations and scalings. More information can be found for example in the book [17].

We begin by defining the pre-shape space. Let

$$\mathbf{S}_d^n = \left\{ (x_1, \dots, x_n) \in (\mathbb{R}^d)^n \left| \sum_{i=1}^n x_i = 0, \sum_{i=1}^n |x_i|^2 = 1 \right. \right\}.$$

This is the space of n points in \mathbb{R}^d , whose centroid is 0 and who are scaled to sit on the unit sphere. Notice that \mathbf{S}_d^n is the intersection of the sphere of dimension $nd - 1$ with a sub-vector space of codimension d . It is therefore a sphere of dimension $(n - 1)d - 1$.

The group $SO(d)$ acts diagonally on \mathbf{S}_d^n , and the resulting quotient is called the *shape space*

$$\Sigma_d^n = \mathbf{S}_d^n / SO(d).$$

Similarly, one can define the *size and shape space* $S\Sigma_d^n$ which is obtained in the same way, but by dropping the condition that the points sit on the $(nd - 1)$ -sphere, see [17, §11.2].

As is pointed out in [10], If we define $T : S\Sigma_d^n \rightarrow \mathbb{R}^n$ by

$$T(x_1, \dots, x_n) = (|x_1 - x_2|, \dots, |x_{n-1} - x_n|, |x_n - x_1|)$$

we get that $\mathcal{M}_d(\ell) = T^{-1}(\{\ell\})$. Furthermore, looking at inverse images of chambers and further stratas in \mathbb{R}^d leads to a decomposition of the size and shape space by configuration spaces of linkages studied in [10].

An even more direct relation between linkage spaces and shape spaces is given by the next proposition.

Proposition A.1. *Let $\ell = (1, \dots, 1, n - 2) \in \mathbb{R}^n$. Then there exists an $SO(d - 1)$ -equivariant homeomorphism*

$$\Phi : \mathcal{C}_d(\ell) \rightarrow \mathbf{S}_{d-1}^{n-1}.$$

In particular, the shape space Σ_{d-1}^{n-1} is homeomorphic to $\mathcal{M}_d(\ell)$.

Proof. Let $(x_1, \dots, x_{n-1}) \in \mathcal{C}_d(\ell)$. If $p_1 : \mathbb{R}^d \rightarrow \mathbb{R}$ is projection to the first coordinate, notice that $p_1(x_i) < 0$ by elementary geometry. Now let $p : \mathbb{R}^d \rightarrow \mathbb{R}^{d-1}$ be projection to the last $d - 1$ coordinates. Let

$$c = \sum_{i=1}^{n-1} |p(x_i)|^2.$$

Then $c > 0$, for otherwise each $x_i = -e_1 \in \mathbb{R}^d$, and

$$x_1 + \cdots + x_{n-1} = (-n + 1, 0, \dots, 0) \neq (-n + 2, 0, \dots, 0).$$

So we can define

$$\Phi(x_1, \dots, x_{n-1}) = (p(x_1)/\sqrt{c}, \dots, p(x_{n-1})/\sqrt{c}).$$

This is injective: If $\Phi(x_1, \dots, x_{n-1}) = \Phi(y_1, \dots, y_{n-1})$, then $p(x_i) = ep(y_i)$ for some $e > 0$ and all $i = 1, \dots, n-1$. If $e > 1$, then $p_1(x_i) < p_1(y_i)$ for all $i = 1, \dots, n-1$, but then

$$\sum_{i=1}^{n-1} p_1(x_i) \neq \sum_{i=1}^{n-1} p_1(y_i),$$

contradicting that both are in $\mathcal{C}_d(\ell)$. The case $e < 1$ leads to a similar contradiction, and $e = 1$ implies $(x_1, \dots, x_{n-1}) = (y_1, \dots, y_{n-1})$.

As $\mathcal{C}_d(\ell)$ is a closed manifold of the same dimension, Φ is also surjective. Equivariance is clear from the construction, so the statement follows. \square

APPENDIX B. POLYNOMIAL RELATIONS

We want to collect a few properties of the sequence of polynomials $Q_n(t)$ given by

$$\begin{aligned} Q_{2m}(t) &= \frac{(t^{m+1} - 1)^2}{(t - 1)^2} \\ Q_{2m+1}(t) &= \frac{(t^{m+2} - 1)(t^{m+1} - 1)}{(t - 1)^2} \end{aligned}$$

for all $m \geq 0$. For convenience, we also add the equations $Q_n(t) = 0$ for $n < 0$.

The next lemma follows directly from the fact that

$$\frac{t^{m+1} - 1}{t - 1} = 1 + t + \cdots + t^m.$$

Lemma B.1. *Let $m \geq 1$, then*

$$\begin{aligned} Q_{2m-1}(t) &= Q_{2m-2}(t) + t^m + \cdots + t^{2m-1} \\ Q_{2m}(t) &= Q_{2m-1}(t) + t^m + \cdots + t^{2m} \end{aligned}$$

Using an induction on Lemma B.1, we get a nice description for the coefficients of $Q_n(t)$.

Lemma B.2. *For $m \geq 0$ we have*

$$\begin{aligned} Q_{2m}(t) &= 1 + 2t + \cdots + (m+1)t^m + mt^{m+1} + \cdots + 2t^{2m-1} + t^{2m} \\ Q_{2m+1}(t) &= 1 + 2t + \cdots + (m+1)t^m + (m+1)t^{m+1} + \cdots + 2t^{2m} + t^{2m+1}. \end{aligned}$$

The next lemmata also follow by induction using Lemma B.1.

Lemma B.3. *For $m \geq 0$ we have*

$$\begin{aligned} Q_m(t) &= \sum_{i=0}^{\lfloor \frac{m}{2} \rfloor} t^i \frac{t^{m+1-2i} - 1}{t - 1} \\ &= \sum_{i=0}^m t^{\lfloor \frac{i+1}{2} \rfloor} \frac{t^{\lfloor \frac{i+2}{2} \rfloor} - 1}{t - 1}. \end{aligned}$$

Lemma B.4. *For $m \geq 0$ we have*

$$Q_m(t^2) + tQ_{m-1}(t^2) = \sum_{i=0}^m t^i \frac{t^{2(n+1-i)} - 1}{t^2 - 1}$$

REFERENCES

- [1] M.M. Cohen, *A course in simple-homotopy theory*, Springer-Verlag, New York-Berlin, 1973.
- [2] M. Farber, *Topology of closed 1-forms*. American Mathematical Society, Providence, RI, 2004.
- [3] M. Farber, *Invitation to topological robotics*, Zurich Lectures in Advanced Mathematics, European Mathematical Society (EMS), Zürich, 2008.
- [4] M. Farber, J.-Cl. Hausmann, D. Schütz, *On the conjecture of Kevin Walker*, J. of Topology and Analysis 1 (2009), 65-86.
- [5] M. Farber, J.-Cl. Hausmann, D. Schütz, *The Walker conjecture for chains in \mathbb{R}^d* , Math. Proc. Cambridge Philos. Soc. 151 (2011), 283-292.
- [6] M. Farber, D. Schütz, *Homology of planar polygon spaces*, Geom. Dedicata 125 (2007), 75-92.
- [7] P. Foth, G. Lozano, *The geometry of polygons in \mathbb{R}^5 and quaternions*, Geom. Dedicata 105 (2004), 209-229.
- [8] J.-Cl. Hausmann, *Sur la topologie des bras articulés*, Algebraic topology Poznan 1989, 146-159, Lecture Notes in Math. 1474, Springer, Berlin, 1991.
- [9] J.-Cl. Hausmann, A. Knutson, *The cohomology ring of polygon spaces*, Ann. Inst. Fourier (Grenoble) 48 (1998), 281-321.
- [10] J.-Cl. Hausmann, E. Rodriguez, *The space of clouds in Euclidean space*, Experiment. Math. 13 (2004), 31-47.
- [11] T.F. Havel, *Some examples of the use of distances as coordinates for Euclidean geometry*, Invariant-theoretic algorithms in geometry (Minneapolis, MN, 1987). J. Symbolic Comput. 11 (1991), 579-593.
- [12] Y. Kamiyama, *Topology of equilateral polygon linkages*, Topology Appl. 68 (1996), 13-31.
- [13] Y. Kamiyama, *Euler characteristic of the moduli space of polygons in higher-dimensional Euclidean space*, Kyushu J. Math. 54 (2000), 333-369.
- [14] Y. Kamiyama, *A perfect Morse function on planar polygon spaces* J. Math. Sci. Adv. Appl. 6 (2010), 125-136.
- [15] Y. Kamiyama, M. Tezuka, T. Toma, *Homology of the configuration spaces of quasi-equilateral polygon linkages*, Trans. Amer. Math. Soc. 350 (1998), 4869-4896.
- [16] M. Kapovich, J. Millson, *The symplectic geometry of polygons in Euclidean space*, J. Differential Geom. 44 (1996), 479-513.
- [17] D. Kendall, D. Barden, T. Carne, H. Le, *Shape and shape theory*. John Wiley & Sons, Ltd., Chichester, 1999.
- [18] A.A. Klyachko, *Spatial polygons and stable configurations of points in the projective line*, Algebraic geometry and its applications, Aspects Math., E25, Vieweg, Braunschweig, 1994, 67-84.
- [19] S.P. Novikov, *Multivalued functions and functionals. An analogue of Morse theory*, Soviet Math. Dokl. 24 (1981), 222-226.
- [20] A. Pajitnov, *Circle-valued Morse theory*. Walter de Gruyter & Co., Berlin, 2006.
- [21] I.J. Schoenberg, *Linkages and distance geometry, I. Linkages*, Indag. Math. 31 (1969), 43-52.
- [22] D. Schütz, *The isomorphism problem of planar polygon spaces*, J. Topol. 3 (2010), 713-742.
- [23] K. Walker, *Configuration spaces of linkages*, Bachelor thesis, Princeton, 1985.
- [24] A. Wasserman, *Equivariant differential topology*, Topology 8 (1969), 127-150.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF DURHAM, SCIENCE LABORATORIES,
SOUTH RD, DURHAM DH1 3LE, UNITED KINGDOM

E-mail address: dirk.schuetz@durham.ac.uk