

THE GEOMETRIC THETA CORRESPONDENCE FOR HILBERT MODULAR SURFACES

JENS FUNKE* AND JOHN MILLSON**

1. INTRODUCTION

In a series of papers [11, 12, 13, 14] we have been studying the geometric theta correspondence (see below) for non-compact arithmetic quotients of symmetric spaces associated to orthogonal groups. It is our overall goal to develop a general theory of geometric theta liftings in the context of the real differential geometry/topology of non-compact locally symmetric spaces of orthogonal and unitary groups which generalizes the theory of Kudla-Millson in the compact case, see [24].

In this paper we study in detail the geometric theta lift for Hilbert modular surfaces. In particular, we will give a new proof and an extension (to all finite index subgroups of the Hilbert modular group) of the celebrated theorem of Hirzebruch and Zagier [16] that the generating function for the intersection numbers of the Hirzebruch-Zagier cycles is a classical modular form of weight 2.¹ In our approach we replace Hirzebruch's smooth complex analytic compactification \tilde{X} of the Hilbert modular surface X with the (real) Borel-Serre compactification \overline{X} . The various algebro-geometric quantities that occur in [16] are then replaced by topological quantities associated to 4-manifolds with boundary. In particular, the "boundary contribution" in [16] is replaced by sums of linking numbers of circles (the boundaries of the cycles) in the 3-manifolds of type Sol (torus bundle over a circle) which comprise the Borel-Serre boundary.

The geometric theta correspondence. We first explain the term "geometric theta correspondence". The Weil (or oscillator) representation gives us a method to construct closed differential forms on locally symmetric spaces associated to groups which belong to dual pairs. Let V be a rational quadratic space of signature (p, q) with for simplicity even dimension. Then the Weil representation induces an action of $\mathrm{SL}_2(\mathbb{R}) \times \mathrm{O}(V_{\mathbb{R}})$ on $\mathcal{S}(V_{\mathbb{R}})$, the Schwartz functions on $V_{\mathbb{R}}$. Let $G = \mathrm{SO}_0(V_{\mathbb{R}})$ and let K be a maximal compact subgroup. We let \mathfrak{g} and \mathfrak{k} be their respective Lie algebras and let $\mathfrak{g} = \mathfrak{p} \oplus \mathfrak{k}$ be the associated Cartan decomposition. Suppose

$$\varphi \in (\mathcal{S}(V_{\mathbb{R}}) \otimes \wedge^r \mathfrak{p}^*)^K$$

is a cocycle in the relative Lie algebra complex for G with values in $\mathcal{S}(V)$. Then φ corresponds to a closed differential r -form $\tilde{\varphi}$ on the symmetric space $D = G/K$ of

Date: May 24, 2011.

* Partially supported by NSF grant DMS-0710228.

** Partially supported by NSF grant DMS-0907446, NSF FRG grant DMS-0554254, and the Simons Foundation.

¹Eichler, [16] p.104, proposed a proof using "Siegel's work on indefinite theta functions". This is what our proof is, though with perhaps more differential geometry than Eichler had in mind.

dimension pq with values in $\mathcal{S}(V)$. For a coset of a lattice \mathcal{L} in V , we define the theta distribution $\Theta = \Theta_{\mathcal{L}}$ by $\Theta = \sum_{\ell \in \mathcal{L}} \delta_{\ell}$, where δ_{ℓ} is the delta measure concentrated at ℓ . It is obvious that Θ is invariant under $\Gamma = \text{Stab}(\mathcal{L}) \subset G$. There is also a congruence subgroup Γ' of $\text{SL}(2, \mathbb{Z})$ such that Θ is also invariant under Γ' . Hence we can apply the theta distribution to $\tilde{\varphi}$ to obtain a closed r -form θ_{φ} on $X = \Gamma \backslash D$ given by

$$\theta_{\varphi}(\mathcal{L}) = \langle \Theta_{\mathcal{L}}, \tilde{\varphi} \rangle.$$

Assume now in addition that φ has weight k under the maximal compact subgroup $\text{SO}(2) \subset \text{SL}_2(\mathbb{R})$. Then θ_{φ} also gives rise to a (in general) non-holomorphic function on the upper half plane \mathbb{H} which is modular of weight k for Γ' . We may then use θ_{φ} as the kernel of a pairing of modular forms f with (closed) differential $(pq - r)$ -forms η or r -chains (cycles) C in X . The resulting pairing in f , η (or C), and φ as these objects vary, we call the **geometric theta correspondence**.

The cocycle of Kudla-Millson. The key point of the work of Kudla and Millson [21, 22] is that they found (in greater generality) a family of cocycles φ_q^V in $(\mathcal{S}(V) \otimes \wedge^q \mathfrak{p}^*)^K$ with weight $(p+q)/2$ for SL_2 . Moreover, these cocycles give rise to Poincaré dual forms for certain totally geodesic, “special” cycles in X . Recently, it has now been shown, first [17] for $\text{SO}(3, 2)$, and then [1] for all $\text{SO}(p, q)$ and $p + q > 6$ (with $p \geq q$) in the cocompact (standard arithmetic) case that the geometric theta correspondence specialized to φ_q^V induces on the adelic level an *isomorphism* from the appropriate space of classical modular forms to $H^q(X)$. In particular, for any congruence quotient, the dual homology groups are spanned by special cycles. This gives further justification to the term geometric theta correspondence and highlights the significance of these cocycles. In [12] we generalize φ_q^V to allow suitable non-trivial coefficient systems (and one has an analogous isomorphism in [1]).

The main results. In the present paper, we consider the case when V has signature $(2, 2)$ with \mathbb{Q} -rank 1. Then $D \simeq \mathbb{H} \times \mathbb{H}$, and X is a Hilbert modular surface. We let \overline{X} be the Borel-Serre compactification of X which is obtained by replacing each isolated cusp associated to a rational parabolic P with a boundary face $e'(P)$ which turns out to be a torus bundle over a circle, a 3-manifold of type Sol. This makes \overline{X} a 4-manifold with boundary. For simplicity, we assume that X has only one cusp so that $\partial \overline{X} = e'(P)$, and we write $k : \partial \overline{X} \hookrightarrow \overline{X}$ for the inclusion. The special cycles C_n^2 in question are now embedded modular and Shimura curves, and are parameterized by $n \in \mathbb{N}$. They define relative homology classes in $H_2(X, \partial X, \mathbb{Q})$.

The geometric theta correspondence of Kudla-Millson [24] for the cocycle φ_2^V in this situation takes the following shape. For a *compact* cycle C in X , we have that

$$(1.1) \quad \langle \theta_{\varphi_2^V}, C \rangle = \int_C \theta_{\varphi_2^V} = \sum_{n \geq 0} (C_n \cdot C) q^n$$

is a holomorphic modular form of weight 2 and is equal to the generating series of the intersection numbers with C_n . Here $q = e^{2\pi i \tau}$ with $\tau \in \mathbb{H}$. (There is a similar statement for the pairing of $\theta_{\varphi_2^V}$ with a closed *compactly supported* differential 2-form on X representing a class in $H_c^2(X)$, see Theorem 7.1). Our first result is

²We distinguish the relative cycles C_n in X from the Hirzebruch-Zagier cycles T_n in \tilde{X} , see below.

Theorem 1.1. *(Theorem 7.3) The differential form $\theta_{\varphi_2^V}$ on X extends to a form on \overline{X} , and the restriction k^* of $\theta_{\varphi_2^V}$ to $\partial\overline{X}$ gives an exact differential form on $\partial\overline{X}$. Moreover, there exists a theta series $\theta_{\phi_1^W}$ for a space W of signature $(1, 1)$ of weight 2 with values in the 1-forms on $\partial\overline{X}$ such that $\theta_{\phi_1^W}$ is a primitive for $k^*\theta_{\varphi_2^V}$:*

$$d(\theta_{\phi_1^W}) = k^*\theta_{\varphi_2^V}.$$

Considering the mapping cone for the inclusion $k : \partial\overline{X} \hookrightarrow \overline{X}$ (see Section 3.3) we then view the pair $[\theta_{\varphi_2^V}, \theta_{\phi_1^W}]$ as an element of the compactly supported cohomology $H_c^2(X)$. Explicitly, let C be a relative cycle in \overline{X} representing a class in $H_2(X, \partial X, \mathbb{Z})$. Then the Kronecker pairing between $[\theta_{\varphi_2^V}, \theta_{\phi_1^W}]$ and C is given by

$$(1.2) \quad \langle [\theta_{\varphi_2^V}, \theta_{\phi_1^W}], C \rangle = \int_C \theta_{\varphi_2^V} - \int_{\partial C} \theta_{\phi_1^W}.$$

In this way, we obtain an extension of the geometric theta lift which captures the non-compact situation.

To describe the geometric interpretation of this extension, we study the cycle C_n at the boundary $\partial\overline{X}$ (Section 4). The intersection of C_n with $\partial\overline{X}$ is a union of circles contained in the torus fibers of Sol. But rationally such circles are homologically trivial. Hence we can find a (suitably normalized) rational 2-chain A_n in $\partial\overline{X}$ whose boundary is the boundary of C_n in $\partial\overline{X}$. ‘‘Capping’’ off C_n by A_n , we obtain a *closed* cycle C_n^c in \overline{X} defining a class in $H_2(X, \mathbb{Q})$. Our main result is the extension of (1.1):

Theorem 1.2. *(Theorem 7.7) Let C be a relative cycle in \overline{X} . Then*

$$\langle [\theta_{\varphi_2^V}, \theta_{\phi_1^W}], C \rangle = \sum_{n \geq 0} (C_n^c \cdot C) q^n$$

is a holomorphic modular form of weight 2 and is equal to the generating series of the intersection numbers with the capped cycles C_n^c . (Similarly for the pairing with an arbitrary closed 2-form on \overline{X} representing a class in $H^2(X)$).

Note that in view of (1.2) the lift of classes of $H_2(X, \partial X)$ or $H^2(X)$ is the sum of two in general non-holomorphic modular forms (see below).

In [13] we systematically study for $O(p, q)$ the restriction of the classes $\theta_{\varphi_q^V}$ (also with non-trivial coefficients) to the Borel-Serre boundary. Whenever the restriction vanishes cohomologically, we can expect that a similar analysis to the one given in this paper will give analogous extensions of the geometric theta correspondence. In fact, aside from this paper we have at present managed to do this for several other cases, namely for modular curves with non-trivial coefficients [14] generalizing work of Shintani [27] and for Picard modular surfaces [15] generalizing work of Cogdell [6].

Linking numbers in 3-manifolds of type Sol. The theta series $\theta_{\phi_1^W}$ at the boundary is of independent interest and has geometric meaning in its own right. Recall that for two disjoint (rationally) homological trivial 1-cycles a and b in a 3-manifold M we can define the *linking number* of a and b as the intersection number

$$\text{Lk}(a, b) = A \cdot b$$

of (rational) chains in M . Here A is a 2-chain in M with boundary a . We show

Theorem 1.3. (Theorem 6.3) *Let c be homologically trivial 1-cycle in $\partial\bar{X}$ which is disjoint from the torus fibers containing components of ∂C_n . Then the holomorphic part of the weight 2 non-holomorphic modular form $\int_c \theta_{\phi_1^W}$ is given by the generating series of the linking numbers $\sum_{n>0} \text{Lk}(\partial C_n, c)q^n$.*

We also give a simple formula in Theorem 4.10 for the linking number of two circles contained in the fiber of a 3-manifold M of type Sol in terms of the glueing homeomorphism for the bundle.

One can reformulate the previous theorem stating that $\sum_{n>0} \text{Lk}(\partial C_n, c)q^n$ is a “mixed Mock modular form” of weight 2; it is the product of a Mock modular form of weight 3/2 with a unary theta series. Such forms, which originate with the famous Ramanujan Mock theta functions, have recently generated great interest.

Theorem 1.3 (and its analogues for the Borel-Serre boundary of modular curves with non-trivial coefficients and Picard modular surfaces) suggest that there is a more general connection between modular forms and linking numbers of nilmanifold subbundles over special cycles in nilmanifold bundles over locally symmetric spaces.

Relation to the work of Hirzebruch and Zagier. In their seminal paper [16], Hirzebruch-Zagier provided a map from the second homology of the smooth compactification of certain Hilbert modular surfaces $j : X \hookrightarrow \tilde{X}$ to modular forms. They introduced the Hirzebruch-Zagier curves T_n in X , which are given by the closure of the cycles C_n in \tilde{X} . They then defined “truncated” cycles T_n^c as the projections of T_n orthogonal to the subspace of $H_2(\tilde{X}, \mathbb{Q})$ spanned by the compactifying divisors of \tilde{X} . The principal result of [16] was that $\sum_{n \geq 0} [T_n^c]q^n$ defines a holomorphic modular form of weight 2 with values in $H_2(\tilde{X}, \mathbb{Q})$. We show $j_*C_n^c = T_n^c$ (Proposition 4.7), and hence the Hirzebruch-Zagier theorem follows easily from Theorem 1.2 above, see Theorem 7.9.

The main work in [16] was to show that the generating function

$$F(\tau) = \sum_{n=0}^{\infty} (T_n^c \cdot T_m)q^n$$

for the intersection numbers in \tilde{X} of T_n^c with a fixed T_m is a modular form of weight 2. The Hirzebruch-Zagier proof of the modularity of F was a remarkable synthesis of algebraic geometry, combinatorics, and modular forms. They explicitly computed the intersection numbers $T_n^c \cdot T_m$ as the sum of two terms, $T_n^c \cdot T_m = (T_n \cdot T_m)_X + (T_n \cdot T_m)_\infty$, where $(T_n \cdot T_m)_X$ is the geometric intersection number of T_n and T_m in the interior of X and $(T_n \cdot T_m)_\infty$ which they called the “contribution from infinity”. They then proved both generating functions $\sum_{n=0}^{\infty} (T_n \cdot T_m)_X q^n$ and $\sum_{n=0}^{\infty} (T_n \cdot T_m)_\infty q^n$ are the holomorphic parts of two non-holomorphic forms F_X and F_∞ with the *same* non-holomorphic part (with opposite signs). Hence combining these two forms gives $F(\tau)$.

We recover this feature of the original Hirzebruch-Zagier proof via (1.2) with $C = C_m$. The first term on the right hand side of (1.2) was studied in the thesis of the first author of this paper [9] and gives the interior intersections $(T_n \cdot T_m)_X$ encoded in F_X . So via Theorem 1.2 the second term on the right hand side of (1.2) must match the boundary contribution F_∞ in [16], that is, we obtain

Theorem 1.4.

$$(T_n \cdot T_m)_\infty = \text{Lk}(\partial C_n, \partial C_m).$$

Hence we give an interpretation for the boundary contribution in [16] in terms of linking numbers in $\partial\bar{X}$. In fact, the construction of $\theta_{\phi_1^W}$ owes a great deal to Section 2.3 in [16], where a scalar-valued version of $\theta_{\phi_1^W}$ is introduced, see also Example 6.4. Using Theorem 4.11 one can also make the connection between our linking numbers and the formulas of the boundary contribution in [16] explicit.

To summarize, we start with the difference of theta integrals (1.2) (which we know a priori is a holomorphic modular form), then by functorial differential topological computations we relate its Fourier coefficients to intersection/linking numbers, and by direct computation of the integrals involved we obtain the explicit formulas of Hirzebruch-Zagier and a “closed form” for their generating function.

Note that Bruinier [4] and Oda [26] use related theta series to consider [16], but their overall approach is different.

Currents. One of the key properties of the cocycle φ_2^V is that the n -th Fourier coefficients of $\theta_{\varphi_2^V}$ represents the Poincaré dual class for the cycle C_n . Kudla-Millson establish this by showing that φ_2^V gives rise to a Thom form for the normal bundle of each of the components of C_n . To prove our main result, Theorem 1.2, we follow a different approach using currents which is implicit in [5] and is closely related to the Green’s function $\Xi(n)$ for the divisors C_n constructed by Kudla [18, 19]. This function plays an important role in the Kudla program (see eg [20]) which considers the analogous generating series for the special cycles in arithmetic geometry. In the non-compact situation however, one needs to modify $\Xi(n)$ to obtain a Green’s function for the cycle T_n^c in \tilde{X} . Discussions with U. Kühn suggest that the constructions in this paper indeed give rise to such a modification of $\Xi(n)$, see Remark 8.5.

We would like to thank Rolf Berndt, Jan Bruinier, Jose Burgos, Misha Kapovich, and Ulf Kühn for fruitful and extensive discussions on the constructions and results of this paper. As always it is a great pleasure to thank Steve Kudla for his interest and encouragement. Each of us began the work of relating theta lifts and special cycles with him.

We dedicate this paper to the memory of Gretchen Taylor Millson, beloved wife of the second author.

2. THE HILBERT MODULAR SURFACE AND ITS BOREL-SERRE COMPACTIFICATION

2.1. The symmetric space and its arithmetic quotient.

2.1.1. *The orthogonal group and its symmetric space.* Let V be a rational vector space of dimension 4 with a non-degenerate symmetric bilinear form $(,)$ of signature $(2, 2)$. We let $\underline{G} = \text{SO}(V)$, viewed as an algebraic group over \mathbb{Q} . We let $G = \underline{G}_0(\mathbb{R}) \simeq \text{SO}_0(2, 2)$ be the connected component of the identity of the real points of \underline{G} . It is most convenient to identify the associated symmetric space $D = D_V$ with the space of negative 2-planes in $V(\mathbb{R})$ on which the bilinear form $(,)$ is negative definite:

$$D = \{z \subset V_{\mathbb{R}}; \dim z = 2 \text{ and } (,)|_z < 0\}.$$

We pick an orthogonal basis $\{e_1, e_2, e_3, e_4\}$ of $V_{\mathbb{R}}$ with $(e_1, e_1) = (e_2, e_2) = 1$ and $(e_3, e_3) = (e_4, e_4) = -1$. We denote the coordinates of a vector x with respect to this basis by x_i . We pick as base point of D the plane $z_0 = [e_3, e_4]$ spanned by e_3 and e_4 , and we let $K \simeq \mathrm{SO}(2) \times \mathrm{SO}(2)$ be the maximal compact subgroup of G stabilizing z_0 . Thus $D \simeq G/K$. Of course, $D \simeq \mathbb{H} \times \mathbb{H}$, the product of two upper half planes.

We let \underline{P} be a rational parabolic subgroup stabilizing a rational isotropic line ℓ and define $P = \underline{P}_0(\mathbb{R})$ as before. We let \underline{N} be its unipotent subgroup and $N = \underline{N}(\mathbb{R})$. We let $u = (e_1 + e_4)/\sqrt{2}$ and $u' = (e_1 - e_4)/\sqrt{2}$ be two isotropic vectors so that $(u, u') = 1$. We assume that u, u' are defined over \mathbb{Q} and obtain a rational Witt decomposition

$$V = \ell \oplus W \oplus \ell'$$

with $\ell = \mathbb{Q}u$, $\ell' = \mathbb{Q}u'$, and a subspace $W = \ell^\perp \cap \ell'^\perp$ such that $W_{\mathbb{R}} = \mathrm{span}_{\mathbb{R}}(e_2, e_3)$. The choice of u' gives a Levi splitting of \underline{P} , and we write

$$P = NAM$$

for the Langlands decomposition. Here, with respect to the basis u, e_2, e_3, u' , we have

$$\begin{aligned} N &= \left\{ n(w) = \begin{pmatrix} 1 & (\cdot, w) & -(w, w)/2 \\ & 1_w & -w \\ & & 1 \end{pmatrix}; w \in W_{\mathbb{R}} \right\}, \\ A &= \left\{ a(t) = \begin{pmatrix} t & & \\ & 1_w & \\ & & t^{-1} \end{pmatrix}; t \in \mathbb{R}_+ \right\}, \\ M &= \left\{ m(s) = \begin{pmatrix} 1 & & \\ & \cosh(s) & \sinh(s) \\ & \sinh(s) & \cosh(s) \\ & & & 1 \end{pmatrix}; s \in \mathbb{R} \right\}. \end{aligned}$$

Note $N \simeq W_{\mathbb{R}}$. We obtain coordinates for D by $z = z(t, s, w)$ where z is the negative two-plane in $V_{\mathbb{R}}$ with $z = [n(w)a(t)m(s)e_3, n(w)a(t)m(s)e_4]$.

2.1.2. Arithmetic Quotient. We let L be an even lattice in V of level N , that is $L \subseteq L^\#$, the dual lattice, $(x, x) \in 2\mathbb{Z}$ for $x \in L$, and $q(L^\#)\mathbb{Z} = \frac{1}{N}\mathbb{Z}$. We fix $h \in L^\#$ and let $\Gamma \subseteq \mathrm{Stab} L$ be a subgroup of finite index of the stabilizer of $\mathcal{L} := L + h$ in G . For each isotropic line $\ell = \mathbb{Q}u$, we assume that u is primitive in the lattice L in V . We will throughout assume that the \mathbb{Q} -rank of \underline{G} is 1, that is, V splits exactly one hyperbolic plane over \mathbb{Q} . Then we define the Hilbert modular surface

$$X = \Gamma \backslash D.$$

Example 2.1. An important example is the following. Let $d > 0$ be the discriminant of the real quadratic field $K = \mathbb{Q}(\sqrt{d})$ over \mathbb{Q} , \mathcal{O}_K its ring of integers. We denote by $x \mapsto x'$ the Galois involution on K . We let $V \subset M_2(K)$ be the space of skew-hermitian matrices in $M_2(K)$, i.e., which satisfy ${}^t x' = -x$. Then the determinant on $M_2(K)$ gives V the structure of a non-degenerate rational quadratic space of signature $(2, 2)$ and \mathbb{Q} -rank 1. We define the integral skew-hermitian matrices by

$$L = \left\{ x = \begin{pmatrix} a\sqrt{d} & \lambda \\ -\lambda' & b\sqrt{d} \end{pmatrix} : a, b \in \mathbb{Z}, \lambda \in \mathcal{O}_K \right\}.$$

Then L is a lattice of level d . We embed $\mathrm{SL}_2(K)$ into $\mathrm{SL}_2 \mathbb{R} \times \mathrm{SL}_2(\mathbb{R})$ by $g \mapsto (g, g')$ so that $\mathrm{SL}_2(\mathcal{O}_K)$ acts on L by $\gamma.x = \gamma x {}^t \gamma'$ as isometries. Hirzebruch and Zagier actually considered this case for $d \equiv 1 \pmod{4}$ a prime.

The quotient space X is in general an oriented uniformizable orbifold with isolated singularities. We will treat X as a manifold - we will use Stokes' Theorem and Poincaré duality over \mathbb{Q} on X . This is justified because in each instance we can pass to a finite normal cover Y of X with Y a manifold. Hence, the formulas we want hold on Y . We then then go back to the quotient by taking invariants or summing over the group Φ of covering transformations. The point is that the de Rham complex of X is the algebra of Φ -invariants in the one of Y and the *rational* homology (cohomology) groups of X are the groups of Φ -coinvariants (invariants) of those of Y .

2.2. Compactifications.

2.2.1. *Admissible Levi decompositions of P .* We let $\Gamma_P = \Gamma \cap P$ and $\Gamma_N = \Gamma_P \cap N$. Then the quotient Γ_P/Γ_N is a non-trivial arithmetic subgroup of $\underline{P}/\underline{N}$ and lies inside the connected component of the identity of the real points of $\underline{P}/\underline{N}$. Furthermore, Γ_P/Γ_N acts as isometries of spinor norm 1 on the anisotropic quadratic space ℓ^\perp/ℓ of signature $(1, 1)$. Hence $\Gamma_P/\Gamma_N \simeq \mathbb{Z}$ is infinite cyclic. Therefore the exact sequence

$$1 \rightarrow \Gamma_N \rightarrow \Gamma_P \rightarrow \Gamma_P/\Gamma_N \rightarrow 1$$

splits. We fix $g \in \Gamma_P$ such that its image \bar{g} generates Γ_P/Γ_N . Then g defines a Levi subgroup M . In fact, the element g generates $\Gamma_M := \Gamma_P \cap M$. Hence

$$\Gamma_P = \Gamma_M \rtimes \Gamma_N.$$

We will say a Levi decomposition $P = NAM$ is admissible if $\Gamma_P = (M \cap \Gamma_P) \rtimes \Gamma_N$. In the following we assume that we have picked an admissible Levi decomposition for each rational parabolic.

2.2.2. *Borel-Serre compactification.* We let \bar{D} be the (rational) Borel-Serre enlargement of D , see [3] or [2], III.9. For any parabolic \underline{P} as before with admissible Levi decomposition $P = NAM$, we define the boundary component

$$e(P) = MN \simeq D_W \times W.$$

Here $D_W \simeq M \simeq \mathbb{R}$ is the symmetric space associated to the orthogonal group of W . Then \bar{D} is given by

$$\bar{D} = D \cup \coprod_{\underline{P}} e(P),$$

where \underline{P} varies over all rational parabolics. The action of Γ on D extends to \bar{D} in a natural way, and we let

$$\bar{X} := \Gamma \backslash \bar{D}$$

be the Borel-Serre compactification of $X = \Gamma \backslash D$. This makes \bar{X} a manifold with boundary such that

$$\partial \bar{X} = \coprod_{[\underline{P}]} e'(P),$$

where for each cusp, the corresponding boundary component is given by

$$e'(P) = \Gamma_P \backslash e(P).$$

Here $[\underline{P}]$ runs over all Γ -conjugacy classes. The space $X_W := \Gamma_M \backslash D_W$ is a circle. Hence $e'(P)$ is a torus bundle over the circle, where the torus T^2 is given by $\Gamma_N \backslash N$.

That is, $e'(P) = X_W \times T^2$, and we have the natural map $\kappa : e'(P) \rightarrow X_W$. We have a natural product neighborhood of $e(P)$ in \overline{D} and hence for $e'(P)$ in \overline{X} given by $[(T, \infty] \times e'(P)]$ for T sufficiently large given by $z(t, s, w)$ with $t > T$. We let $i : X \hookrightarrow \overline{X}$ and $i_P : e'(P) \hookrightarrow \overline{X}$ be the natural inclusions.

It is one of the fundamental properties of the Borel-Serre compactification \overline{X} that it is homotopic equivalent to X itself. Hence their (co)homology groups coincide.

2.2.3. Hirzebruch's smooth compactification. We let X' be the Baily-Borel compactification of X , which is obtained by collapsing in \overline{X} each boundary component $e'(P)$ to a single point or topologically by taking a cone on each component of the Borel-Serre boundary. It is well known that X' is a projective algebraic variety. We let \tilde{X} be Hirzebruch's smooth resolution of the cusp singularities and $\pi : \tilde{X} \rightarrow X'$ be the natural map collapsing the compactifying divisors for each cusp. We let $j : X \hookrightarrow \tilde{X}$ be the natural embedding. Note that the Borel-Serre boundary separates \tilde{X} into two pieces, the (connected) inside X^{in} , which is isomorphic to X and the (disconnected) outside X^{out} , which for each cusp is a neighborhood of the compactifying divisors. Note that we can view $e'(P)$ as lying in both X^{in} and X^{out} since the intersection $X^{in} \cap X^{out}$ is equal to $\coprod_P e(P)$.

3. (CO)HOMOLOGY

In this section we describe the relationship between the (co)homology of the various compactifications.

3.1. The homology of the boundary components. Every element of $\Gamma_N = \pi_1(T^2)$ is a rational multiple of a commutator in Γ_P and accordingly the image of $H_1(T^2, \mathbb{Q})$ in $H_1(e'(P), \mathbb{Q})$ is trivial. Let $a_P \in H_1(e'(P), \mathbb{Z})$ be the class of the identity section of $\kappa : e'(P) \rightarrow X_W$ and $b_P \in H_2(e'(P), \mathbb{Z})$ be the class of the torus fiber of κ . It is clear that the intersection number of a_P and b_P is 1 (up to sign) whence a_P and b_P are nontrivial primitive classes. Furthermore, a_P generates $H_1(e'(P), \mathbb{Q})$ and $H_2(e'(P), \mathbb{Z}) \cong \mathbb{Z}$, generated by b_P . So

Lemma 3.1. (i) *The first rational homology group of $e'(P)$ is generated by a_P .*
(ii) *The second homology group of $e'(P)$ is generated by b_P .*

Remark 3.2. To compute the homology over \mathbb{Z} one has only to use the Wang sequence for a fiber bundle over a circle, see [25], page 67.

Let Ω_P be the unique P -invariant 2-form on $e'(P)$ such that

$$(3.1) \quad \int_{b_P} \Omega_P = 1.$$

Since b_P is the image of the fundamental class of T^2 inside $H_2(e'(P), \mathbb{Z})$, we see that that the restriction of Ω_P to T^2 lifts to the area form on $W_{\mathbb{R}} \simeq N$ normalized such that $T^2 = \Gamma_N \backslash N$ has area 1.

3.2. Homology and cohomology of X and \tilde{X} . Accordingly to the discussion in Section 2.2.3 we have the Mayer-Vietoris sequence

$$0 \rightarrow \oplus_P H_2(e'(P)) \rightarrow H_2(X) \oplus (\oplus_P S_P) \rightarrow H_2(\tilde{X}) \rightarrow 0.$$

Here S_P denotes the span of the classes defined by compactifying divisors at the cusp associated to P . The zero on the left comes from $H_3(\tilde{X}) = 0$ and the zero on the right comes from the fact that for each P the class a_P injects into $H_1(X^{out})$, see [28], II.3. Since the generator b_P has trivial intersection with each of the compactifying divisors, b_P bounds on the outside so a fortiori it bounds in \tilde{X} . Thus the above short exact sequence is the sum of the two short exact sequences $\oplus_P H_2(e'(P)) \rightarrow H_2(X) \rightarrow j_* H_2(X)$ and $0 \rightarrow \oplus_P S_P \rightarrow \oplus_P S_P$. By adding the third terms of the two sequences and equating them to $H_2(\tilde{X})$ we obtain the orthogonal splittings (for the intersection pairing) - see also [28], p.123,

$$H_2(\tilde{X}) = j_* H_2(X) \oplus (\oplus_{[P]} S_P), \quad H^2(\tilde{X}) = j_{\#} H_c^2(X) \oplus (\oplus_{[P]} S_P^{\vee}).$$

Here $j_{\#}$ is the push-forward map. Furthermore, the pairings on each summand are non-degenerate. Considering $\oplus_P H_2(e'(P)) \rightarrow H_2(X) \rightarrow j_* H_2(X)$ we also obtain

Proposition 3.3. $H_2(\partial\bar{X})$ is the kernel of j_* so that

$$j_* H_2(X) \simeq H_2(X) / \sum_{[P]} H_2(e'(P)).$$

3.3. Compactly supported cohomology and the cohomology of the mapping cone. We briefly review the mapping-cone-complex realization of the cohomology of compact supports of X . For a more detailed discussion, see [14], section 5.

We let $A_c^{\bullet}(X)$ be the complex of compactly supported differential forms on X which gives rise to $H_c^{\bullet}(X)$, the cohomology of compact supports. We now represent the compactly-supported cohomology of X by the cohomology of the mapping cone C^{\bullet} of i^* , see [29], p.19, where as before $i : X \hookrightarrow \bar{X}$. However, we will change the sign of the differential on C^{\bullet} and shift the grading down by one. Thus we have

$$C^i = \{(a, b), a \in A^i(\bar{X}), b \in A^{i-1}(\partial\bar{X})\}$$

with $d(a, b) = (da, i^*a - db)$. If (a, b) is a cocycle in C^{\bullet} we will use $[[a, b]]$ to denote its cohomology class. We have

Proposition 3.4. *The cochain map $A_c^{\bullet}(X) \rightarrow C^{\bullet}$ given by $c \mapsto (c, 0)$ is a quasi-isomorphism.*

We now give a cochain map from C^{\bullet} to $A_c^{\bullet}(X)$ which induces the inverse to the above isomorphism. We let V be a product neighborhood of $\partial\bar{X}$ as in Section 2.2.2, and we let $\pi : V \rightarrow \partial\bar{X}$ be the projection. If b is a form on $\partial\bar{X}$ we obtain a form π^*b on V . Let f be a smooth function of the geodesic flow coordinate t which is 1 near $t = \infty$ and zero for $t \leq T$ for some sufficiently large T . We may regard f as a function on V by making it constant on the $\partial\bar{X}$ factor. We extend f to all of \bar{X} by making it zero off of V . Let (a, b) be a cocycle in C^i . Then there exist a compactly supported closed form α and a form μ which vanishes on $\partial\bar{X}$ such that

$$a - d(f\pi^*b) = \alpha + d\mu.$$

We define the cohomology class $[a, b]$ in the compactly supported cohomology $H_c^i(X)$ to be the class of α , and the assignment $[[a, b]] \mapsto [a, b]$ gives the desired inverse. From this we obtain the following integral formulas for the Kronecker pairings with $[a, b]$.

Lemma 3.5. *Let η be a closed form on \bar{X} and C a relative cycle in \bar{X} of appropriate degree. Then*

$$\langle [a, b], [\eta] \rangle = \int_{\bar{X}} a \wedge \eta - \int_{\partial \bar{X}} b \wedge i^* \eta, \quad \text{and} \quad \langle [a, b], C \rangle = \int_C a - \int_{\partial C} b.$$

4. CAPPED SPECIAL CYCLES AND LINKING NUMBERS IN SOL

For $x \in V$ such that $(x, x) > 0$, we define

$$D_x = \{z \in D; z \perp x\}.$$

Then D_x is an embedded upper half plane in D . We let $\Gamma_x \subset \Gamma$ be the stabilizer of x and define the special or Hirzebruch-Zagier cycle by

$$C_x = \Gamma_x \backslash D_x,$$

and by slight abuse identify C_x with its image in X . These are modular or Shimura curves. For positive $n \in \mathbb{Q}$, we write $\mathcal{L}_n = \{x \in \mathcal{L}; \frac{1}{2}(x, x) = n\}$. Then the composite cycles C_n are given by

$$C_n = \sum_{x \in \Gamma \backslash \mathcal{L}_n} C_x.$$

Since the divisors define in general relative cycles, we take the sum in $H_2(X, \partial X, \mathbb{Z})$.

4.1. The closure of special cycles in the Borel-Serre boundary and the capped cycle C_x^c . We now study the closure of C_x in $\partial \bar{X}$, which is the same as the intersection of \bar{C}_x or ∂C_x with the union of the hypersurfaces $e'(P)$. A straightforward calculation gives

Proposition 4.1. *If $(x, u) \neq 0$ then there exists a neighborhood U_∞ of $e(P)$ such that*

$$D_x \cap U_\infty = \emptyset.$$

If $(x, u) = 0$, then $\bar{D}_x \cap e(P)$ is contained in the fiber of p over $s(x)$, where $s(x)$ is the unique element of \mathbb{R} satisfying

$$(x, m(s(x))e_3) = 0.$$

At $s(x)$ the intersection $\bar{D}_x \cap e(P)$ is the affine line in W given by

$$\{w \in W : (x, w) = (u', x)\}.$$

We define $c_x \subset \partial C_x$ to be the closed geodesic in the fiber over $s(x)$ which is the image of $\bar{D}_x \cap e(P)$ under the covering $e(P) \rightarrow e'(P)$. We have

- Proposition 4.2.**
- (i) *The 1-cycle ∂C_x is a finite union of circles.*
 - (ii) *At a cusp associated to P , each circle is contained in a fiber of the map $\kappa : e'(P) \rightarrow X_W$ and hence is a rational boundary (by Lemma 3.1).*
 - (iii) *Two boundary circles c_x and c_y are parallel if they are contained in the same fiber. In particular, $c_x \cap c_y \neq \emptyset \iff c_x = c_y$.*

We now describe the intersection of \overline{C}_n or ∂C_n with $e'(P)$. For $\mathcal{L}_V = \mathcal{L} = L + h$ we can write

$$\mathcal{L}_W = \mathcal{L}_{W_P} = (\mathcal{L} \cap u^\perp) / (\mathcal{L} \cap u) \simeq \coprod_k (L_{W,k} + h_{W,k})$$

for some lattices $L_{W,k} \subset W$ and vectors $h_{W,k} \in L_{W,k}^\#$.

Via the isomorphism $W \simeq N$, we can identify $\Gamma_N = N \cap \Gamma$ with a lattice Λ_W in W . Since u is primitive in L and $n(w)x = x + (w, x)u$ for a vector $x \in u^\perp$ we see that \mathcal{L}_W is contained in the dual lattice of Λ_W .

Lemma 4.3. *The intersection $\partial C_n \cap e'(P)$ is given by*

$$\coprod_{\substack{x \in \Gamma_M \backslash \mathcal{L}_W \\ (x,x)=2n}} \coprod_{0 \leq k < \min'_{\lambda \in \Lambda_W} |(\lambda, x)|} c_{x+ku}.$$

Here \min' denotes that we take the minimum over all nonzero values of $|(\lambda, x)|$.

Proof. We will first prove $\partial C_{n,P} := \partial C_n \cap e'(P)$ is a disjoint union

$$(4.1) \quad \partial C_{n,P} = \coprod_{y \in \Gamma_P \backslash \mathcal{L}_{n,u}} c_y,$$

where $\mathcal{L}_{n,u} = \{x \in \mathcal{L} \cap u^\perp; (x, x) = 2n\}$. Indeed, first note that by Proposition 4.1 only vectors in $\mathcal{L}_{n,u}$ can contribute to $\partial C_{n,P}$. The action of Γ on V induces an equivalence relation \sim_Γ on the set $\Gamma_P \backslash \mathcal{L}_{n,u} \subset V$ which is consequently a union of equivalence classes $[x_i] = [x_i]_P$, $1 \leq i \leq k$. We may accordingly organize the union R on the right-hand side of (4.1) as $R = \coprod_{i=1}^k \coprod_{y \in [x_i]} c_y$. But it is clear that $(\partial C_{x_i})_P = \coprod_{y \in [x_i]} c_y$ and hence we have the equality of 1-cycles in $e'(P)$ and ∂X

$$(4.2) \quad (\partial C_{x_i})_P = \sum_{y \in [x_i]_P} c_y \quad \text{and} \quad \partial C_{x_i} = \sum_{[P]} \sum_{y \in [x_i]_P} c_y,$$

since an element $y \in [x_i]$ gives rise to the lift D_y of C_{x_i} to D that intersects $e(P)$ and this intersection projects to c_y . Thus we may rewrite the right-hand side of (4.1) as $R = \coprod_{\sim_\Gamma \backslash \mathcal{L}_{n,u}} \partial C_{x_i}$. But it is clear that this latter union is $\partial C_{n,P}$ and (4.1) follows. Finally, we easily see that $\coprod_{\substack{x \in \Gamma_M \backslash \mathcal{L}_W \\ (x,x)=2n}} \coprod_{0 \leq k < \min'_{\lambda \in \Lambda_W} |(\lambda, x)|} x + ku$ is a complete set of representatives of Γ_P -equivalence classes in $\mathcal{L}_{n,u}$. These give the circles c_{x+ku} above. \square

Proposition 4.4. *Let $x \in \mathcal{L}_{n,u}$ with $n > 0$. Then there exists a rational 2-chain a_x in $e'(P)$ such that*

- (1) $\partial a_x = c_x$
- (2) $\int_{a_x} \Omega_P = 0$, here Ω_P is the area form for the fibers (see (3.1))

Proof. Except for the rationality of the cap this follows immediately from Proposition 4.2. The problem is to find a cap a_x such that $\int_{a_x} \Omega_P \in \mathbb{Q}$. We will prove this in Section 4.3 below. \square

We will define $(A_x)_P$ by $(A_x)_P = \sum_{y \in [x]} a_x$. Then sum over the components $e'(P)$ to obtain A_x a rational 2-chain in ∂X . Then we have (noting that $(\partial C_x)_P = \sum_{y \in [x]} c_y$)

$$\partial A_x = \partial C_x.$$

Definition 4.5. We define the rational absolute 2-cycle in \bar{X} by

$$C_x^c = C_x \cup (-A_x)$$

with the 2-chain A_x in $\partial \bar{X}$ as in Proposition 4.4. In particular, C_x^c defines a class in $H_2(\bar{X}) = H_2(X)$. In the same way we obtain C_n^c .

4.2. The closure of the special cycles in \tilde{X} and the cycle T_n^c . Following Hirzebruch-Zagier we let T_n be the cycle in \tilde{X} given by the closure of the cycle C_n in \tilde{X} . Hence T_n defines a class in $H_2(\tilde{X})$.

Definition 4.6. Consider the decomposition $H_2(\tilde{X}) = j_* H_2(X) \oplus (\oplus_{[P]} S_p)$, which is orthogonal with respect to the intersection pairing on \tilde{X} . We let T_n^c be the image of T_n under orthogonal projection onto the summand $j_* H_2(X)$.

Proposition 4.7. *We have*

$$j_* C_n^c = T_n^c.$$

Proof. For simplicity, we assume that X has only one cusp. The 3-manifold $e'(P)$ separates T_n and we can write $T_n = T_n \cap X^{in} + T_n \cap X^{out}$ as (appropriately oriented) 2-chains in \tilde{X} . It is obvious that we have $j_* \bar{C}_n = T_n \cap X^{in}$ as 2-chains. We write $B_n = T_n \cap X^{out}$. We have $\partial C_n = -\partial B_n$. Hence we can write $T_n = j_* C_n^c + B_n^c$, the sum of two 2-cycles in \tilde{X} . Here B_n^c is obtained by ‘capping’ B_n in $e'(P)$ with the negative of the cap A_n of C_n . Since $j_* C_n^c$ is clearly orthogonal to S_P (since it lies in X^{in}) and $B_n^c \in S_P$ (since it lies in X^{out}) the decomposition $T_n = j_* C_n^c + B_n^c$ is just the decomposition of T_n relative to the splitting $H_2(\tilde{X}) = j_* H_2(X) \oplus S_P$. Hence $T_n^c = j_* C_n^c$, as claimed. \square

4.3. Rationality of the cap. We will now prove Proposition 4.4. In fact we will show that it holds for any circle α contained in a torus fiber of $e(P)$ and passing through a rational point. We would like to thank Misha Kapovich for simplifying our original argument. The idea is to construct, for each component of ∂C_x , a 2-chain A with that component as boundary so that A is a sum $P+T+\mathcal{M}(\gamma_0)$ of three simplicial 2-chains in M . We then verify that the ‘‘parallelogram’’ P and the ‘‘triangle’’ T have rational area and the period of Ω over the ‘‘monodromy chain’’ $\mathcal{M}(\gamma_0)$ is zero.

In what follows we will pass from pictures in the plane involving directed line segments, triangles and parallelograms to identities in the space of simplicial 1-chains $C_1(T^2)$ on T^2 . The principal behind this is that any k -dimensional subcomplex S of a simplicial complex Y which is the fundamental cycle of an oriented k -submanifold $|S|$ (possibly with boundary) of Y corresponds in a *unique* way to a sum of oriented k -simplices in $C_k(Y)$.

In this subsection we will work with a general 3-manifold M with Sol geometry. Of course this includes all the manifolds $e'(P)$ that occur in this paper. Let $f \in \text{SL}(2, \mathbb{Z})$

be a hyperbolic element. We will then consider the 3-manifold M obtained from $\mathbb{R} \times T^2$ (with the 2-torus $T^2 = W/\mathbb{Z}^2$) given by the relation

$$(4.3) \quad (s, w) \sim (s + 1, f(w)).$$

We let $\pi : \mathbb{R} \times T^2 \rightarrow M$ be the resulting infinite cyclic covering.

We now define notation we will use below. We will use Greek letters to denote closed geodesics on T^2 , a subscript c will indicate that the geodesic starts at the point c on T^2 . We will use the analogous notation for geodesic arcs on W . We will use $[\alpha]$ to denote the corresponding homology class of a closed geodesic α on T^2 . If x and y are points on W we will use \overrightarrow{xy} to denote the oriented line segment joining x to y and \overline{xy} to denote the corresponding (free) vector i.e. the equivalence class of \overrightarrow{xy} modulo parallel translation.

We first take care of the fact that α does not necessarily pass through the origin. For convenience we will assume α is in the fiber over the base-point $z(x)$ corresponding to $s = 0$. Let α_0 be the parallel translate of α to the origin. Then we can find a cylinder P , image of an oriented parallelogram \tilde{P} under the universal cover $W \rightarrow T^2$ with rational vertices, such that in $Z_1(T^2, \mathbb{Q})$, the group of rational 1-cycles, we have

$$(4.4) \quad \partial P = \alpha - \alpha_0.$$

Since \tilde{P} has rational vertices we find $\int_P \Omega = \int_{\tilde{P}} \Omega \in \mathbb{Q}$.

Now we take care of the harder part of finding A as above. The key is the construction of “monodromy 2-chains”. For any closed geodesic $\gamma_0 \subset T^2$ starting at 0 we define the monodromy 2-chain $\mathcal{M}(\gamma_0)$ to be the image of the cylinder $\gamma_0 \times [0, 1] \subset T^2 \times \mathbb{R}$ in M . The reader will verify using (4.3) that in $Z_1(T^2, \mathbb{Q})$ we have

$$(4.5) \quad \partial \mathcal{M}(\gamma_0) = f^{-1}(\gamma_0) - \gamma_0.$$

Since f preserves the origin, the geodesic $f^{-1}(\gamma_0)$ is also a closed geodesic starting at the origin. Since f^{-1} is hyperbolic we have $|\text{tr}(f^{-1})| > 2$ and hence $\det(f^{-1} - I) = \det(I - f) = \text{tr}(f) - 2 \neq 0$. Put $N = \det(f^{-1} - I)$ and define $[\gamma_0] \in H_1(T^2, \mathbb{Z})$ by

$$(4.6) \quad f^{-1}([\gamma_0]) - [\gamma_0] = N[\alpha_0].$$

Note that $[\gamma_0] = N\{(f^{-1} - I)^{-1}([\alpha_0])\}$ is necessarily an integer homology class. Also note that is an equation in the first *homology*, it is not an equation in the group of 1-cycles $Z_1(T^2, \mathbb{Q})$. Since any homology class contains a unique closed geodesic starting at the origin we obtain a closed geodesic $\gamma_0 \in [\gamma_0]$ and a corresponding monodromy 2-chain $\mathcal{M}(\gamma_0)$ whence (4.5) holds in $Z_1(T^2, \mathbb{Q})$. We now solve

Problem 4.8. Find an equation in $Z_1(T^2, \mathbb{Z})$ which descends to (4.6).

Let h_1 resp. h_2 denote the covering transformation of π corresponding to the element α_0 resp. γ_0 of the fundamental group of T^2 . Define c_1 and c_2 in W by $c_1 = Nh_1(0)$ and $c_2 = h_2(0)$. Define $d \in W$ by $d = f^{-1}(c_2)$ in W . Let \tilde{T} be the oriented triangle with vertices $0, c_2, d$. Then

$$(i) \quad \pi(\overline{0c_1}) = N\alpha_0 \quad (ii) \quad \pi(\overline{0c_2}) = \gamma_0 \quad (iii) \quad \pi(\overline{0d}) = \pi(f^{-1}(\overline{0c_2})) = f^{-1}(\gamma_0).$$

We now leave it to the reader to combine the homology equation (4.6) and the three equations to show the equality of directed line segments

$$(4.7) \quad h_2(\overline{0c_1}) = \overline{c_2d}.$$

With this we can solve the problem. We see that if we consider \tilde{T} as an oriented 2-simplex we have the following equality of one chains

$$\partial\tilde{T} = \overline{0c_2} + \overline{c_2d} - \overline{0d}.$$

Let T be the image of \tilde{T} under π . Take the direct image of the previous equation under π and use equation (4.7) which implies that the second edge $\overline{c_2d}$ is equivalent under h_2 in the covering group to the directed line segment $\overline{0c_1}$ which maps to $N\alpha_0$. Hence $\overline{c_2d}$ also maps to $N\alpha_0$. We obtain the following equation in $Z_1(T^2, \mathbb{Z})$

$$(4.8) \quad \partial T = \gamma_0 + N\alpha_0 - f^{-1}(\gamma_0),$$

and we have solved the above problem. Combining (4.5) and (4.8) we have

$$\partial(\mathcal{M}(\gamma_0) + T) = f^{-1}(\gamma_0) - \gamma_0 + \gamma_0 + \alpha_0 - f^{-1}(\gamma_0) = N\alpha_0.$$

Combining this with (4.4) and setting where $A_0 = \mathcal{M}(\gamma_0) + T$ we obtain

$$(4.9) \quad \partial(NP + A_0) = N\alpha,$$

in $Z_1(M, \mathbb{Z})$. Hence if we define A to be the *rational* chain $A = \frac{1}{N}(NP + A_0) = P + \frac{1}{N}T + \frac{1}{N}\mathcal{M}(\gamma_0)$ in M we have the following equation in $Z_1(M, \mathbb{Q})$:

$$\partial A = \alpha.$$

Finally, the integral of Ω over A is rational. Indeed, the integral over P is rational. Since all vertices of \tilde{T} are integral the area of \tilde{T} is integral, the integral of Ω over T is integral. Thus it suffices to observe that the restriction of Ω to $\mathcal{M}(c)$ is zero. With this we have completed the proof of Proposition 4.4.

4.4. Linking numbers in Sol. In the introduction we defined the linking number of two disjoint homologically trivial 1-cycles a and b in a closed 3-manifold M as $\text{Lk}(a, b) = \langle A, b \rangle$, where A is any rational 2-chain in M with boundary a . Since b defines a trivial homology class in M , the link is well-defined, ie, does not depend on the choice of A .

We let M be the Sol manifold as before realized as in Section 4.3 via (4.3) and consider the case when a and b are two contained in two torus fibers. Then by the previous section they are homologically trivial. If a and b are contained in the same fiber we move b to the right (i.e. in the direction of positive s) to a nearby fiber. We take $a, b \in H_1(T^2, \mathbb{Z})$, and in this section we are allowed to confuse a and b with their representatives in the lattice \mathbb{Z}^2 and the unique closed geodesic in T^2 passing through the origin that represents them. We will write for the image of a and b in $\mathbb{R} \times T^2$ and M $a = a(0) = 0 \times a$ and $b = b(\epsilon) = \epsilon \times b$. Our goal is to compute the linking number $Lk(a, b(\epsilon))$. By the explicit construction of the cap A in Section 4.3 we obtain

Lemma 4.9.

$$Lk(a, b(\epsilon)) = M(c) \cdot b(\epsilon) = c(\epsilon) \cdot b(\epsilon) = c \cdot b.$$

Here c is the rational one cycle obtained by solving $(f^{-1} - I)(c) = a$ and $M(c)$ is the (rational) monodromy 2-chain associated to c (see above) with boundary $\partial M(c) = (f^{-1} - I)(c) = a$. Here the first \cdot is the intersection of chains in M , the next \cdot is the intersection number of 1-cycles in the fiber $\epsilon \times T^2$ and the last \cdot is the intersection number of 1-cycles in $0 \times T^2$.

Noting that this last intersection number coincides with the intersection number of the underlying homology classes which in term coincides with the symplectic form $\langle \cdot, \cdot \rangle$ on $H_1(T^2, \mathbb{Q})$ we have found our desired formula for the linking number.

Theorem 4.10. $Lk(a, b(\epsilon)) = \langle (f^{-1} - I)^{-1}(a), b \rangle$.

It is a remarkable fact that there is a simple formula involving only the action of the glueing homeomorphism $f \in \mathrm{SL}(2, \mathbb{Z})$ on $H_1(T^2, \mathbb{Z})$ for linking numbers for 1-cycles contained in fiber tori T^2 of in Sol (unlike the case of linking numbers in \mathbb{R}^3).

This immediately leads to an explicit formula for the numbers $Lk(\partial C_n, \partial C_m)$. Using Lemma 4.3 we obtain

Theorem 4.11. Let $g = (f^{-1} - I)^{-1}$. Then

$$Lk((\partial C_n)_P, (\partial C_m)_P) = \sum_{\substack{x \in \Gamma_M \setminus \mathcal{L}_W \\ (x, x) = 2n}} \sum_{\substack{x' \in \Gamma_M \setminus \mathcal{L}_W \\ (x', x') = 2m}} (\min'_{\lambda \in \Lambda_W} |(\lambda, x)|) (\min'_{\mu \in \Lambda_W} |(\mu, x')|) \langle g(Jx), Jx' \rangle.$$

Here Jx is properly oriented primitive vector in Λ_W such that $(Jx, x) = 0$.

Example 4.12. We consider the integral skew Hermitian matrices in Example 2.1. Let $u = \begin{pmatrix} \sqrt{p} & 0 \\ 0 & 0 \end{pmatrix}$, so that $W = \{ \begin{pmatrix} 0 & \lambda \\ -\lambda' & 0 \end{pmatrix}; \lambda \in K \} \simeq K$. The symplectic form on K is given by $\langle \lambda, \mu \rangle = \frac{1}{\sqrt{p}}(\lambda\mu' - \lambda'\mu)$. The action of the unipotent radical $N = \{n(\lambda) = \begin{pmatrix} 1 & \lambda \\ 0 & 1 \end{pmatrix}\}$ on a vector $\mu \in K$ is now slightly different, namely, $n(\lambda)\mu = \mu + \langle \lambda, \mu \rangle u$. Hence in these coordinates, ∂C_μ is given by the image of the line $\mathbb{R}\mu = \{ \lambda \in K_{\mathbb{R}}; \langle \lambda, \mu \rangle = 0 \}$, and $(\min'_{\lambda \in \mathcal{O}_K} |(\lambda, \mu)|)\mu$ is a primitive generator in \mathcal{O}_K for that line. We let ϵ be a generator of U_+ , the totally positive units in \mathcal{O}_K , and we assume that the glueing map f is realized by multiplication with ϵ' . For $d \equiv 1 \pmod{4}$ a prime and $m = 1$, C_1 has only component arising from $x = 1 \in K$ and $C_1 \simeq \mathrm{SL}_2(\mathbb{Z}) \backslash \mathbb{H}$. Then Theorem 4.11 becomes (the \min' -term is now wrt $\langle \cdot, \cdot \rangle$)

$$Lk((\partial C_n)_P, (\partial C_1)_P) = 2 \sum_{\substack{\mu \in U_+ \setminus \mathcal{O}_K \\ \mu\mu' = n, \mu \gg 0}} \left\langle \frac{\mu}{\epsilon - 1}, 1 \right\rangle = 2 \sum_{\substack{\mu \in U_+ \setminus \mathcal{O}_K \\ \mu\mu' = n, \mu \gg 0}} = \frac{2}{\sqrt{p}} \frac{\mu + \mu'\epsilon}{\epsilon - 1}.$$

This is (twice) the “boundary contribution” in [16], Section 1.4, see also Section 7.5.

5. SCHWARTZ FUNCTIONS AND FORMS

Let U be a non-degenerate rational quadratic space of signature (p, q) and even dimension m . We will later apply the following to $U = V$ and $U = W$. Changing notation from before, we let $G = \mathrm{SO}_0(U_{\mathbb{R}})$ with maximal compact subgroup K and write $D = G/K$ for the associated symmetric space. We let $\mathcal{S}(U_{\mathbb{R}})$ be the space of Schwartz functions on $U_{\mathbb{R}}$ on which $\mathrm{SL}_2(\mathbb{R})$ acts via the Weil representation ω .

5.1. Extending certain Schwartz functions to functions of $\tau \in \mathbb{H}$ and $z \in D$. Let $\varphi \in \mathcal{S}(U_{\mathbb{R}})$ be an eigenfunction under the maximal compact $\mathrm{SO}(2)$ of $\mathrm{SL}_2(\mathbb{R})$ of weight r . Define $g'_\tau \in \mathrm{SL}_2(\mathbb{R})$ by $g'_\tau = \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix} \begin{pmatrix} v^{1/2} & 0 \\ 0 & v^{-1/2} \end{pmatrix}$. Then we have $\omega(g'_\tau)\varphi(x) = v^{m/4}\varphi(\sqrt{v}x)e^{\pi i(x,x)u}$. Accordingly we define

$$(5.1) \quad \varphi(x, \tau) = v^{-r/2}\omega(g'_\tau)\varphi(x) = v^{-r/2+m/4}\varphi^0(\sqrt{v}x)e^{\pi i(x,x)\tau}.$$

Here we have also defined $\varphi^0(x) = \varphi(x)e^{\pi(x,x)}$. Let E be a G -module and let $g_z \in G$ be any element that carries the basepoint z_0 in D to $z \in D$. Then define for $\varphi \in [\mathcal{S}(U_{\mathbb{R}}) \otimes E]^K$, the E -valued K -invariant Schwartz functions on $U_{\mathbb{R}}$, the functions $\varphi(x, z)$ and $\varphi(x, \tau, z)$ for $x \in U, z \in D, \tau \in \mathbb{H}$ by

$$\varphi(x, z) = g_z\varphi(g_z^{-1}x) \quad \text{and} \quad \varphi(x, \tau, z) = g_z\varphi(g_z^{-1}x, \tau).$$

We will continue to use these notational conventions for other (not necessarily Schwartz) functions that arise in this paper.

5.2. Schwartz forms for V . Let \mathfrak{g} be the Lie algebra of G and $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ be the Cartan decomposition of \mathfrak{g} associated to K . We identify $\mathfrak{g} \simeq \bigwedge^2 V_{\mathbb{R}}$ as usual via $(v_1 \wedge v_2)(v) = (v_1, v)v_2 - (v_2, v)v_1$. We write $X_{ij} = e_i \wedge e_j \in \mathfrak{g}$ and note that \mathfrak{p} is spanned by X_{ij} with $1 \leq i \leq 2$ and $3 \leq j \leq 4$. We write ω_{ij} for their dual. We orient D such that $\omega_{13} \wedge \omega_{14} \wedge \omega_{23} \wedge \omega_{24}$ gives rise to the G -invariant volume element on D .

5.2.1. Special forms for V . The Kudla-Millson form φ_2 is an element in

$$[\mathcal{S}(V_{\mathbb{R}}) \otimes \mathcal{A}^2(D)]^G \simeq [\mathcal{S}(V_{\mathbb{R}}) \otimes \bigwedge^2 \mathfrak{p}^*]^K,$$

where the isomorphism is given by evaluation at the base point. Here $\mathcal{A}^2(D)$ denotes the differential 2-forms on D . Note that G acts diagonally in the natural fashion. At the base point φ_2 is given by

$$\varphi_2 = \frac{1}{2} \prod_{\mu=3}^4 \sum_{\alpha=1}^2 \left(x_\alpha - \frac{1}{2\pi} \frac{\partial}{\partial x_\alpha} \right) \varphi_0 \otimes \omega_{\alpha\mu}.$$

Here $\varphi_0(x) := e^{-\pi(x,x)_0}$, where $(x, x)_0 = \sum_{i=1}^4 x_i^2$ is the minimal majorant associated to the base point in D . Note that φ_2 has weight 2, see [21]. There is another Schwartz form ψ_1 of weight 0 which lies in $[\mathcal{S}(V_{\mathbb{R}}) \otimes \mathcal{A}^1(D)]^G \simeq [\mathcal{S}(V_{\mathbb{R}}) \otimes \mathfrak{p}^*]^K$ and is given by

$$(5.2) \quad \psi_1 = -x_1x_3\varphi_0(x) \otimes \omega_{14} + x_1x_4\varphi_0(x) \otimes \omega_{13} - x_2x_3\varphi_0(x) \otimes \omega_{24} + x_2x_4\varphi_0(x) \otimes \omega_{23}.$$

The key relationship is (see [24], §8)

Theorem 5.1.

$$\omega(L)\varphi_2 = d\psi_1.$$

Here $\omega(L)$ is the Weil representation action of the SL_2 -lowering operator $L = \frac{1}{2} \begin{pmatrix} 1 & -i \\ -i & -1 \end{pmatrix} \in \mathfrak{sl}_2(\mathbb{R})$ on $\mathcal{S}(V_{\mathbb{R}})$, while d denotes the exterior differentiation on D .

On the upper half plane \mathbb{H} , the action of L corresponds to the action of the classical Maass lowering operator which we also denote by L . For a function f on \mathbb{H} , we have

$$Lf = -2iv^2 \frac{\partial}{\partial \bar{\tau}} f.$$

When made explicit using (5.1) Theorem 5.1 translates to

$$(5.3) \quad v \frac{\partial}{\partial v} \varphi_2^0(\sqrt{v}x) = d(\psi_1^0(\sqrt{v}x)).$$

5.2.2. *The singular form $\tilde{\psi}_1$.* We define the singular form $\tilde{\psi}_1$ by

$$(5.4) \quad \tilde{\psi}_1(x) = - \left(\int_1^\infty \psi_1^0(\sqrt{r}x) \frac{dr}{r} \right) e^{-\pi(x,x)} = - \frac{1}{2\pi(x_3^2 + x_4^2)} \psi_1(x).$$

for $x \neq 0$, and as before $\tilde{\psi}_{2,0}^0(x) = \tilde{\psi}_1(x)e^{\pi(x,x)}$ and $\tilde{\psi}_1(x, z)$. We see that $\tilde{\psi}_1$ is defined for $x \notin \text{span}[e_3, e_4]^\perp$. Formulated differently, $\tilde{\psi}_1(x, z)$ for fixed x is defined for $z \notin D_x$. Furthermore, as if $\tilde{\psi}_1$ was a Schwartz function of weight 2, we define

$$(5.5) \quad \tilde{\psi}_1(x, \tau, z) = \tilde{\psi}_1^0(\sqrt{v}x, z) e^{\pi i(x,x)\tau} = - \left(\int_v^\infty \psi_1^0(\sqrt{r}x, z) \frac{dr}{r} \right) e^{\pi i(x,x)\tau}.$$

Proposition 5.2. $\tilde{\psi}_1(x, z)$ is a differential 1-form with singularities along D_x . Outside D_x , we have

$$d\tilde{\psi}_1(x, z) = \varphi_2(x, z).$$

Here d denotes the exterior differentiation on D . In particular, for $(x, x) \leq 0$, we see that $\varphi_2(x)$ is exact. Furthermore,

$$L\tilde{\psi}_1(x, \tau) = \psi_1(x, \tau).$$

Proof. Using (5.4) and (5.3), we see

$$d\tilde{\psi}_1^0(x, z) = - \int_1^\infty d(\psi_1^0(\sqrt{r}x, z)) \frac{dr}{r} = - \int_1^\infty \frac{\partial}{\partial r} (\varphi_2^0(\sqrt{r}x, z)) \frac{dr}{r} = \varphi_2^0(x, z),$$

as claimed. The formula $L\tilde{\psi}_1(x, \tau) = \psi_1(x, \tau)$ follows easily from (5.5). \square

Remark 5.3. The construction of the singular form $\tilde{\psi}$ works in much greater generality for $O(p, q)$ whenever we have two Schwartz forms ψ and φ (of weight $r - 2$ and r resp.) such that

$$d\psi = L\varphi.$$

Then the analogous construction of $\tilde{\psi}$ then immediately yields $d\tilde{\psi} = \varphi$ outside a singular set. The main example for this are the general Kudla-Millson forms φ_q and ψ_{q-1} , see [24]. For these forms, this construction is already implicit in [5]. In particular, the proof of Theorem 7.2 in [5] shows that $\tilde{\psi}$ gives rise to a differential character for the analogous cycle C_x , see also Section 8 of this paper. The unitary case will be considered in [10].

5.3. **Schwartz forms for W .** Let $W \subset V$ be the rational quadratic space of signature $(1, 1)$ obtained from the Witt decomposition of V . We will refer to the nullcone of W as the light-cone. We write $\mathfrak{m} \simeq \mathbb{R}$ for the Lie algebra of $M = \text{SO}_0(W_\mathbb{R})$. Then $X_{23} = e_2 \wedge e_3$ is its natural generator with dual ω_{23} . We identify the associated symmetric space D_W to M with the space of lines in $W_\mathbb{R}$ on which the bilinear form $(,)$ is negative definite:

$$D_W = \{\mathfrak{s} \subset W_\mathbb{R}; \dim \mathfrak{s} = 1 \text{ and } (,)|_{\mathfrak{s}} < 0\}.$$

We pick as base point of D_W the line \mathfrak{s}_0 spanned by e_3 . We set $x(s) := m(s)e_3 = \sinh(s)e_2 + \cosh(s)e_3$. This realizes the isomorphism $D_W \simeq \mathbb{R}$. Namely, $\mathfrak{s} = \text{span } x(s)$. Accordingly, we frequently write s for \mathfrak{s} and vice versa. A vector $x \in W$ of positive length defines a point $D_{W,x}$ in D via $D_{W,x} = \{\mathfrak{s} \in D; \mathfrak{s} \perp x\}$. So $\mathfrak{s} = D_{W,x}$ if and only if $(x, x(\mathfrak{s})) = 0$. We also write $\mathfrak{s}(x) = D_{W,x}$.

5.3.1. *Special forms for W .* We carry over the conventions from section 5.1. We first consider the Schwartz form $\varphi_{1,1}$ on $W_{\mathbb{R}}$ constructed in [12] (in much greater generality) with values in $\mathcal{A}^1(D_W) \otimes W_{\mathbb{C}}$. More precisely,

$$\varphi_{1,1} \in [\mathcal{S}(W_{\mathbb{R}}) \otimes \mathcal{A}^1(D_W) \otimes W_{\mathbb{C}}]^M \simeq [\mathcal{S}(W_{\mathbb{R}}) \otimes \mathfrak{m}^* \otimes W_{\mathbb{C}}],$$

Here M acts diagonally on all three factors. Explicitly at the base point, we have

$$\varphi_{1,1}(x) = \frac{1}{2^{3/2}} \left(4x_2^2 - \frac{1}{\pi} \right) e^{-\pi(x_2^2 + x_3^2)} \otimes \omega_{23} \otimes e_2.$$

Note that $\varphi_{1,1}$ has weight 2, see [12], Theorem 6.2. We define $\varphi_{1,1}(x, s)$ and $\varphi_{1,1}^0$ as before. There is another Schwartz function $\psi_{0,1}$ of weight 0 given by

$$\begin{aligned} \psi_{0,1}(x) &= -\frac{1}{\sqrt{2}} x_2 x_3 e^{-\pi(x_2^2 + x_3^2)} \otimes 1 \otimes e_2 + \frac{1}{4\sqrt{2}\pi} e^{-\pi(x_2^2 + x_3^2)} \otimes 1 \otimes e_3 \\ &\in [\mathcal{S}(W_{\mathbb{R}}) \otimes \bigwedge^0 \mathfrak{m}^* \otimes W_{\mathbb{C}}], \end{aligned}$$

and also $\psi_{0,1}(x, s)$ and $\psi_{0,1}^0$. Note that the notation differs from [12], section 6.5. The function $\psi_{0,1}$ defined here is the term $-\psi_{1,1} - \frac{1}{2}\Lambda_{1,1}$ given in Theorem 6.11 in [12]. The key relation between $\varphi_{1,1}$ and $\psi_{0,1}$ (correcting a sign mistake in [12]) is given by

Theorem 5.4. ([12], Theorem 6.2)

$$\omega(L)\varphi_{1,1} = d\psi_{0,1}.$$

When made explicit, we have, again using (5.1),

$$(5.6) \quad v^{3/2} \frac{\partial}{\partial v} (v^{-1/2} \varphi_{1,1}^0(\sqrt{v}x, s)) = d(\psi_{0,1}^0(\sqrt{v}x, s)).$$

5.3.2. *The singular Schwartz function $\tilde{\psi}_{0,1}$.* In the same way as for V we define

$$(5.7) \quad \tilde{\psi}_{0,1}(x) = - \left(\int_1^\infty \psi_{0,1}^0(\sqrt{r}x) r^{-3/2} dr \right) e^{-\pi(x,x)}$$

for all $x \in W$, including $x = 0$. Define $\tilde{\psi}_{0,1}^0(x)$, $\tilde{\psi}_{0,1}^0(x, s)$ as before and also

$$\tilde{\psi}_{0,1}(x, \tau, s) = v^{-1/2} \tilde{\psi}_{0,1}^0(\sqrt{v}x, s) e^{\pi i(x,x)\tau} = - \left(\int_v^\infty \psi_{0,1}^0(\sqrt{r}x, s) r^{-3/2} dr \right) e^{\pi i(x,x)\tau}.$$

Note that $\tilde{\psi}_{0,1}(x, s)$ has a singularity at $D_{w,x}$. Define functions A and B by

$$\tilde{\psi}_{0,1}(x) = A(x) \otimes 1 \otimes e_2 + B(x) \otimes 1 \otimes e_3$$

and note

$$(5.8) \quad -X_{23}B(x) = A(x).$$

We extend these functions to D_W as before. We see by integrating by parts

Lemma 5.5.

$$A(x) = \frac{1}{2\sqrt{\pi}} x_2 \frac{x_3}{|x_3|} \Gamma\left(\frac{1}{2}, 2\pi x_3^2\right) e^{-\pi(x,x)}$$

$$B(x) = -\frac{1}{2\sqrt{2}\pi} e^{-\pi(x_2^2+x_3^2)} + \frac{1}{2\sqrt{\pi}} |x_3| \Gamma\left(\frac{1}{2}, 2\pi x_3^2\right) e^{-\pi(x,x)}.$$

Here $\Gamma\left(\frac{1}{2}, a\right) = \int_a^\infty e^{-u} u^{-1/2} du$ is the incomplete Γ -function at $s = 1/2$.

It is now immediate that B is continuous and bounded on D_W . Since A is clearly bounded we find that A and B are locally integrable on D_W and integrable and square-integrable on W . The singularities of A and B are given as follows.

Lemma 5.6. (i) $B(x) - (1/2)|x_3|e^{-\pi(x,x)}$ is C^2 on the Minkowski plane W .
 (ii) $A(x) - (1/2)x_2 \frac{x_3}{|x_3|} e^{-\pi(x,x)}$ is C^1 on the Minkowski plane W .

Proof. Use Lemma 5.5, expand the incomplete gamma function around $x_3 = 0$, and observe that $|x|x^n$ is C^n for $n > 0$. \square

The key properties of $\tilde{\psi}_{0,1}$ analogous to Lemma 5.2 are given by

Lemma 5.7. Outside $D_{W,x}$,

$$d\tilde{\psi}_{0,1}(x, s) = \varphi_{1,1}(x, s) \quad \text{and} \quad L\tilde{\psi}_{0,1}(x, \tau) = \psi_{0,1}(x, \tau).$$

5.3.3. *The singular function $\tilde{\psi}'_{0,1}$.* Inspired by [16], section 2.3, we define a functions $A'(x)$ and $B'(x)$ on W by

$$(5.9) \quad B'(x) = \begin{cases} \frac{1}{2} \min(|x_2 - x_3|, |x_2 + x_3|) e^{-\pi(x,x)} & \text{if } x_2^2 - x_3^2 > 0, \\ 0 & \text{otherwise,} \end{cases}$$

$$A'(x) = -X_{23}B'(x) = -\text{sgn}(x_2x_3)B'(x).$$

Lemma 5.8. (i) $B'(x) + \frac{1}{2}|x_3|e^{-\pi(x,x)}$ is C^2 on the complement of the light-cone in W and C^2 on nonzero M -orbits.

(ii) $A'(x) + \frac{1}{2}x_2 \frac{x_3}{|x_3|} e^{-\pi(x,x)}$ is C^1 on the complement of the light-cone in W and C^1 on nonzero M -orbits.

We define $\tilde{\psi}'_{0,1}$ by

$$\tilde{\psi}'_{0,1}(x) = A'(x) \otimes 1 \otimes e_2 + B'(x) \otimes 1 \otimes e_3$$

and $\tilde{\psi}'_{0,1}(x, \tau, s) = v^{-1/2} m(s) \tilde{\psi}'_{0,1}(m^{-1}(s)\sqrt{v}x) e^{\pi i(x,x)\tau}$. A little calculation shows that $\tilde{\psi}'_{0,1}(x)$ is locally constant on D_W with a singularity at $D_{W,x}$ and holomorphic in τ :

Lemma 5.9. Outside $D_{W,x}$ we have

$$d\tilde{\psi}'_{0,1}(x) = 0 \quad \text{and} \quad L\tilde{\psi}'_{0,1}(x, \tau) = 0.$$

Remark 5.10. The functions $\tilde{\psi}_{0,1}(x)$ and $\tilde{\psi}'_{0,1}(x)$ define currents on D_W . One can show, similarly to Section 6.5, that for $(x, x) > 0$ we have

$$d[\tilde{\psi}_{0,1}(x)] = \delta_{D_{W,x} \otimes x} + [\varphi_{1,1}(x)], \quad d[\tilde{\psi}'_{0,1}(x)] = -\delta_{D_{W,x} \otimes x},$$

where $D_{W,x} \otimes x$ is the 0-cycle $D_{W,x}$ ‘with coefficient $x \in W$ ’ defined in [12].

5.3.4. *The form $\phi_{0,1}$ on W .* We now combine $\tilde{\psi}_{0,1}$ and $\tilde{\psi}'_{0,1}$ to obtain an integrable and also square-integrable W -valued function

$$\phi_{0,1} \in [L^2(W_{\mathbb{R}}) \otimes \bigwedge^0 \mathfrak{m}^* \otimes W_{\mathbb{C}}]$$

by

$$\phi_{0,1}(x) = \tilde{\psi}_{0,1}(x) + \tilde{\psi}'_{0,1}(x)$$

and then also $\phi_{0,1}(x, s)$. Combining Lemmas 5.6, 5.8 and (5.8), (5.9) we obtain

Proposition 5.11. (i) $B(x) + B'(x)$ is C^2 on the complement of the light-cone in W and C^2 on nonzero M -orbits.

(ii) $A(x) + A'(x)$ is C^1 on the complement of the light-cone in W and C^1 on nonzero M orbits.

(iii) $X_{23}(B + B') = -(A + A')$ on all of W .

So for given x , the function $\phi_{0,1}(x, s)$ is a C^1 -function on D_W with values in $W_{\mathbb{C}}$.

The following theorem is fundamental for us. It is an immediate consequence of the Lemmas 5.7 and 5.9.

Theorem 5.12. *The form $\varphi_{1,1}$ on D_W is exact. Namely,*

$$d\phi_{0,1} = \varphi_{1,1}.$$

Furthermore,

$$L\phi_{0,1} = d\psi_{0,1}.$$

Proposition 5.13. *The function $\phi_{0,1}$ is an eigenfunction of $K' = \text{SO}(2)$ of weight 2 under the Weil representation. More precisely,*

$$\omega(k')\phi_{0,1} = \chi^2(k')\phi_{0,1},$$

where χ is the standard character of $\text{SO}(2) \simeq U(1)$.

Proof. It suffices to show this for one component of $\phi_{0,1}$, that is, the function $B(x) + B'(x)$. Then the assertion has been already proved in §2.3 by showing that $B(x) + B'(x)$ is an eigenfunction under the Fourier transform. We give here an infinitesimal proof. Since $\omega(k')$ acts essentially as Fourier transform and $B + B'$ is L^1 , we see that $\omega(k')(B + B')$ is continuous. Hence it suffices to establish the corresponding current equality $[\omega(k')(B + B')] = \chi^2(k')[B + B']$, since continuous functions coincide when they induce the same current. The infinitesimal generator of K' acts by $H := \frac{-i}{4\pi} \left(\frac{\partial^2}{\partial x_2^2} - \frac{\partial^2}{\partial x_3^2} \right) + \pi i(x_2^2 - x_3^2)$, and a straightforward calculation immediately shows

$$HB' = 2iB' \quad \text{and} \quad HB = 2iB,$$

outside the singularity $x_2^2 - x_3^2 = 0$. Now we consider the currents $H[B]$ and $H[B']$. An easy calculation using that B and B' are C^2 up to $|x_3|e^{-\pi(x_2^2 - x_3^2)}$ shows that for a test function f on W we have

$$\begin{aligned} H[B](f) &= [HB](f) + \int_0^\infty e^{-\pi x_2^2} f(x_2, 0) dx_2, \\ H[B'](f) &= [HB'](f) - \int_0^\infty e^{-\pi x_2^2} f(x_2, 0) dx_2. \end{aligned}$$

Thus $H[B + B'] = [H(B + B')] = 2i[B + B']$ as claimed. \square

5.3.5. *The map ι_P .* We define a map

$$\iota_P : \mathcal{S}(W_{\mathbb{R}}) \otimes \bigwedge^i \mathfrak{m}^* \otimes W_{\mathbb{C}} \rightarrow \mathcal{S}(W_{\mathbb{R}}) \otimes \bigwedge^{i+1} (\mathfrak{m}^* \oplus \mathfrak{n}^*)$$

by

$$\iota_P(\varphi \otimes \omega \otimes w) = \varphi \otimes ((\omega \wedge (w \wedge u'))).$$

Here we used the isomorphism $\mathfrak{n} \simeq W \wedge \mathbb{R}u \in \bigwedge^2 V_{\mathbb{R}} \simeq \mathfrak{g}$ and identify W with its dual via the bilinear form $(,)$ so that $\mathfrak{n}^* \simeq W \wedge \mathbb{R}u'$. In [13], Section 6.2 we explain that ι_P is a map of Lie algebra complexes. Hence we obtain a map of complexes

$$[\mathcal{S}(W_{\mathbb{R}}) \otimes \mathcal{A}^i(D_W) \otimes W_{\mathbb{C}}]^M \rightarrow [\mathcal{S}(W_{\mathbb{R}}) \otimes \mathcal{A}^{i+1}(e(P))]^{NM},$$

which we also denote by ι_P . Here N acts trivially on $\mathcal{S}(W_{\mathbb{R}})$. Explicitly, the vectors e_2 and e_3 in W map under ι_P to the left-invariant 1-forms

$$e_2 \mapsto \cosh(s)dw_2 - \sinh(s)dw_3 \quad e_3 \mapsto \sinh(s)dw_2 - \cosh(s)dw_3$$

with the coordinate functions w_2, w_3 on W defined by $w = w_2e_2 + w_3e_3$. We apply ι_P to the forms on W of this section, and we obtain $\varphi_{1,1}^P, \phi_{0,1}^P, \psi_{0,1}^P$, and $\psi'_{0,1}^P$.

6. THE BOUNDARY THETA LIFT AND LINKING NUMBERS IN SOL

6.1. **Global theta functions for W .** We let \mathcal{L}_W be a Γ_P -invariant (coset of a) lattice in W , where Γ_N acts trivially on W . For $\varphi_{1,1}$, we define its theta function by

$$\theta_{\varphi_{1,1}}(\tau, \mathcal{L}_W) = \sum_{x \in \mathcal{L}_W} \varphi_{1,1}(x, \tau)$$

and similarly for $\psi_{0,1}$, and $\phi_{0,1}$. Then the usual theta machinery gives that $\theta_{\varphi_{1,1}}(\tau, \mathcal{L}_W)$ and $\theta_{\phi_{0,1}}(\mathcal{L}_W)$ both transform like (non)-holomorphic modular forms of weight 2 for some congruence subgroup of $\mathrm{SL}_2(\mathbb{Z})$.

Remark 6.1. The claim is not obvious for $\theta_{\phi_{0,1}}$, since $\phi_{0,1}$ is not a Schwartz function. In that case, we use Proposition 5.11. The component $B + B'$ of $\phi_{0,1}$ is C^2 outside the light cone. Since W is anisotropic we can then apply Poisson summation, and this component transforms like a modular form. Then apply the differential operator X_{23} to obtain the same for the other component $A + A'$ of $\phi_{0,1}$.

In fact, if W is isotropic and \mathcal{L}_W intersects non-trivially with the light cone, then $\theta_{\phi_{0,1}}$ is not quite a modular form. The case, when the \mathbb{Q} -rank of V is 2 is interesting in its own right. We will discuss this elsewhere.

Via the map ι_P from Section 5.3.5 we can view all theta functions for W as functions resp. differential forms on $e'(P)$. We set $\theta_{\varphi_{1,1}}^P = \theta_{\varphi_{1,1}^P}$, and similarly $\theta_{\psi_{0,1}}^P$ and $\theta_{\phi_{0,1}}^P$. Since ι_P is a map of complexes we immediately see by Theorem 5.4 and Theorem 5.12

Proposition 6.2.

$$L\theta_{\varphi_{1,1}}^P = d\theta_{\psi_{0,1}}^P \quad \text{and} \quad L\theta_{\phi_{0,1}}^P = \theta_{\psi_{0,1}}^P.$$

We now interpret the (holomorphic) Fourier coefficients of the boundary theta lift associated to $\theta_{\varphi}^P(\tau, \mathcal{L}_W)$. They are given by linking numbers. We have

Theorem 6.3. *Let c a homological trivial 1-cycle in $e'(P)$ which is disjoint from the torus fibers containing components of ∂C_n or for $c = \partial C_y$ for C_y one of the components of C_n , we have*

$$\int_c \theta_{\phi_{0,1}}^P(\tau, \mathcal{L}_{W_P}) = \sum_{n=1}^{\infty} \text{Lk}((\partial C_n)_P, c) q^n + \sum_{n \in \mathbb{Q}} \int_c \tilde{\psi}_{0,1}^P(n)(\tau).$$

So the Fourier coefficients of the holomorphic part of $\int_c \theta_{\phi}^P(\tau, \mathcal{L}_{W_P})$ are the linking numbers of the cycles c and ∂C_n at the boundary component $e'(P)$.

Theorem 6.3 follows from $\phi_{0,1} = \tilde{\psi}_{0,1} + \tilde{\psi}'_{0,1}$ combined with Theorem 6.7 below.

Example 6.4. In the situation of Examples 2.1 and 4.12, we obtain

$$\int_{\partial C_1} \theta_{\phi_{0,1}}^P(\tau, \mathcal{L}_{W_P}) = \frac{1}{\sqrt{2d}} \sum_{\substack{\lambda \in \mathcal{O}_K \\ \lambda \lambda' > 0}} \min(|\lambda|, |\lambda'|) e^{-2\pi \lambda \lambda' \tau} - \frac{\sqrt{2}}{\sqrt{dv}} \sum_{\lambda \in \mathcal{O}_K} \beta(\pi v (\lambda - \lambda')^2) e^{-2\pi \lambda \lambda' \tau},$$

where $\beta(s) = \frac{1}{16\pi} \int_1^{\infty} e^{-st} t^{-3/2} dt$. This is (up to a constant) exactly Zagier's function $\mathcal{W}(\tau)$ in [16], §2.3.

6.2. Linking numbers, de Rham cohomology and linking duals. We begin with a general discussion of integral formulas for linking numbers. Such formulas go back to the classical Gauss-Ampère formula for \mathbb{R}^3 , see [8], p.79-81, and [7] for its generalization to S^3 and H^3 . Suppose now that c is a 1-cycle in an oriented compact 3-manifold M that is a rational boundary and U is a tubular neighborhood of c .

Definition 6.5. We will say any closed form β in $M - U$ is a *linking dual* (relative to U) of the bounding 1-cycle c if for any 1-cycle a in $M - U$ which is a rational boundary in M we have

$$\int_a \beta = \text{Lk}(a, c).$$

We will prove that given a cycle c that bounds rationally then linking duals for c exist for all tubular neighborhoods U of c . Let η be a Thom form for c compactly supported in U . This means that η is closed and has integral 1 over any normal disk to c . Let η_M be the extension of η to M by zero. It is standard in topology (the extension of the Thom class by zero is the Poincaré dual of the zero section of the normal bundle) that the form η_M represents the 2-dimensional cohomology class on M which is Poincaré dual to c . Since c is a rational boundary there exists a 1-form β on M such that $d\beta = \eta_M$. We will now see that β is a linking dual of c . To this end, suppose a is a 1-cycle in $M - U$ which is a rational boundary in M , hence there exists a rational chain A with $\partial A = a$. We may suppose η_M vanishes in a neighborhood V of a which is disjoint from U . Then the restriction η_{M-V} of η_M to $M - V$ represents the (relative) Poincaré dual of the absolute cycle c in $(M - V, \partial(M - V))$. Using this the reader will show that

$$\int_A \eta_M = \int_{A \cap (M-V)} \eta_{M-V} = A \cdot c = \text{Lk}(a, c).$$

Note that restriction of β to $M - U$ is closed. Then

$$\int_a \beta = \int_A \eta_M = \text{Lk}(a, c).$$

Hence we have

Proposition 6.6. *β is a linking dual of c .*

6.3. The 1-form $e^{2\pi n}\tilde{\psi}'_{0,1}(n)$ is a linking dual of $(\partial C_n)_P$. We now return to the case in hand. In what follows, we drop subscript and superscript P 's since we are fixing a boundary component $e(P)$. We let F_n be the union of the fibers containing components of ∂C_n , and we let F_x be the fiber containing c_x . Recall that c_x is the image of $D_x \cap e(P)$ in $e'(P)$.

Theorem 6.7. *Let $n > 0$. The 1-form $e^{2\pi n}\tilde{\psi}'_{0,1}(n)$ is a linking dual for ∂C_n in $e'(P)$ relative to any neighborhood U of F_n . Hence, for c a rational 1-boundary in $e'(P)$ which is disjoint from F_n we have*

$$(6.1) \quad \int_c \tilde{\psi}'_{0,1}(n) = \text{Lk}(\partial C_n, c)e^{-2\pi n}.$$

Furthermore, (6.1) holds when $c = c_y$ contained in one fiber F_x of ∂C_n .

We will first deal with the case in which c is disjoint from F_n (which we will refer to in what follows as case (i)), then at the end of this section we will reduce the case in which $c = c_y$ (which we will refer to as case (ii)) to case (i) by a Stokes' Theorem argument. Thus we will now assume we are in case (i).

The key step is

Proposition 6.8. *Let $n > 0$ and let η be an exact 2-form in $e'(P)$ which is compactly supported in the complement of F_n . Then*

$$(6.2) \quad \int_{e'(P)} \eta \wedge \tilde{\psi}'_{0,1}(n) = \left(\int_{A_n} \eta \right) e^{-2\pi n}.$$

Remark 6.9. Note that (6.2) also holds in case $\eta = \Omega_P$. In this case the right-hand side is zero by the normalization of the cap A_n and the left-hand side is zero because $\Omega \wedge \tilde{\psi}'_{0,1}(n) = 0$ since Ω has bidegree $(0, 2)$ and $\tilde{\psi}'_{0,1}(n)$ has bidegree $(0, 1)$ (here we use the obvious base/fiber bigrading on the de Rham algebra of $e'(P)$).

6.4. Proof of Proposition 6.8.

Lemma 6.10. *Under the hypothesis on η in Proposition 6.8 we have*

$$\int_{A_n} \eta = \sum_{\substack{x \in \Gamma_M \setminus \mathcal{L}_W \\ (x, x) = 2n}} \min_{\lambda \in \Lambda_W} |(\lambda, x)| \int_{a_x} \eta$$

Proof. We use Lemma 4.3. Write $\eta = d\omega$ for some 1-form ω which by the support condition on η is closed in F_n . Since c_{x+ku} and c_x are parallel hence homologous circles in F_x , we see $\int_{a_{x+ku}} \eta = \int_{c_{x+ku}} \omega = \int_{c_x} \omega = \int_{a_x} \eta$. \square

Since

$$\tilde{\psi}'_{0,1}(n) = \sum_{\substack{x \in \Gamma_M \setminus \mathcal{L}_W \\ (x,x)=2n}} \sum_{\gamma \in \Gamma_M} \gamma^* \tilde{\psi}'_{0,1}(x),$$

Proposition 6.8 will now follow from

Proposition 6.11. *Under the hypothesis on η in Proposition 6.8, we have for any positive length vector $x \in \mathcal{L}_W$*

$$\int_{e'(P)} \eta \wedge \sum_{\gamma \in \Gamma_M} \gamma^* \tilde{\psi}'_{0,1}(x) = \left(\min_{\lambda \in \Lambda_W} |(\lambda, x)| \right) \left(\int_{a_x} \eta \right) e^{-\pi(x,x)}.$$

Proof. By choosing appropriate coordinates we can assume that $x = \mu e_2$ with $\mu = \pm \sqrt{2n}$, so that the singularity of $\sum_{\gamma \in \Gamma_M} \gamma^* \tilde{\psi}'_{0,1}(x)$ in $e'(P)$ occurs at $s = 0$. We pick a tubular neighborhood $U_\varepsilon = (-\varepsilon, \varepsilon) \times T^2$ in $e'(P)$ around F_x . Then we have first

$$\int_{e'(P)} \eta \wedge \sum_{\gamma \in \Gamma_M} \gamma^* \tilde{\psi}'_{0,1}^P(x) = \lim_{\varepsilon \rightarrow 0} \int_{e'(P) - U_\varepsilon} \eta \wedge \sum_{\gamma \in \Gamma_M} \gamma^* \tilde{\psi}'_{0,1}^P(x).$$

Since $\eta \wedge \tilde{\psi}'_{0,1}(x) = d(\omega \wedge \tilde{\psi}'_{0,1}(x))$ outside U_ε and $\partial(e'(P) - U_\varepsilon) = -\partial U_\varepsilon$ we see by Stokes' theorem

$$\begin{aligned} (6.3) \quad \int_{e'(P) - U_\varepsilon} \eta \wedge \sum_{\gamma \in \Gamma_M} \gamma^* \tilde{\psi}'_{0,1}(x) &= - \int_{\partial U_\varepsilon} \omega \wedge \sum_{\gamma \in \Gamma_M} \gamma^* \tilde{\psi}'_{0,1}(x) \\ &= \sum_{\gamma \in \Gamma_M} \int_{T^2} \left[\omega(-\varepsilon, w) \wedge \tilde{\psi}'_{0,1}(\gamma^{-1}x, -\varepsilon, w) - \omega(\varepsilon, w) \wedge \tilde{\psi}'_{0,1}(\gamma^{-1}x, \varepsilon, w) \right]. \end{aligned}$$

For $\gamma \neq 1$ we note that $\omega(s, w) \wedge \tilde{\psi}'_{0,1}(\gamma^{-1}x, s, w)$ is continuous at $s = 0$, while for $\gamma = 1$, we have

$$(6.4) \quad \tilde{\psi}'_{0,1}(\mu e_2, s, w) = \frac{1}{2} |\mu| (\operatorname{sgn}(s) dw_2 - dw_3) e^{-\pi \mu^2}.$$

Hence taking the limit in the last term of (6.3) we obtain

$$|\mu| e^{-\pi \mu^2} \int_{T^2} \omega_3(0, w) dw_2 dw_3 = |\mu| e^{-\pi \mu^2} \int_{T^2/c_{e_2}} \left(\int_{c_{e_2}} \omega(0, w_2, w_3) \right) dw_2.$$

In the expression T^2/c_{e_2} (and for the rest of this proof) we have abused notation and identified the cycle c_{e_2} with the subgroup $0 \times S^1$ of T^2 .

Here ω_3 is the dw_3 component of ω and we used that ∂D_x is the w_3 -line in W . Note that the inner integral on the right is the period of ω over (homologous) horizontal translates of the cycle c_{e_2} . But the restriction of ω to F_x is closed so $\int_{c_{e_2}} \omega(0, w_2, w_3)$ is independent of w_2 and the last integral becomes $\left(\int_{T^2/c_{e_2}} dw_2 \right) \left(\int_{c_{e_2}} \omega \right) e^{-\pi \mu^2}$. But $\int_{c_{e_2}} \omega = \int_{A_{e_2}} \eta$. The proposition is then a consequence of

$$|\mu| \int_{T^2/\partial c_{e_2}} dw_2 = |\mu| \min_{\lambda \in \Lambda_W} |(\lambda, e_2)| = \min_{\lambda \in \Lambda_W} |(\lambda, x)|,$$

which follows from the fact that the map $W \rightarrow \mathbb{R}$ given by $w \mapsto (w, e_2)$ induces an isomorphism $T^2/\partial C_{e_2} \simeq \mathbb{R}/\min'_{\lambda \in \Lambda_W} |(\lambda, e_2)|\mathbb{Z}$. \square

6.5. Proof of Theorem 6.7. We now prove Theorem 6.7. First we will assume that we are in case(i). We need to show

$$\int_c \tilde{\psi}'_{0,1}(n) = \text{Lk}(\partial C_n, c) e^{-2\pi n} = (A_n \cdot c) e^{-2\pi n}.$$

The theorem will be a consequence of the following discussion. We may assume that c is an embedded loop in $e'(P)$ (note that since any loop in a manifold of dimension 3 or more is homotopic to an embedded loop by transversality any homology class of degree 1 in $e'(P)$ is represented by an embedded loop).

Choose a tubular neighborhood $N(c)$ of c such that $N(c)$ is disjoint from F_n . Let η_c be a closed 2-form which is supported inside $N(c)$ and has integral 1 on the disk fibers of $N(c)$ (a Thom class for the normal disk bundle $N(c)$). Then we have proved in Subsection 6.2

Lemma 6.12.

$$(6.5) \quad \int_{A_n} \eta_c = A_n \cdot c = \text{Lk}(\partial C_n, c).$$

We then have

Lemma 6.13.

$$\int_c \tilde{\psi}'_{0,1}(n) = \left(\int_{A_n} \eta_c \right) e^{-2\pi n}.$$

Proof. To prove the Lemma we compute $\int_{e'(P)} \eta_c \wedge \tilde{\psi}'_{0,1}(n) = \int_{e'(P)} \tilde{\psi}'_{0,1}(n) \wedge \eta_c$ in two different ways. First we apply Proposition 6.8 with $\eta = \eta_c$. We deduce

$$\int_{e'(P)} \eta_c \wedge \tilde{\psi}'_{0,1}(n) = \left(\int_{A_n} \eta_c \right) e^{-2\pi n}.$$

Next choose a tubular neighborhood V_n of the fibers F_n such that $e'(P) - V_n$ contains $N(c)$. Then $\tilde{\psi}'_{0,1}(n)$ is smooth on $e'(P) - V_n \supset \text{supp}(\eta_c)$. Also, since η_c is the extension of a Thom class by zero, the restriction of η_c to $e'(P) - V_n$ represents the Poincaré dual $PD(c)$ of the absolute cycle c in $e'(P) - V_n$. The lemma now follows from

$$\int_{e'(P)} \tilde{\psi}'_{0,1}(n) \wedge \eta_c = \int_{e'(P) - V_n} \tilde{\psi}'_{0,1}(n) \wedge \eta_c = \int_{e'(P) - V_n} \tilde{\psi}'_{0,1}(n) \wedge PD(c) = \int_c \tilde{\psi}'_{0,1}(n). \quad \square$$

By Lemma 6.12 this concludes the proof of Theorem 6.7 in the case when c is disjoint from the fibers F_n .

It remains to treat case (ii). Thus we now assume that $c = c_y$ which is contained in a fiber F_x containing a component of ∂C_n . We first prove

Lemma 6.14.

$$\int_c \tilde{\psi}'_{0,1}(x) = \int_{c(\epsilon)} \tilde{\psi}'_{0,1}(x).$$

Proof. We can take $x = \mu e_2$ and hence c is contained in the fiber over the image of $e_3 \in W$. Hence, by Proposition 4.1, c is the circle in the torus fiber at $s(x) = 0$ in the e_3 -direction, i.e., parallel to the image of $(0, \mathbb{R}e_3)$ in $e'(P)$. We note that by (6.4) even though $\tilde{\psi}'_{0,1}(x)$ is not defined on the whole fiber over $s = 0$ its restriction to c is smooth. Hence the left hand side is well-defined since all the other terms in the sum are defined on the whole fiber and in fact in a neighborhood of that fiber. Hence the locally constant form $\sum_{\gamma \in \Gamma_M} \gamma^* \tilde{\psi}'_{0,1}(x)$ is closed on the cylinder $[0, \epsilon] \times c$, and its integrals over the circles $s \times c$ all coincide. But $\epsilon \times c = c(\epsilon)$. The lemma follows. \square

Summing over x and using case (i) we obtain

$$\int_c \tilde{\psi}'_{0,1}(n) = \int_{c(\epsilon)} \tilde{\psi}'_{0,1}(n) = \text{Lk}(\partial C_n, c(\epsilon)),$$

since $c(\epsilon)$ is disjoint from all the components of F_n . Thus it suffices to prove

$$(6.6) \quad \text{Lk}(\partial C_n, c) = \text{Lk}(\partial C_n, c(\epsilon)).$$

To this end suppose that $c \subset F_x \subset F_n$ and c_1, \dots, c_k are the components of ∂C_n contained in F_x . Hence c and c_i , $1 \leq i \leq k$, are all parallel. Since the fibers containing all other components of ∂C_n are disjoint from c , (6.6) will follow from

$$\text{Lk}(c_i, c) = \text{Lk}(c_i, c(\epsilon)), 1 \leq i \leq k.$$

If $c_i = c$ then the previous equation is the definition of $Lk(c, c)$. Thus we may assume c_i is parallel to and disjoint from c . In this case their linking number is already topologically defined. But since c is disjoint from c_i the circles c and $c(\epsilon)$ are homologous in the complement of c_i (by the product homology $c \times [0, \epsilon]$) and since the linking number with c_i is a homological invariant of the complement of c_i in $e'(P)$ we have $\text{Lk}(c_i, c(\epsilon)) = Lk(c_i, c)$.

We this Theorem 6.7 is proved.

7. THE GENERATING SERIES OF THE CAPPED CYCLES

In this section, we show that the generating series of the ‘capped’ cycles C_n^c gives rise to a modular form, extending Theorem 7.1 to a lift of the full cohomology $H^2(X)$ of X . In particular, we give our new proof of the theorem of Hirzebruch and Zagier and show how a remarkable feature of their proof appears from our point of view.

7.1. The theta series associated to φ_2 . We define the theta series

$$\theta_{\varphi_2}(\tau, \mathcal{L}) = \sum_{x \in \mathcal{L}} \varphi_2(x, \tau, z).$$

In the following we will often drop the argument $\mathcal{L} = L + h$. For $n \in \mathbb{Q}$, we also set

$$\varphi_2(n) = \sum_{n \in \mathcal{L}_n, x \neq 0} \varphi_2(x).$$

Clearly, $\theta_{\varphi_2}(\tau, \mathcal{L})$ and $\varphi_2(n)$ descend to closed differential 2-forms on X . Furthermore, $\theta_{\varphi_2}(\tau, \mathcal{L})$ is a non-holomorphic modular form in τ of weight 2 for the principal congruence subgroup $\Gamma(N)$. In fact, for $\mathcal{L} = L$ as in Example 2.1, $\theta_{\varphi_2}(\tau, \mathcal{L})$ transforms like a form for $\Gamma_0(d)$ of nebentypus.

Theorem 7.1 (Kudla-Millson [24]). *We have*

$$[\theta_{\varphi_2}(\tau)] = -\frac{1}{2\pi}\delta_{h_0}[\omega] + \sum_{n>0} \text{PD}[C_n]q^n \in H^2(X, \mathbb{Q}) \otimes M_2(\Gamma(N)).$$

That is, for any closed 2-form η on X with compact support,

$$\Lambda(\eta, \tau) := \int_X \eta \wedge \theta_{\varphi_2}(\tau, \mathcal{L}) = -\frac{1}{2\pi}\delta_{h_0} \int_X \eta \wedge \omega + \sum_{n>0} \left(\int_{C_n} \eta \right) q^n.$$

Here δ_{h_0} is Kronecker delta, and ω is the Kähler form on D normalized such that its restriction to the base point is given by $\omega_{13} \wedge \omega_{14} + \omega_{23} \wedge \omega_{24}$. We obtain a map

$$(7.1) \quad \Lambda : H_c^2(X, \mathbb{C}) \rightarrow M_2(\Gamma(N))$$

from the cohomology with compact supports to the space of holomorphic modular forms of weight 2 for the principal congruence subgroup $\Gamma(N) \subset \text{SL}_2(\mathbb{Z})$. Alternatively, for C an absolute 2-cycle in X defining a class in $H_2(X, \mathbb{Z})$, the lift $\Lambda(C, \tau)$ is given by (1.1) with C_0 the class given by $-\frac{1}{2\pi}\delta_{h_0}[\omega]$.

The key fact for the proof of the Fourier expansion is that for $n > 0$, the form $\varphi_2(n)$ is a Poincaré dual form of C_n , while $\varphi_2(n)$ is exact for $n \leq 0$, see also Section 8.

7.2. The restrictions of the global theta functions.

Theorem 7.2. *The differential forms $\theta_{\varphi_2}(\mathcal{L}_V)$ and $\theta_{\psi_1}(\mathcal{L}_V)$ on X extend to the Borel-Serre compactification \overline{X} . More precisely, for the restriction i_P^* to the boundary face $e'(P)$ of \overline{X} , we have*

$$i_P^* \theta_{\varphi_2}(\mathcal{L}_V) = \theta_{\varphi_{1,1}}^P(\mathcal{L}_{W_P}) \quad \text{and} \quad i_P^* \theta_{\psi_1}(\mathcal{L}_V) = \theta_{\psi_{0,1}}^P(\mathcal{L}_{W_P}).$$

Proof. The restriction of $\theta_{\varphi_2}(\mathcal{L}_V)$ is the theme (in much greater generality) of [13]. For $\theta_{\psi_1}(\mathcal{L}_V)$ one proceeds in the same way. In short, one detects the boundary behaviour of the theta functions by switching to a mixed model of the Weil representation. For a model calculation see the proof of Theorem 7.4 below. \square

We conclude by Proposition 6.2

Theorem 7.3. *The restriction of $\theta_{\varphi_2}(\mathcal{L}_V)$ to the boundary of \overline{X} is exact and*

$$i_P^* \theta_{\varphi_2}(\mathcal{L}_V) = d \left(\theta_{\varphi_{0,1}}^P(\mathcal{L}_{W_P}) \right).$$

We also have a crucial restriction result for the singular form $\tilde{\psi}_{0,1}$. However, one needs to be careful in forming the naive theta series associated to $\tilde{\psi}_{0,1}$ by summing over all (non-zero) lattice elements. This would give a form on X with singularities on a dense subset of X . Instead we define $\tilde{\psi}_{2,0}(n)$ in the same way as for $\varphi_2(n)$ by summing over all non-zero $x \in \mathcal{L}_V$ of length $n \in \mathbb{Q}$. This gives a 1-form on X which for $n > 0$ has singularities along the locally finite cycle C_n . Similarly, we define

$$\tilde{\psi}_{0,1}^P(n) = \sum_{x \in \mathcal{L}_{W_P}, (x,x)=2n} \tilde{\psi}_{0,1}^P(x),$$

which descends to a 1-form on $e'(P)$ with singularities. We also define $\tilde{\psi}'_{0,1}(n)$ and $\phi_{0,1}^P(n)$ in the same way. We have

Proposition 7.4. *The restriction of the 1-form $\tilde{\psi}_1(n)$ to $e'(P)$ is given by*

$$i_P^* \tilde{\psi}_1(n) = \tilde{\psi}_{0,1}^P(n).$$

Proof. We assume that P is the stabilizer of the isotropic line $\ell = \mathbb{Q}u$. For $x = au + x_W + bu'$, we have for the majorant at $z = (w, t, s)$ the formula

$$(x, x)_z = \frac{1}{t^2} (a - (x_W, w) - bq(w))^2 + (x_w + bw, x_w + bw)_s + b^2 t^2.$$

Here $(\cdot, \cdot)_s$ is the majorant associated to W . Hence by (5.4) and (5.2) we see that the sum of all $x \in \mathcal{L}_V$ with $b \neq 0$ in $\tilde{\psi}_1(n)$ is uniformly rapidly decreasing as $t \rightarrow \infty$. Now fix an element $x_W \in \mathcal{L}_W$. Then $x_W + (a+h)u \in \mathcal{L}_V$ for all $a \in \mathbb{Z}$ for some $h \in \mathbb{Q}/\mathbb{Z}$; in fact all elements in $\mathcal{L}_V \cap u^\perp$ are of this form. We consider $\sum_{a \in \mathbb{Z}} \tilde{\psi}_1(x_W + (a+h)u, z)$ as $t \rightarrow \infty$. By considerations as in [13], sections 4 and 9, we can assume $w = 0$ and $s = 0$. We apply Poisson summation for the sum on $a \in \mathbb{Z}$ and obtain

$$\sum_{a \in \mathbb{Z}} \tilde{\psi}_1(x_W + au, z) = \sum_{k \in \mathbb{Z}} \left(\int_1^\infty P(x, t, r) e^{-2\pi x_3^2 r + t^2 k^2 / r} \frac{dr}{r} \right) e^{-2\pi i k h} e^{-\pi(x_W, x_W)},$$

where

$$P(x, t, r) = \frac{x_2 x_3 \sqrt{r}}{\sqrt{2}} dw_2 + \frac{1}{2\sqrt{2}} \left(\frac{1}{2\pi} - \frac{t^2 k^2}{r} \right) dw_3 - \frac{i x_3 k}{\sqrt{2}} dt + \frac{i x_2 k t}{\sqrt{2}} ds.$$

Now the sum over all $k \neq 0$ is rapidly decreasing while for $k = 0$ we obtain $\tilde{\psi}_{0,1}(x_W)$. If $x_W = 0$, i.e., for $n = 0$ one needs to argue slightly differently. Then we have

$$\sum_{a \neq 0} \tilde{\psi}_1(au, z) = \frac{1}{2\sqrt{2}\pi} \sum_{a \neq 0} e^{-\pi a^2 / t^2} \frac{dw_3}{t} = \frac{1}{2\sqrt{2}\pi} \left(\sum_{k \in \mathbb{Z}} e^{-\pi t^2 k^2} \right) dw_3 - \frac{1}{2\sqrt{2}\pi} \frac{dw_3}{t},$$

which goes to $\frac{1}{2\sqrt{2}\pi} dw_3 = \tilde{\psi}_{0,1}(0)$. This proves the proposition. \square

7.3. Main result. In the previous sections, we constructed a closed 2-form θ_{φ_2} on \overline{X} such that the restriction of θ_{φ_2} to the boundary $\partial \overline{X}$ was exact with primitive $\sum_{[P]} \theta_{\phi_{0,1}}^P$. From now on we usually write φ for φ_2 and ϕ for $\phi_{0,1}$ if it does not cause any confusion. By the definition of the differential for the mapping cone complex C^\bullet we immediately obtain by Theorem 7.2 and Theorem 7.3

Proposition 7.5. *The pair $(\theta_{\varphi_2}(\mathcal{L}_V), \sum_{[P]} \theta_{\phi_{0,1}}^P(\mathcal{L}_{W_P}))$ is a 2-cocycle in C^\bullet .*

We write for short $(\theta_\varphi, \theta_\phi)$. We obtain a class $[[\theta_\varphi, \theta_\phi]]$ in $H^2(C^\bullet)$ and hence a class $[\theta_\varphi, \theta_\phi]$ in $H_c^2(X)$. The pairing with $[\theta_\varphi, \theta_\phi]$ then defines a lift Λ^c on differential 2-forms on \overline{X} , which factors through $H^2(\overline{X}) = H^2(X)$. By Lemma 3.5 it is given by

$$\Lambda^c(\eta, \tau) = \int_{\overline{X}} \eta \wedge \theta_{\varphi_2} - \sum_{[P]} \int_{e'(P)} i^* \eta \wedge \theta_{\phi_{0,1}}^P.$$

Theorem 7.6. *The class $[[\theta_\varphi, \theta_\phi]]$ is holomorphic, that is,*

$$L(\theta_\varphi, \theta_\phi) = d(\theta_{\psi_1}, 0).$$

Hence $[\theta_\varphi, \theta_\phi]$ is a holomorphic modular form with values in the compactly supported cohomology of X , so that the lift Λ^c takes values in the holomorphic modular forms.

Proof. By Theorem 7.2 and Theorem 6.2 we calculate

$$d(\theta_{\psi_1}, 0) = (d\theta_{\psi_1}, i^*\theta_{\psi_1}) = \left(L\theta_{\varphi_2}, \sum_{[P]} \theta_{\psi_{0,1}}^P \right) = L \left(\theta_{\varphi_2}, \sum_{[P]} \theta_{\phi_{0,1}}^P \right). \quad \square$$

It remains to compute the Fourier expansion in τ of $[\theta_\varphi, \theta_\phi](\tau)$. We will carry this out in Section 8.

Theorem 7.7. *We have*

$$[\theta_\varphi, \theta_\phi](\tau) = -\frac{1}{2\pi} \delta_{h_0}[\omega] + \sum_{n>0} \text{PD}[C_n^c] q^n \in H_c^2(X, \mathbb{Q}) \otimes M_2(\Gamma(N)).$$

That is, for any closed 2-form η on \bar{X}

$$\Lambda^c(\eta, \tau) = -\frac{1}{2\pi} \delta_{h_0} \int_X \eta \wedge \omega + \sum_{n>0} \left(\int_{C_n^c} \eta \right) q^n,$$

In particular, the map takes values in the holomorphic modular forms and factors through cohomology. We obtain a map

$$(7.2) \quad \Lambda^c : H^2(X) \rightarrow M_2(\Gamma(N))$$

from the cohomology with compact supports to the space of holomorphic modular forms of weight 2 for the principal congruence subgroup $\Gamma(N) \subseteq \text{SL}_2(\mathbb{Z})$. Alternatively, for C any relative 2-cycle in X defining a class in $H_2(\bar{X}, \partial\bar{X}, \mathbb{Z})$, we have

$$\Lambda^c(C, \tau) = -\frac{1}{2\pi} \delta_{h_0} \text{vol}(C) + \sum_{n>0} (C_n^c \cdot C) q^n \in M_2(\Gamma(N)).$$

Remark 7.8. In the theorem we now consider the Kähler form ω representing a class in the compactly supported cohomology. In fact, our mapping cone construction gives an explicit coboundary by which ω is modified to become rapidly decreasing.

7.4. The Hirzebruch-Zagier Theorem. We now view $[\theta_\varphi, \theta_\phi]$ as a class in $H^2(\tilde{X})$ via the map $j_\# : H_c^2(X) \rightarrow H^2(\tilde{X})$. We recover the Hirzebruch-Zagier-Theorem.

Theorem 7.9. *We have*

$$j_\#[\theta_\varphi, \theta_\phi](\tau) = -\frac{1}{2\pi} \delta_{h_0}[\omega] + \sum_{n>0} [T_n^c] q^n \in H^2(\tilde{X}, \mathbb{Q}) \otimes M_2(\Gamma(N)).$$

In particular,

$$-\frac{1}{2\pi} \delta_{h_0} \text{vol}(T_m) + \sum_{n>0} (T_n^c \cdot T_m)_{\tilde{X}} q^n \in M_2(\Gamma(N)).$$

This is the result Hirzebruch-Zagier proved for certain Hilbert modular surfaces (Example 2.1) by explicitly computing the intersection numbers $T_m \cdot T_n^c$.

Proof. This follows from Theorem 7.7 since $j_*C_n^c = T_n^c$ (Proposition 4.7), combined with the following general principle. Suppose ω is a compactly supported form on X such that the cohomology class of ω is the Poincaré dual of the homology class of a cycle C : $[\omega] = \text{PD}(C)$. Then we have $j_\#[\omega] = \text{PD}(j_*C)$. To see this we have only to replace ω by a cohomologous ‘Thom representative’ of $\text{PD}(C)$, namely a closed

form $\tilde{\omega}$ supported in a tubular neighborhood $N(C)$ of C in X such that the integral of $\tilde{\omega}$ over any disk of $N(C)$ is one. Then it is a general fact from algebraic topology (extension by zero of a Thom class) that $\tilde{\omega}$ represents the Poincaré dual of C in any manifold M containing $N(C)$, in particular for $M = \tilde{X}$. \square

Remark 7.10. If one is only interested in recovering the statement of this theorem, then there is also a different way of deriving this from the Kudla-Millson theory. Namely, the lift Λ on $H_2(X)$ (Theorem 7.1) factors through the quotient of $H_2(X)$ by $H_2(\partial X)$ since the restriction of θ_{φ_2} is exact (Theorem 7.3). But by Proposition 3.3 we have $j_*H_2(X) \simeq H_2(X)/H_2(\partial X)$, and the Hirzebruch-Zagier result exactly stipulates the modularity of the lift of classes in $j_*H_2(X)$. However, in that way one misses the remarkable extra structure coming from ∂X as we will explain in the next subsection.

7.5. The lift of special cycles. We now consider the lift of a special cycle C_y . By Theorem 7.7 and Lemma 3.5 we see

$$(7.3) \quad \begin{aligned} \Lambda^c(C_y, \tau, \mathcal{L}_V) &= -\frac{1}{2\pi} \delta_{h0} \text{vol}(C_y) + \sum_{n>0} (C_n^c \cdot C_y) q^n \\ &= \int_{C_y} \theta_{\varphi_2}(\tau, \mathcal{L}_V) - \sum_{[P]} \int_{(\partial C_y)_P} \theta_{\phi_{0,1}}^P(\tau, \mathcal{L}_{W_P}). \end{aligned}$$

The two terms on the right, the integrals over C_y and ∂C_y , are both non-holomorphic modular forms (see below) whose difference is holomorphic (by Theorem 7.6). So the generating series series of $(C_n^c \cdot C_y)$ is the sum of two non-holomorphic modular forms. We now give geometric interpretations for the two individual non-holomorphic forms.

Following [16] we define the *interior* intersection number of two special cycles by

$$(C_n \cdot C_y)_X = (C_n \cdot C_y)^{tr} + \text{vol}(C_n \cap C_y),$$

the sum of the transversal intersections and the volume of the 1-dimensional (complex) intersection of C_n and C_y which occur if one of the components of C_n is equal to C_y .

Theorem 7.11. *We have*

$$\int_{C_y} \theta_{\varphi_2}(\tau, \mathcal{L}_V) = -\frac{1}{2\pi} \delta_{h0} \text{vol}(C_y) + \sum_{n=1}^{\infty} (C_n \cdot C_y)_X q^n + \sum_{n \in \mathbb{Q}} \sum_{[P]} \int_{(\partial C_y)_P} \tilde{\psi}_{0,1}^P(n)(\tau).$$

So the Fourier coefficients of the holomorphic part of the non-holomorphic modular form $\int_{C_y} \theta_{\varphi_2}$ are the interior intersection numbers of the cycles C_y and C_n .

Proof. This is essentially [9], section 5, where more generally $O(p, 2)$ is considered. There the interpretation of the holomorphic Fourier coefficients as interior intersection number is given. (For more details of an analogous calculation see [14], section 8). A little calculation using the formulas in [9] gives the non-holomorphic contribution. A more conceptual proof would use the relationship between φ_2 and $\tilde{\psi}_1$ (see Proposition 5.2 and Section 8) and the restriction formula for $\tilde{\psi}_1(n)$ (Theorem 7.4). \square

By slight abuse of notation we write $\text{Lk}(C_n, C_y) = \sum_{[P]} \text{Lk}((\partial C_n)_P, (\partial C_y)_P)$ for the total linking number of ∂C_n and ∂C_y . Then by Theorem 6.3 we obtain

Theorem 7.12.

$$\sum_{[P]} \int_{(\partial C_y)_P} \theta_{\phi,1}^P(\tau, \mathcal{L}_{W_P}) = \sum_{n>0} \text{Lk}(C_n, C_y) q^n + \sum_{n \in \mathbb{Q}} \sum_{[P]} \int_{(\partial C_y)_P} \tilde{\psi}_{0,1}^P(n)(\tau).$$

So the Fourier coefficients of the holomorphic part of $\int_{(\partial C_y)_P} \theta_{\phi}^P(\tau, \mathcal{L}_{W_P})$ are the linking numbers of the cycles ∂C_y and ∂C_n at the boundary component $e'(P)$.

Remark 7.13. There is also another ‘‘global’’ proof for Theorem 7.12. The cycle C_y intersects $e'(P)$ transversally (when pushed inside) and hence also the cap A_n . From this it is not hard to see that we can split the intersection number $C_n^c \cdot C_y$ as

$$C_n^c \cdot C_y = (C_n \cdot C_y)_X - \text{Lk}(C_n, C_y).$$

Hence Theorem 7.12 also follows from combining (7.3) and Theorem 7.11.

Hirzebruch-Zagier also obtain the modularity of the functions given in Theorems 7.11 and 7.12, but by quite different methods. In particular, they explicitly calculate the intersection number $T_n^c \cdot T_m$. They split the intersection number into the interior part $(T_n \cdot T_m)_X$ and a ‘boundary contribution’ $(T_n \cdot T_m)_{\infty}$ given by

$$(T_n \cdot T_m)_{\infty} = (T_n \cdot T_m)_{\bar{X}-X} - (T_m - T_m^c) \cdot (T_n - T_n^c).$$

Now by Theorem 7.9 and its proof we have

$$T_n^c \cdot T_m = C_n^c \cdot C_m.$$

We have (per definition) $(T_n \cdot T_m)_X = (C_n \cdot C_m)_X$, so Theorem 7.11 gives the generating series for $(T_n \cdot T_m)_X$. Note that Theorem 5.4 in [9] also compares the explicit formulas in [16] for $(T_n \cdot T_m)_X$ with the ones obtained via $\int_{C_y} \theta_{\varphi_2}(\tau, \mathcal{L}_V)$. All this implies

$$(T_n \cdot T_m)_{\infty} = \text{Lk}(C_n \cdot C_m).$$

Independently, we also obtain this from comparing the explicit formulas for the boundary contribution in [16], Section 1.4 with our formulas for the linking numbers, Theorem 4.11 and Example 4.12.

8. A CURRENT APPROACH FOR THE SPECIAL CYCLES

In this section we prove Theorem 7.7, the crucial Fourier coefficient formula for our lift Λ^c . As a consequence of our approach we will also obtain Theorem 6.7, the linking number interpretation for the lift at the boundary.

8.1. A differential character for C_n^c . The key step for the entire Kudla-Millson theory is that for $n > 0$ the form $\varphi_2(n)$ is a Poincaré dual form for the cycle C_n , i.e.,

Theorem 8.1 ([22, 23]). *Let η be a closed rapidly decreasing 2-form. Then*

$$\int_X \eta \wedge \varphi_2(n) = \left(\int_{C_n} \eta \right) e^{-2\pi n}.$$

To show this they employ at some point a homotopy argument which requires η to be rapidly decaying. Since we require η to be any closed 2-form on the compactification \overline{X} , their approach is not applicable in our case. Instead, we use a differential character argument for φ_2 which implicitly already occurred in [5], Section 7 for general signature (p, q) . Namely, we have

Theorem 8.2. ([5], Section 7) *Let $n > 0$. The singular form $\tilde{\psi}_1(n)$ is a differential character in the sense of Cheeger-Simons for the cycle C_n . More precisely, $\tilde{\psi}_1(n)$ is a locally integrable 1-form on X , and for any compactly supported 2-form η we have*

$$\int_X \eta \wedge \varphi_2(n) = \left(\int_{C_n} \eta \right) e^{-2\pi n} - \int_X d\eta \wedge \tilde{\psi}_1(n).$$

Proof. This is the content of the proofs of Theorem 7.1 and Theorem 7.2 in [5]. There the analogous properties for a singular theta lift associated to ψ is established. However, the proofs boil down to establish the claims for $\tilde{\psi}_1$. The form $\tilde{\psi}$ there is indeed the form $\tilde{\psi}_1$ of this paper. \square

Remark 8.3. The form $\tilde{\psi}_1$ is closely related to Kudla's Green function ξ [18, 19] (more generally for $O(p, 2)$) which is given by

$$\xi(x) = \left(\int_1^\infty \varphi_0^0(\sqrt{rx}) \frac{dr}{r} \right) e^{-\pi(x,x)}.$$

Then $\Xi(n) = \sum_{x \in \mathcal{L}_n} \xi(x)$ gives rise to a Green's function for the divisor C_n and moreover $dd^c \xi = \varphi_2$. Here $d^c = \frac{1}{4\pi i}(\partial - \bar{\partial})$. This suggests $d^c \xi = \tilde{\psi}_1$, which indeed follows from $d^c \varphi_0 = -\psi_1$, see [5], Remark 4.5.

For $n \in \mathbb{Q}$ we define

$$\varphi_2^c(n) := \varphi_2(n) - \sum_{[P]} d(f\pi^*\phi_{0,1}^P(n))$$

and follow the current approach to show that for $n > 0$ the form $\varphi_2^c(n)$ is a Poincaré dual form for the cycle C_n^c . Here we follow the notation of subsection 3.3. That is, $\pi^*\phi_{0,1}^P(n)$ is the pullback to a product neighborhood V of $\partial\overline{X}$, and f is a smooth function on V of the geodesic flow coordinate t which is 1 near $t = \infty$ and zero else. Note that $\varphi_2^c(n)$ is exactly the n -th Fourier coefficient of the mapping cone element $[\theta_\varphi, \theta_\phi]$, when realized as a rapidly decreasing form on X . We also define

$$\tilde{\psi}_1^c(n) = \tilde{\psi}_1(n) - f\pi^*\phi_{0,1}^P(n).$$

We call a differential form η on \overline{X} special if in a neighborhood of each boundary component $e'(P)$ it is the pullback of a form η_P on $e'(P)$ under the geodesic retraction and if the pullback of the form η_P to the universal cover $e(P)$ is N -left-invariant. The significance of the forms lies in the fact that the complex of special forms also computes the cohomology of \overline{X} . Note that the proof of Theorem 7.2 shows that θ_{φ_2} is 'almost' special; it only differs from a special form by a rapidly decreasing form.

Theorem 8.4. *Let $n > 0$. The form $\tilde{\psi}_1^c(n)$ is a differential character for the cycle C_n^c . More precisely, $\tilde{\psi}_1^c(n)$ is a locally integrable 1-form on X and satisfies the following current equation on special 2-forms on \bar{X} :*

$$d[\tilde{\psi}_1^c(n)] + \delta_{C_n} e^{-2\pi n} = [\varphi_2^c(n)].$$

That is, for any special 2-form η on \bar{X} we have

$$\int_X \eta \wedge \varphi_2^c(n) = \left(\int_{C_n^c} \eta \right) e^{-2\pi n} - \int_X d\eta \wedge \tilde{\psi}_2^c(n).$$

This implies Theorem 7.7 for the positive Fourier coefficients. For $n \leq 0$, the form $\varphi_2^c(n)$ is exact with primitive $\tilde{\psi}_2^c(n)$ which by Theorem 7.4 is decaying. So Theorem 8.4 holds also for $n \leq 0$ with $C_n^c = \emptyset$. Hence for these coefficients only the term $x = 0$ contributes, which gives the integral of η against the Kähler form.

Remark 8.5. In view of Remark 8.3 it is very natural question to ask how one can modify Kudla's Green's function $\Xi(n)$ to obtain a Green's function for the cycle T_n^c in \tilde{X} . Extensive discussions with Kühn suggest that (if X has only one cusp)

$$\Xi(n) - t \sum_{\substack{x \in \mathcal{L}_W \\ (x,x)=2n}} f\pi^*(B(x) + B'(x))$$

is such a Green's function, but we have not checked all details.

8.2. Proof of Theorem 8.4. For simplicity assume that X has only one cusp and continue to drop the superscript P . We let ρ_T be a family of smooth functions on a standard fundamental domain \mathcal{F} of Γ in D only depending on t which is 1 for $t \leq T$ and 0 for $T + 1$. We then have

$$\int_X \eta \wedge \varphi_2^c(n) = \lim_{T \rightarrow \infty} \int_X \rho_T \eta \wedge (\varphi_2(n) - d(f\pi^*\phi_{0,1}(n))).$$

We apply Theorem 8.2 for the compactly supported form $\rho_T \eta$ and obtain

$$(8.1) \quad \int_X \eta \wedge \varphi_2^c(n) = \lim_{T \rightarrow \infty} \left[\left(\int_{C_n} \rho_T \eta \right) e^{-2\pi n} - \int_X d(\rho_T \eta) \wedge \tilde{\psi}_1(n) \right. \\ \left. - \int_X d(\rho_T \eta \wedge (f\pi^*\phi_{0,1}(n))) - d(\rho_T \eta) \wedge f\pi^*\phi_{0,1}(n) \right]$$

The first term on the right hand side of (8.1) goes to $\left(\int_{C_n} \eta \right) e^{-2\pi n}$ as $T \rightarrow \infty$, while the third vanishes for any T by Stokes' theorem. For the two remaining terms of (8.1) we first note $d(\rho_T \eta) = \rho'_T(t) dt \wedge \eta + \rho_T d\eta$ and $\rho'_T(t) = 0$ outside $[T, T + 1]$. We obtain for these two terms

$$(8.2) \quad - \int_X (d\eta) \wedge \left(\tilde{\psi}_1(n) - f\pi^*\phi_{0,1}(n) \right) \\ - \lim_{T \rightarrow \infty} \int_T^{T+1} \int_{e'(P)} \rho'_T(t) dt \wedge \eta \wedge \left(\tilde{\psi}_1(n) - f\pi^*\phi_{0,1}(n) \right).$$

It remains to compute the second term in the previous equation. For T sufficiently large we have $f \equiv 1$. Furthermore by Theorem 7.4 and its proof we have $\tilde{\psi}_1(n) = \pi^* \tilde{\psi}_{0,1}(n) + O(e^{-Ct})$. As $\phi_{0,1}(n) = \tilde{\psi}_{0,1}(n) + \tilde{\psi}'_{0,1}(n)$, we can replace $\tilde{\psi}_1(n) - f\pi^* \phi_{0,1}(n)$ by $-\pi^* \tilde{\psi}'_{0,1}(n)$. Since η is special it does not depend on the t -variable near the boundary. For the last term in (8.2)

$$\lim_{T \rightarrow \infty} \int_T^{T+1} \rho'_T(t) dt \int_{e'(P)} \eta \wedge \pi^* \tilde{\psi}'_{0,1}(n) = - \int_{e'(P)} \eta \wedge \tilde{\psi}'_{0,1}(n) = - \left(\int_{A_n} \eta \right) e^{-2\pi n}.$$

Indeed, for $\eta = \Omega$ this is Remark 6.9. Otherwise, η is exact with special primitive ω , and it is not hard to see that the proof of Proposition 6.8 carries over to this situation. Since $C_n^c = C_n \coprod (-A_n)$ collecting all terms completes the proof of Theorem 8.4.

REFERENCES

- [1] N. Bergeron, J. Millson, and C. Moeglin, *Hodge type theorems for arithmetic manifolds associated to orthogonal groups*, preprint.
- [2] A. Borel and L. Ji, *Compactifications of symmetric and locally symmetric spaces*, Birkhäuser, 2006.
- [3] A. Borel and J.-P. Serre, *Corners and arithmetic groups*, Commentarii Mathematici Helvetici **48** (1973), 436-491.
- [4] J. Bruinier, *Hilbert modular forms and their applications*, in: The 1-2-3 of Modular Forms, Springer-Verlag (2008).
- [5] J. Bruinier and J. Funke, *On two geometric theta lifts*, Duke Math J. **125** (2004), 45-90.
- [6] J. Cogdell. *Arithmetic cycles on Picard modular surfaces and modular forms of Nebentypus*, J. Reine u. Angew. Math. **357** (1985), 115-137.
- [7] D. DeTurck and H. Gluck, *Electrodynamics and the Gauss linking integral on the 3-sphere and in hyperbolic 3-space*, J. Math. Phys. **49**, (2008)
- [8] H. Flanders, *Differential forms with applications to the physical sciences*, Mathematics in Science and Engineering **11** (1963), Academic Press.
- [9] J. Funke, *Heegner divisors and nonholomorphic modular forms*, Compositio Math. **133** (2002), 289-321.
- [10] J. Funke, *Singular theta liftings for unitary groups and the construction of Green currents for special cycles*, in preparation.
- [11] J. Funke and J. Millson, *Cycles in hyperbolic manifolds of non-compact type and Fourier coefficients of Siegel modular forms*, Manuscripta Math. **107** (2002), 409-449.
- [12] J. Funke and J. Millson, *Cycles with local coefficients for orthogonal groups and vector-valued Siegel modular forms*, American J. Math. **128**, 899-948 (2006)
- [13] J. Funke and J. Millson, *Boundary behavior of special cohomology classes arising from the Weil representation*, preprint.
- [14] J. Funke and J. Millson, *Spectacle cycles with coefficients and modular forms of half-integral weight*, to appear in: Arithmetic Geometry and Automorphic forms, Volume in honor of the 60th birthday of Stephen S. Kudla, Advanced Lectures in Mathematics series. International Press and the Higher Education Press of China (2011).
- [15] J. Funke and J. Millson, in preparation.
- [16] F. Hirzebruch and D. Zagier, *Intersection numbers of curves on Hilbert modular surfaces and modular forms of Nebentypus*, Inv. Math. **36** (1976), 57-113.
- [17] W. Hoffman and H. He, *Picard groups of Siegel modular threefolds and theta lifting*, preprint.
- [18] S. Kudla, *Central derivatives of Eisenstein series and height pairings*, Ann. of Math. **146** (1997), 545-646.
- [19] S. Kudla, *Integrals of Borcherds forms*, Compositio Math. **137** (2003), 293-349.

- [20] S. Kudla, *Special cycles and derivatives of Eisenstein series*, in: Heegner points and Rankin L -series, 243-270, Math. Sci. Res. Inst. Publ., **49**, Cambridge Univ. Press, 2004.
- [21] S. Kudla and J. Millson, *The theta correspondence and harmonic forms I*, Math. Ann. **274** (1986), 353-378.
- [22] S. Kudla and J. Millson, *The Theta Correspondence and Harmonic Forms II*, Math. Ann. **277** (1987), 267-314.
- [23] S. Kudla and J. Millson, *Tubes, cohomology with growth conditions and application to the theta correspondence*, Canad. J. Math. **40** (1988), 1-37.
- [24] S. Kudla and J. Millson, *Intersection numbers of cycles on locally symmetric spaces and Fourier coefficients of holomorphic modular forms in several complex variables*, IHES Pub. **71** (1990), 121-172.
- [25] J. Milnor, *Singularities of Complex Hypersurfaces*, Annals of Math. Studies **61**, Princeton University Press, 1968.
- [26] T. Oda, *On modular forms associated with indefinite quadratic forms of signature $(2, n - 2)$* , Math. Annalen **231** (1977), 97-144.
- [27] T. Shintani, *On the construction of holomorphic cusp forms of half integral weight*, Nagoya Math. J. **58** (1975), 83-126.
- [28] G. van der Geer, *Hilbert modular surfaces*, Ergebnisse der Math. und ihrer Grenzgebiete (3), vol. **16**, Springer, 1988.
- [29] C. A. Weibel, *An introduction to homological algebra*, Cambridge studies in advanced mathematics, vol. **38**, Cambridge University Press, 1994.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF DURHAM, SCIENCE LABORATORIES, SOUTH RD, DURHAM DH1 3LE, UNITED KINGDOM

E-mail address: jens.funke@durham.ac.uk

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND, COLLEGE PARK, MD 20742, USA

E-mail address: jjm@math.umd.edu