

From Little Strings to M5-branes via a Quasi-Topological Sigma Model on Loop Group

Meng-Chwan Tan

Durham, 14 Aug 2018

LMS Higher Structures in M-Theory



Outline of Talk

- Introduction and Motivation
- Summary of Results
- Main Body of the Talk
- Conclusion and Future Directions

Introduction and Motivation

In this talk, we will discuss a **quasi-topological twist** of a 2d $\mathcal{N} = (2, 2)$ nonlinear sigma model (NLSM) on $\mathbb{C}P^1$ with target space the based loop group $\Omega SU(k)$.

The motivations for doing so are to:

- Describe the ground and half-excited states of the 6d A_{k-1} $\mathcal{N} = (2, 0)$ **little string theory**.
- Obtain a physical **derivation and generalization** of a mathematical relation by Braverman-Finkelberg which defines a **geometric Langlands correspondence for surfaces**.
- Elucidate the **1/2 and 1/4 BPS sectors** of the **M5-brane worldvolume theory**.

Introduction and Motivation

This talk is based on

- M.-C. Tan et al., *Little Strings, Quasi-Topological Sigma Model on Loop Group, and Toroidal Lie Algebras*, Nucl.Phys. B928, 469-498 (2018)

Built on earlier insights in

- M.-C. Tan, *Two-Dimensional Twisted Sigma Models And The Theory of Chiral Differential Operators*, Adv.Theor.Math.Phys. 10, 759-851 (2006).
- M.-C. Tan, *Five-Branes in M-Theory and a Two-Dimensional Geometric Langlands Duality*, Adv.Theor.Math.Phys. 14, 179-224 (2010).
- R. Dijkgraaf, *The Mathematics of Fivebranes*, Documenta Mathematica, 133-142 (1998).

Summary of Results

1. In the quasi-topological sigma model with target $\Omega SU(k)$, there is a scalar supercharge \overline{Q}_+ which generated supersymmetry survives on a $\mathbb{C}P^1$ worldsheet, whereby in the \overline{Q}_+ -cohomology, we have the following currents that generate the following toroidal algebra $\mathfrak{su}(k)_{\text{tor}}$:

$$[J_{m_1}^{an_1}, J_{m_2}^{bn_2}] = if_c^{ab} J_{m_1+m_2}^{c\{n_1+n_2\}} + c_1 n_1 \delta^{ab} \delta^{\{n_1+n_2\}0} \delta_{\{m_1+m_2\}0} + c_2 m_1 \delta^{ab} \delta^{\{n_1+n_2\}0} \delta_{\{m_1+m_2\}0}$$

2. In the topological subsector of the sigma model, we have instead the following affine algebra $\mathfrak{su}(k)_{\text{aff}}$:

$$[J_0^{an_1}, J_0^{bn_2}] = if_c^{ab} J_0^{c\{n_1+n_2\}} + c_1 n_1 \delta^{ab} \delta^{\{n_1+n_2\}0}$$

Summary of Results

3. Via a theorem by Atiyah in [1], the left-excited states (in the DLCQ) of the 6d $A_{k-1} \mathcal{N} = (2, 0)$ little string theory (LST) on $\mathbb{R}^{5,1}$ can be related to the \overline{Q}_+ -cohomology of the quasi-topological sigma model. In turn, we find that

left-excited spectrum of 6d $A_{k-1} (2, 0)$ LST = modules of $\mathfrak{su}(k)_{\text{tor}}$

4. Likewise, the ground states (in the DLCQ) of the 6d $A_{k-1} \mathcal{N} = (2, 0)$ LST on $\mathbb{R}^{5,1}$ can be related to the topological subsector of the sigma model. In turn, we find that

ground spectrum of 6d $A_{k-1} (2, 0)$ LST = modules of $\mathfrak{su}(k)_{\text{aff}}$

Summary of Results

5. This means (via the ground states) that we have

$$\mathbb{H}^*(\mathcal{M}_{SU(k)}^N(\mathbb{R}^4)) = \widehat{su}(k)_{c_1}^N$$

i.e., the intersection cohomology of the moduli space $\mathcal{M}_{SU(k)}^N(\mathbb{R}^4)$ of $SU(k)$ -instantons forms a finite submodule over $\mathfrak{su}(k)_{\text{aff}}$. This is just the Braverman-Finkelberg relation in [2]

6. This also means (via the left-excited states) that we have

$$H_{\check{C}ech}^*(\widehat{\Omega}_{\mathcal{M}_{SU(k)}^N(\mathbb{R}^4)}^{ch}) = \widehat{su}(k)_{c_1, c_2}^N$$

i.e., the Čech-cohomology of the sheaf $\widehat{\Omega}_{\mathcal{M}_{SU(k)}^N(\mathbb{R}^4)}^{ch}$ of Chiral de Rham Complex on $\mathcal{M}_{SU(k)}^N(\mathbb{R}^4)$ forms a submodule over $\mathfrak{su}(k)_{\text{tor}}$. This is a novel, physically-derived generalization of the Braverman-Finkelberg relation.

Summary of Results

7. Using the relevant SUSY algebras, one can show the correspondence between the ground states of the little string and the 1/2 BPS sector of the M5-brane worldvolume theory, from which we can compute the 1/2 BPS sector partition function to be

$$Z_{1/2} = \sum_{\widehat{\lambda}'} \chi_{\widehat{su}(k)_{c_1}}^{\widehat{\lambda}'}(p)$$

It is a cousin of a modular form which transforms as a representation of $SL(2, \mathbb{Z})$.

There is an intrinsic $SL(2, \mathbb{Z})$ symmetry in the M5-brane worldvolume theory on $\mathbb{R}^{5,1}$!

Emerges as gauge-theoretic S-duality of 4d $\mathcal{N} = 4$ SYM after compactifying on T^2 .

Summary of Results

8. Likewise, one can show the correspondence between the left-excited states of the little string and the 1/4 BPS sector of the M5-brane worldvolume theory, from which we can compute the 1/4 BPS sector partition function to be

$$Z_{1/4} = q^{\frac{1}{24}} \sum_{\widehat{\lambda}} \chi_{\widehat{su}(k)_{c_1}}^{\widehat{\lambda}}(p) \frac{1}{\eta(\tau)}$$

It is a cousin of an automorphic form which transforms as a representation of $SO(2, 2; \mathbb{Z})$.

There is an intrinsic $SO(2, 2; \mathbb{Z})$ symmetry of the M5-brane worldvolume theory on $\mathbb{R}^{5,1}$!

Emerges as string-theoretic T-duality of little strings after compactifying on T^2 .

LET'S EXPLAIN HOW WE GOT THESE RESULTS

Based Loop Group ΩG

A **loop group** LG is the group consisting of maps from the unit circle S^1 to a (Lie) group G :

$$f : S^1 \rightarrow G. \quad (1)$$

Parametrize S^1 by $e^{i\theta}$. If we impose the based point condition

$$f(\theta = 0) \rightarrow I, \quad (2)$$

we get the **based** loop group ΩG . One can show that

$$\Omega G = LG/G, \quad (3)$$

i.e., it is a G -equivariant subset of LG endowed with an LG -action.

ΩG also admits a closed nondegenerate symplectic two-form ω . The complex and symplectic structures of ΩG are compatible, and conspire to make it an infinite-dimensional Kähler manifold.

Based Loop Group ΩG

Let ξ and η be elements of $\Omega\mathfrak{g}$, the based loop algebra. Then, expanding them in the $L\mathfrak{g}$ basis gives

$$\begin{aligned}\xi(\theta) &= \xi_n e^{in\theta} = \xi_{an} T^a e^{in\theta}, \\ \eta(\theta) &= \eta_n e^{in\theta} = \eta_{an} T^a e^{in\theta},\end{aligned}\tag{4}$$

where $n \in \mathbb{Z}$ and $a = 1, \dots, \dim \mathfrak{g}$. The based point condition (2), which can be written as $e^{i\xi(\theta=0)} = 1$, then translates to $\sum_n \xi_{an} T^a = 0$.

The metric of ΩG is

$$g_{am, bn} = |n| \delta_{n+m, 0} \operatorname{Tr}(T_a T_b).\tag{5}$$

If we denote $T^a e^{im\theta} \equiv T^{am}$, we have

$$[T^{am}, T^{bn}] = if_c^{ab} T^{c(m+n)}.\tag{6}$$

The $\mathcal{N} = (2, 2)$ Sigma Model on $\mathbb{C}P^1$ with Target $\Omega SU(k)$

The action of the $\mathcal{N} = (2, 2)$ supersymmetric sigma model on $\mathbb{C}P^1$ with $\Omega SU(k)$ target space is

$$S = \int d^2z \left(g_{am, b\bar{n}} \left(\frac{1}{2} \partial_z \phi^{am} \partial_z \bar{\phi}^{b\bar{n}} + \frac{1}{2} \partial_z \phi^{am} \partial_z \bar{\phi}^{b\bar{n}} + \bar{\psi}_-^{b\bar{n}} D_z \psi_-^{am} + \psi_+^{am} D_z \bar{\psi}_+^{b\bar{n}} \right) - R_{am, c\bar{p}, bn, d\bar{q}} \psi_+^{am} \psi_-^{bn} \bar{\psi}_-^{c\bar{p}} \bar{\psi}_+^{d\bar{q}} \right), \quad (7)$$

where

$$\begin{aligned} \phi^{a(-n)} &= \bar{\phi}^{an}, \\ \psi_{\mp}^{a(-n)} &= \bar{\psi}_{\pm}^{an}. \end{aligned} \quad (8)$$

and

$$\begin{aligned} D_z \psi_-^{am} &= \partial_z \psi_-^{am} + \Gamma_{bn, cp}^{am} \partial_z \phi^{bn} \psi_-^{cp}, \\ D_z \bar{\psi}_+^{a\bar{m}} &= \partial_z \bar{\psi}_+^{a\bar{m}} + \Gamma_{b\bar{n}, c\bar{p}}^{a\bar{m}} \partial_z \bar{\phi}^{b\bar{n}} \bar{\psi}_+^{c\bar{p}}. \end{aligned} \quad (9)$$

Quasi-Topological A-Model on $\mathbb{C}P^1$ with Target $\Omega SU(k)$

We may twist the $\mathcal{N} = (2, 2)$ sigma model, i.e., shift the spin of the fields by their $U(1)_R$ -charges. Let us consider the A-twist. The fermionic fields then become the following scalars/one-forms

$$\begin{aligned}\psi_+^{am} &\rightarrow \rho_{\bar{z}}^{am}, \\ \bar{\psi}_+^{am} &\rightarrow \bar{\chi}^{am}, \\ \psi_-^{am} &\rightarrow \chi^{am}, \\ \bar{\psi}_-^{am} &\rightarrow \bar{\rho}_z^{am},\end{aligned}\tag{10}$$

and we can write

$$\begin{aligned}S &= \int d^2z \left(g_{am,bn} (\partial_{\bar{z}} \phi^{am} \partial_z \bar{\phi}^{bn} + \bar{\rho}_z^{bn} D_{\bar{z}} \chi^{am} + \rho_{\bar{z}}^{am} D_z \bar{\chi}^{bn}) \right. \\ &\quad \left. - R_{cp,bn,dq,am} \bar{\rho}_z^{cp} \chi^{bn} \bar{\chi}^{dq} \rho_{\bar{z}}^{am} + \int \Phi^* \omega \right) \\ &= S_{pert.} + \int \Phi^* \omega,\end{aligned}\tag{11}$$

where the map $\Phi : \mathbb{C}P^1 \rightarrow \Omega SU(k)$ is of integer degree N .

Quasi-Topological A-Model on $\mathbb{C}P^1$ with Target $\Omega SU(k)$

Like the fermion fields, there are two (nilpotent) scalar supercharges \bar{Q}_+ and Q_- , which SUSYs are therefore preserved on a worldsheet of any genus. In particular, \bar{Q}_+ generates the transformations

$$\begin{aligned}\delta\phi^{am} &= 0, \\ \delta\bar{\phi}^{am} &= \bar{\epsilon}_- \bar{\chi}^{am}, \\ \delta\rho_z^{am} &= -\bar{\epsilon}_- \partial_z \phi^{am}, \\ \delta\bar{\rho}_z^{am} &= -\bar{\epsilon}_- \bar{\Gamma}_{bn,cp}^{am} \bar{\chi}^{bn} \bar{\rho}_z^{cp}, \\ \delta\chi^{am} &= 0, \\ \delta\bar{\chi}^{am} &= 0,\end{aligned}\tag{12}$$

where $\bar{\epsilon}_-$ is a scalar grassmanian parameter.

Quasi-Topological A-Model on $\mathbb{C}P^1$ with Target $\Omega SU(k)$

- The action (11) can be cast into the form

$$S = \int d^2z \{ \bar{Q}_+, W'(t) \} + \cdots + tN \quad (13)$$

where $W'(t)$ is a metric-dependent combination of fields with metric scale t , and the ellipsis indicates additional terms which are metric-independent but depend on the complex structure of the target space.

- Although the stress tensor T_{zz} (i.e. $\delta S / \delta g_{zz}$) is \bar{Q}_+ -closed, it is generically **not** \bar{Q}_+ -exact; only $T_{\bar{z}\bar{z}}$ is \bar{Q}_+ -exact. So, the correlation function of \bar{Q}_+ -closed (but not exact) observables $\tilde{\mathcal{O}}$ is not completely independent of arbitrary deformations the worldsheet metric g . This is the **quasi-topological** A-model.
- Path integral localizes to \bar{Q}_+ -fixed points, and from (12), these are **holomorphic maps** from $\mathbb{C}P^1$ to $\Omega SU(k)$.

Quasi-Topological A-Model on $\mathbb{C}P^1$ with Target $\Omega SU(k)$

- The \overline{Q}_+ -cohomology of the model has **ground and left-excited states**, and the relevant operator observables $\tilde{\mathcal{O}}$ of **holomorphic dimension zero and positive** are Čech cohomology classes of the sheaf $\widehat{\Omega}^{ch}$ of chiral de Rham complex on $\mathcal{M}(\mathbb{C}P^1 \xrightarrow[N]{hol.} \Omega SU(k))$ [3].
- A correlation function of observables $\tilde{\mathcal{O}}$ has the form

$$\langle \prod_{\gamma} \tilde{\mathcal{O}}_{\gamma} \rangle = \sum_N e^{-tN} \left(\int_{F_N} \mathcal{D}\phi \mathcal{D}\bar{\phi} \mathcal{D}\rho_z \mathcal{D}\bar{\rho}_z \mathcal{D}\chi \mathcal{D}\bar{\chi} e^{-\int d^2z (\{\overline{Q}_+, W'(t)\} + \dots)} \prod_{\gamma} \tilde{\mathcal{O}}_{\gamma} \right). \quad (14)$$

Notice that

$$\frac{d}{dt} \left(\int_{F_N} \mathcal{D}\phi \dots \mathcal{D}\bar{\chi} e^{-\int d^2z (\{\overline{Q}_+, W'(t)\} + \dots)} \prod_{\gamma} \tilde{\mathcal{O}}_{\gamma} \right) = \langle \{\overline{Q}_+, \dots\} \rangle_{pert.} = 0 \quad (15)$$

so we can compute the path integral over F_N in (14), henceforth denoted as $\langle \prod_{\gamma} \tilde{\mathcal{O}}_{\gamma} \rangle_{pert.}$, at any convenient value of t , whilst keeping the original value of t in the constant factor e^{-tN} (due to worldsheet instantons).

Appearance of Toroidal and Affine $SU(k)$ Algebra in the \overline{Q}_+ -Cohomology

- **Isometries** of the target space inherited as **worldsheet symmetries** of the sigma model.
- Since $\Omega G \cong LG/G$, our sigma model ought to have an $LSU(k)$ symmetry on the worldsheet.
- Indeed, the corresponding **Noether currents**, the J 's, which charges generate a symmetry of the sigma model, can be shown to obey a current algebra associated with $LSU(k)$.
- As the J 's generate a symmetry, they act to leave the \overline{Q}_+ -cohomology of operator observables invariant. Thus, they ought to be \overline{Q}_+ -closed (but not exact), and are therefore also **in the \overline{Q}_+ -cohomology**, as one can verify.

Appearance of Toroidal and Affine $SU(k)$ Algebra in the \overline{Q}_+ -Cohomology

- We can conveniently compute the correlation functions of the J 's and T_{zz} via a large t limit, as explained in (14)-(15), and as OPEs, they are (in worldsheet instanton sector N)

$$J_z^{an_1}(z) J_z^{bn_2}(w) \sim \frac{if_c^{ab} J_z^{c\{n_1+n_2\}}(w)}{z-w}, \quad (16)$$

and

$$T_{zz}(z) J_z^{ak}(w) \sim \frac{J_z^{ak}(w)}{(z-w)^2} + \frac{\partial J_z^{ak}(w)}{(z-w)}. \quad (17)$$

Appearance of Toroidal and Affine $SU(k)$ Algebra in the \overline{Q}_+ -Cohomology

- Laurent expanding, these correspond to the **double loop algebra** $LL\mathfrak{su}(k)$

$$[J_{m_1}^{an_1}, J_{m_2}^{bn_2}] = if_c^{ab} J_{m_1+m_2}^{c\{n_1+n_2\}}, \quad (18)$$

and

$$[L_n, J_m^{ak}] = -mJ_{n+m}^{ak}. \quad (19)$$

- In the **holomorphic dimension zero sector**, the corresponding operator $L_0 = \oint dz z T_{zz}$ must act trivially, i.e., be \overline{Q}_+ -exact, and from (19), we see that $m = 0$, whence $LL\mathfrak{su}(k)$ reduces to the **loop algebra** $L\mathfrak{su}(k)$:

$$[J_0^{an_1}, J_0^{bn_2}] = if_c^{ab} J_0^{c\{n_1+n_2\}}. \quad (20)$$

This is also the **topological sector**, since T_{zz} is also \overline{Q}_+ -exact.

Appearance of Toroidal and Affine $SU(k)$ Algebra in the \overline{Q}_+ -Cohomology

- Our aforementioned J 's were derived from a classical Lagrangian density, and there would be quantum corrections.
- This means that the aforementioned algebras ought to be modified as well. Specifically, they will acquire central extensions.
- This leads us to a **toroidal lie algebra** $\mathfrak{su}(k)_{\text{tor}}$:

$$[J_{m_1}^{an_1}, J_{m_2}^{bn_2}] = if_c^{ab} J_{m_1+m_2}^{c\{n_1+n_2\}} + c_1 n_1 \delta^{ab} \delta^{\{n_1+n_2\}0} \delta_{\{m_1+m_2\}0} + c_2 m_1 \delta^{ab} \delta^{\{n_1+n_2\}0} \delta_{\{m_1+m_2\}0} \quad (21)$$

and **affine Lie algebra** $\mathfrak{su}(k)_{\text{aff}}$:

$$[J_0^{an_1}, J_0^{bn_2}] = if_c^{ab} J_0^{c\{n_1+n_2\}} + c_1 n_1 \delta^{ab} \delta^{\{n_1+n_2\}0} \quad (22)$$

in the \overline{Q}_+ -cohomology (for some $c_{1,2}$).

Modules over the Toroidal and Affine $SU(k)$ Algebra in the \overline{Q}_+ -Cohomology

- Now, acting on a ground state $|0\rangle$ (which is \overline{Q}_+ -closed) with the generators of $\mathfrak{su}(k)_{\text{tor}}$, we have the states

$$J_{-m_1}^{a\{-n_1\}} J_{-m_2}^{b\{-n_2\}} J_{-m_3}^{c\{-n_3\}} \dots |0\rangle, \quad (23)$$

where $m_j, n_i \geq 0$.

- They span a module $\widehat{\mathfrak{su}}(k)_{c_1, c_2}^N$ over the toroidal Lie algebra $\mathfrak{su}(k)_{\text{tor}}$ of levels c_1 and c_2 .
- These states have nonzero holomorphic dimension (according to (19)), and can be shown to be elements of the \overline{Q}_+ -cohomology.
- Thus, via the state-operator correspondence, we have

$$H_{\text{Čech}}^* \left(\widehat{\Omega}_{\mathcal{M}(\mathbb{C}P^1 \xrightarrow[N]{\text{hol.}} \Omega SU(k))}^{ch} \right) = \widehat{\mathfrak{su}}(k)_{c_1, c_2}^N. \quad (24)$$

Modules over the Toroidal and Affine $SU(k)$ Algebra in the \overline{Q}_+ -Cohomology

- In the topological sector where $m_i = 0$, the states are

$$J_0^{a\{-n_1\}} J_0^{b\{-n_2\}} J_0^{c\{-n_3\}} \dots |0\rangle, \quad (25)$$

where $n_i \geq 0$.

- They span a module $\widehat{su}(k)_{c_1}^N$ over the affine Lie algebra $\mathfrak{su}(k)_{\text{aff}}$ of level c_1 .
- These states have zero holomorphic dimension (according to (19)), and persist as elements of the \overline{Q}_+ -cohomology.
- Thus, via the state-operator correspondence, and the fact that the zero holomorphic dimension chiral de Rham complex is just the de Rham complex [3], we have

$$H_{L^2}^*(\mathcal{M}(\mathbb{C}P^1 \xrightarrow[\text{hol.}]{N} \Omega SU(k))) = \widehat{su}(k)_{c_1}^N. \quad (26)$$

The 6d A_{k-1} $\mathcal{N} = (2, 0)$ Little String Theory

- Little string theories (LST) exist in 6d spacetimes, and reduce to interacting local QFTs when string length $l_s \rightarrow 0$.
- The 6d A_{k-1} $\mathcal{N} = (2, 0)$ LST, in particular, reduces to the 6d A_{k-1} $\mathcal{N} = (2, 0)$ superconformal field theory - has **no known classical action**. Rich theory, so corresponding LST must be at least just as rich.
- It is also the worldvolume theory of a stack of NS5-branes in type IIA string theory, whereby the fundamental strings which reside within the branes with coupling $g_s \rightarrow 0$ (whence bulk d.o.f., including gravity, decouple) and $l_s \not\rightarrow 0$, are the little strings.

The Ground and Left-Excited Spectrum of the 6d A_{k-1} $(2,0)$ LST

- The discrete lightcone quantization (DLCQ) of the LST on $\mathbb{R}^{5,1}$ describes it as a 2d $\mathcal{N} = (4,4)$ sigma model on $S^1 \times R$ with target $\mathcal{M}_{SU(k)}^N(\mathbb{R}^4)$, the moduli space of $SU(k)$ N -instantons on \mathbb{R}^4 . Here, $k =$ no. of branes, $N =$ units of discrete momentum along the S^1 [4].
- The ground states of the LST are given by sigma model states annihilated by all the supercharges, i.e., they correspond to harmonic forms and thus L^2 -cohomology classes of $\mathcal{M}_{SU(k)}^N(\mathbb{R}^4)$.
- The left-excited states of the LST are given by sigma model states annihilated by the four chiral supercharges, i.e., they correspond to Čech cohomology classes of the sheaf $\widehat{\Omega}_{\mathcal{M}_{SU(k)}^N(\mathbb{R}^4)}^{ch}$.

The Ground and Left-Excited Spectrum of the $6d A_{k-1}$ $(2,0)$ LST

- According to Atiyah [1], we have the identification

$$\mathcal{M}_G^N(\mathbb{R}^4) \cong \mathcal{M}(\mathbb{C}P^1 \xrightarrow[\text{hol.}]{N} \Omega G). \quad (27)$$

- In turn, this means we can identify

$$H_{L^2}^*(\mathcal{M}_{SU(k)}^N(\mathbb{R}^4)) \cong H_{L^2}^*(\mathcal{M}(\mathbb{C}P^1 \xrightarrow[\text{hol.}]{N} \Omega SU(k))) \quad (28)$$

and

$$H_{\check{C}ech}^*(\mathcal{M}_{SU(k)}^N(\mathbb{R}^4)) \cong H_{\check{C}ech}^*(\widehat{\Omega}^{ch} \mathcal{M}(\mathbb{C}P^1 \xrightarrow[\text{hol.}]{N} \Omega SU(k))) \quad (29)$$

The Ground and Left-Excited Spectrum of the 6d A_{k-1} $(2, 0)$ LST

- Thus, from (28) and (26), we find that

$$\boxed{\text{ground spectrum of 6d } A_{k-1} (2, 0) \text{ LST} = \text{modules of } \mathfrak{su}(k)_{\text{aff}}}$$

(30)

- Similarly, from (29) and (24), we find that

$$\boxed{\text{left-excited spectrum of 6d } A_{k-1} (2, 0) \text{ LST} = \text{modules of } \mathfrak{su}(k)_{\text{tor}}}$$

(31)

Deriving the Braverman-Finkelberg Relation and its Generalization

- From (26) and (28), we also find (c.f. [5]) that

$$\boxed{\mathrm{IH}^*(\mathcal{M}_{SU(k)}^N(\mathbb{R}^4)) = \widehat{su}(k)_{c_1}^N,} \quad (32)$$

This is the **Braverman-Finkelberg relation** in [2].

- From (24) and (29), we also find that

$$\boxed{H_{\check{\mathrm{C}}\mathrm{ech}}^*(\widehat{\Omega}_{\mathcal{M}_{SU(k)}^N(\mathbb{R}^4)}^{ch}) = \widehat{su}(k)_{c_1, c_2}^N.} \quad (33)$$

This is a novel **generalization of the Braverman-Finkelberg relation**.

The M5-brane Worldvolume Theory

- The setup of the k NS5-branes with type IIA fundamental strings bound to it as little strings, has an M-theoretic interpretation.
- They can be regarded as k M5-branes with M2-branes ending on them in one spatial direction (as M-strings) and wrapping the 11th circle of radius R in the other spatial direction, observed at low energy scales $\ll R^{-1}$.
- As such, the low energy DLCQ of this **worldvolume theory of k M5-branes** can also be understood via the LST described as a $\mathcal{M}_{SU(k)}^N(\mathbb{R}^4)$ sigma model [6].

1/2 BPS sector of M5-brane Worldvolume Theory

- Using the relevant SUSY algebras, we can show that the low energy **1/2 BPS sector** of the M5-brane theory is captured by the **ground states** of the LST.
- Thus, according to (30), the partition function of the 1/2 BPS sector ought to be given by summing representations of $\mathfrak{su}(k)_{\text{aff}}$. In particular, it is computed to be

$$Z_{1/2} = \sum_{\widehat{\lambda}'} \chi_{\widehat{\mathfrak{su}(k)_{c_1}}}^{\widehat{\lambda}'}(p) \quad (34)$$

where χ is a character of the module; $\widehat{\lambda}'$ a dominant highest weight; $p = e^{2\pi i\tau}$; and τ is the complex structure of an auxiliary torus.

- This is a cousin of a modular form which transforms as a representation of $SL(2, \mathbb{Z})$.
- There is an intrinsic $SL(2, \mathbb{Z})$ symmetry in the M5-brane worldvolume theory on $\mathbb{R}^{5,1}$, which emerges as gauge-theoretic S-duality of 4d $\mathcal{N} = 4$ SYM after compactifying on T^2 !

1/4 BPS sector of M5-brane Worldvolume Theory

- Using the relevant SUSY algebras, we can show that the low energy **1/4 BPS sector** of the M5-brane theory is captured by the **left-excited states** of the LST.
- Thus, according to (31), the partition function of the 1/4 BPS sector ought to be given by summing representations of $\mathfrak{su}(k)_{\text{tor}}$. In particular, it is computed to be

$$Z_{1/4} = q^{\frac{1}{24}} \sum_{\hat{\lambda}} \chi_{\widehat{\mathfrak{su}(k)}_{c_1}}^{\hat{\lambda}}(p) \frac{1}{\eta(\tau)} \quad (35)$$

where η is the Dedekind eta function; $q = e^{2\pi i\sigma}$; and σ is the Kähler structure of an auxiliary torus.

- This is a cousin of an automorphic form which transforms as a representation of $SO(2, 2; \mathbb{Z})$.
- There is an intrinsic $SO(2, 2; \mathbb{Z})$ symmetry of the M5-brane worldvolume theory on $\mathbb{R}^{5,1}$, which emerges as string-theoretic T-duality of little strings after compactifying on T^2 !

Conclusion

- We have explained how a quasi-topological $\Omega SU(k)$ sigma model can be used to help us (i) understand the 6d $A_{k-1}(2,0)$ LST; (ii) derive and generalize the Braverman-Finkelberg relation; (iii) understand the M5-brane worldvolume theory.
- Notably, we find that the chiral spectrum of the little string is furnished by representations of a toroidal algebra, and the BPS spectrums of the M5-brane worldvolume theory are closely related to modular and automorphic forms.
- Consistent with these aforementioned physical results is a geometric Langlands correspondence for surfaces – the Braverman-Finkelberg relation – and its generalization, which we also physically derived.
- We see a nice interconnection between string theory, M-theory, geometric representation theory and number theory.

- To ascertain the **full chiral plus anti-chiral spectrum** of the the 6d $A_{k-1} (2, 0)$ LST. We expect it to be furnished by representations of a holomorphic plus antiholomorphic (positive-moded) toroidal algebra.
- Gauge the $\Omega SU(k)$ sigma model to obtain a derivation and **generalization of the AGT correspondence**, which we expect will relate the equivariant Cech-cohomology of the sheaf of chiral de Rham complex on $\mathcal{M}_{SU(k)}^N(\mathbb{R}^4)$ to toroidal W -algebras.
- Go **beyond the BPS sector** of the M5-brane worldvolume theory as captured by the full spectrum of the LST. We expect the corresponding worldvolume partition function to consist of the $1/4$ BPS partition function with an extra Dedekind eta function in $\bar{\tau}$.

THANKS FOR LISTENING!

- [1] M. F. Atiyah, *Instantons in Two and Four Dimensions*, *Communications in Mathematical Physics* **93** (Dec., 1984) 437–451.
- [2] A. Braverman and M. Finkelberg, *Pursuing the Double Affine Grassmannian I: Transversal Slices via Instantons on A_k -singularities*, *Duke Mathematical Journal* **152** (Apr., 2010) 175–206.
- [3] M.-C. Tan, *Two-Dimensional Twisted Sigma Models And The Theory of Chiral Differential Operators*, *Adv.Theor.Math.Phys.* 10, 759-851 (2006).
- [4] O. Aharony, *A Brief Review of 'Little String Theories'*, *Classical and Quantum Gravity* **17** (Mar., 2000) 929–938.
- [5] M. Goresky, *L^2 Cohomology is Intersection Cohomology*, in *The Zeta Functions of Picard Modular Surfaces* (ed. R. P. Langlands and D. Ramakrishnan), Centre de Recherches Mathématiques, Montréal, 1992.
- [6] R. Dijkgraaf, “The Mathematics of Fivebranes”, *Documenta Mathematica*, 133-142 (1998).