

The gauged WZ term with boundary

José Figueroa-O'Farrill

Edinburgh Mathematical Physics Group



LMS Durham Symposium, 29 July 2005

A quaternion of sigma models

WZ	bWZ
gWZ	gbWZ

... and one of cohomology theories

de Rham	relative de Rham
equivariant	relative equivariant

References

- [hep-th/9407149](#)
- *Phys. Lett.* **B341** (1994) 153–159, [hep-th/9407196](#)
- *JHEP* **01** (2001) 006, [hep-th/0008038](#)

with Sonia Stanciu

References

- [hep-th/9407149](#)
- *Phys. Lett.* **B341** (1994) 153–159, [hep-th/9407196](#)
- *JHEP* **01** (2001) 006, [hep-th/0008038](#)

with Sonia Stanciu

- [hep-th/0506049](#)

with Nouri Mohammedi

Sigma models

- Two oriented pseudo-riemannian manifolds: Σ^d, X^n
- $\partial\Sigma$ may or may not be empty
- $\varphi : \Sigma \rightarrow X, d\varphi \in \Omega^1(\Sigma, \varphi^*TX)$
- sigma model action

$$I_\sigma = \int_\Sigma \frac{1}{2} |d\varphi|^2$$

defines variational problem for harmonic maps $d \star_\Sigma d\varphi = 0$



The Wess–Zumino term

- $H \in \Omega_{\text{closed}}^{d+1}(X)$
- assume $\partial\Sigma = \emptyset$ and $\varphi(\Sigma) \subset X$ bounds

$$\varphi(\Sigma) = \partial M \quad \exists M \subset X$$

- Wess–Zumino term

$$I_{\text{WZ}} = \int_M H$$



- $dH = 0 \implies$ field equations for φ

$$\delta I_{\text{WZ}} = \int_M \mathcal{L}_{\delta\varphi} H = \int_M d\iota_{\delta\varphi} H = \int_{\varphi(\Sigma)} \iota_{\delta\varphi} H$$

\therefore classical theory is independent of choice of M

- quantum theory depends on I_{WZ} modulo $2\pi\mathbb{Z}$

\therefore quantum theory is independent of M if $[\frac{1}{2\pi}H]$ is an integral class in $H_{\text{dR}}^{d+1}(X)$



Example: the WZW model

- Σ two-dimensional
- X a compact simple Lie group
- $H_2(X) = 0$ so $\varphi(\Sigma)$ always bounds [Cartan]
- $[\frac{1}{2\pi}H]$ is k times the generator of $H^3(X; \mathbb{Z}) \cong \mathbb{Z}$, where k is the level of the WZW model



The presence of a boundary

- suppose $\partial\Sigma \neq \emptyset \implies$ need to specify boundary conditions
- let $i : Y \hookrightarrow X$, $i^*H = dB$ for some $B \in \Omega^d(Y)$
- BCs: $\varphi(\partial\Sigma) \subset Y$

\implies a theory of relative maps

$$\varphi : (\Sigma, \partial\Sigma) \rightarrow (X, Y)$$



The boundary Wess–Zumino term

- $\varphi(\Sigma) \subset X$ is not a cycle, but it is a cycle modulo Y :

$$\partial\varphi(\Sigma) = \varphi(\partial\Sigma) \subset Y$$

so suppose that it bounds modulo Y :

$$\exists M \subset X, \quad D \subset Y \quad \text{s.t.} \quad \partial M = \varphi(\Sigma) + D$$

whence $\partial D = -\varphi(\partial\Sigma)$



- boundary Wess–Zumino term

$$I_{\text{bWZ}} = \int_M H - \int_D B$$

- B only enters in the boundary conditions:

$$\delta I_{\text{bWZ}} = \int_M d\iota_{\delta\varphi} H - \int_D \mathcal{L}_{\delta\varphi} B = \int_{\varphi(\Sigma)} \iota_{\delta\varphi} H + \int_{\varphi(\partial\Sigma)} \iota_{\delta\varphi} B$$

and field equations are not otherwise changed



\implies classical theory is again independent on choice of M and D , but quantum theory is not unless $[\frac{1}{2\pi}H]$ is an integral class in the relative de Rham cohomology $H_{\text{dR}}^{d+1}(X, Y)$



Example: the boundary WZW model

- $Y \subset X$ a conjugacy class
- $[\frac{1}{2\pi}H] \in H^3(X, Y; \mathbb{Z})$ selects a discrete set of conjugacy classes corresponding to unitary highest weight representations of the affine Lie algebra at level k



Symmetries of sigma model with WZ term

- G a connected Lie group, with Lie algebra \mathfrak{g} with basis Z_a
- G acts on X isometrically preserving H
- $Z_a \mapsto v_a$ a Killing vector, $[v_a, v_b] = f_{ab}{}^c v_c$
- let $\iota_a := \iota_{v_a}$ and $\mathcal{L}_a := \mathcal{L}_{v_a}$, then $\mathcal{L}_a = d\iota_a + \iota_a d$
- $\mathcal{L}_a H = 0$, equivalently $d\iota_a H = 0$



Gauging a sigma model

- means coupling it to a gauge field $A \in \Omega^1(\Sigma, \mathfrak{g})$ so that the action is invariant under (infinitesimal) gauge transformations:

$$\delta_\lambda A = d\lambda + [A, \lambda]$$

$$\delta_\lambda \omega = d\lambda^a \wedge \iota_a \omega + \lambda^a \mathcal{L}_a \omega$$

where $\omega \in \Omega(X)$ and $\lambda \in C^\infty(\Sigma, \mathfrak{g})$

- can write $\delta_\lambda A = d\lambda^a \wedge \iota_a A + \lambda^a \mathcal{L}_a A$ by defining

$$\iota_a A^b = \delta_a^b \quad \iota_a F^b = 0 \quad \text{and} \quad \mathcal{L}_a = d\iota_a + \iota_a d$$

where $F = dA + \frac{1}{2}[A, A]$ is the field-strength



Minimal coupling

- $I_\sigma = \int_\Sigma \frac{1}{2} |d\varphi|^2$ can be gauged by minimal coupling:

$$d\varphi \mapsto d_A\varphi := d\varphi - A^a \iota_a d\varphi$$

but I_{WZ} is a different matter: the minimally coupled H need not be closed



Gauging the WZ term

- means extending H to a closed gauge-invariant form \mathcal{H} :

$$\mathcal{H} = H + \text{terms involving } A \text{ and } F$$

such that $d\mathcal{H} = 0$ and

$$\delta_\lambda \mathcal{H} = d\lambda^a \wedge \iota_a \mathcal{H} + \lambda^a \mathcal{L}_a \mathcal{H} = 0$$

equivalently $\iota_a \mathcal{H} = 0$ (and $\mathcal{L}_a \mathcal{H} = 0$)



Differential graded algebras

- a \mathfrak{g} -dga \mathfrak{A} :

- ★ $\mathfrak{A} = \bigoplus_{i \geq 0} \mathfrak{A}^i$

- ★ (associative, supercommutative) product

$$\wedge : \mathfrak{A}^i \otimes \mathfrak{A}^j \rightarrow \mathfrak{A}^{i+j}$$

- ★ derivation $d : \mathfrak{A}^i \rightarrow \mathfrak{A}^{i+1}$

- ★ derivation $\iota_a : \mathfrak{A}^i \rightarrow \mathfrak{A}^{i-1}$

- ★ derivation $\mathcal{L}_a = d\iota_a + \iota_a d$ defines \mathfrak{g} -action



- $\Omega(X)$ is a \mathfrak{g} -dga
- so is the Weyl algebra

$$\mathfrak{W} = \Lambda \mathfrak{g}^* \otimes \mathfrak{S} \mathfrak{g}^*$$

with generators $\mathcal{A}^a \in \Lambda^1 \mathfrak{g}^*$ and $\mathcal{F}^a \in \mathfrak{S}^1 \mathfrak{g}^*$ and where $\iota_a \mathcal{A}^b = \delta_a^b$ and $\iota_a \mathcal{F}^b = 0$ and $d\mathcal{A}^a = \mathcal{F}^a - \frac{1}{2} f_{bc}^a \mathcal{A}^b \wedge \mathcal{A}^c$

- Weyl homomorphism $w : \mathfrak{W} \rightarrow \Omega(\Sigma, \mathfrak{g})$ defined by $\mathcal{A}^a \mapsto A^a$ and $\mathcal{F}^a \mapsto F^a$
- $\Omega(X) \otimes \mathfrak{W}$ is a \mathfrak{g} -dga



Equivariant forms

- \mathfrak{A} a \mathfrak{g} -dga, then $\phi \in \mathfrak{A}$ is
 - ★ horizontal, if $\iota_a \phi = 0$
 - ★ invariant, if $\mathcal{L}_a \phi = 0$
 - ★ equivariant, if both
 - $\Omega_{\mathfrak{g}}(X)$: subcomplex of equivariant elements of $\Omega(X) \otimes \mathfrak{W}$
 - $\{w(\phi) \mid \phi \in \Omega_{\mathfrak{g}}(X)\}$ are gauge-invariant
- \therefore gauging the WZ term is finding an equivariant closed extension
 $\mathcal{H} \in \Omega_{\mathfrak{g}}^{d+1}(X)$ of $H \in \Omega^{d+1}(X)$



The Cartan model

- in a local gauge-invariant quantity, A only appears in minimally coupled expressions (or through F)
- this suggests defining

$$\Omega_C(X) := (\Omega(X) \otimes \mathfrak{S}\mathfrak{g}^*)^{\mathfrak{g}}$$

called the **Cartan model** of $\Omega_{\mathfrak{g}}(X)$

- indeed, $\Omega_C(X) \cong \Omega_{\mathfrak{g}}(X)$



- $\pi : \Omega_{\mathfrak{g}}(X) \xrightarrow{\cong} \Omega_C(X)$ consists in setting $\mathcal{A} = 0$
- $\pi^{-1} : \Omega_C(X) \xrightarrow{\cong} \Omega_{\mathfrak{g}}(X)$ consists of minimal coupling
- induced differential $d_C = \pi \circ d \circ \pi^{-1} : \Omega_C^p(X) \rightarrow \Omega_C^{p+1}(X)$ is

$$d_C \mathcal{F}^a = 0 \quad \text{and} \quad d_C \omega = d\omega - \iota_a \omega \mathcal{F}^a$$

for $\omega \in \Omega(X)$

- gauging WZ term is equivalent to finding a d_C -closed extension $\mathcal{H}_C \in \Omega_C(X)$



The Hull–Spence obstructions

- write the most general \mathcal{H}_C in the Cartan model

$$\mathcal{H}_C = H + \theta_a \mathcal{F}^a + \frac{1}{2} \theta_{ab} \mathcal{F}^a \mathcal{F}^b + \dots$$

where $\theta_a \in \Omega^{d-1}(X)$, $\theta_{ab} \in \Omega^{d-3}(X), \dots$ satisfy

$$\mathcal{L}_a \theta_b = f_{ab}^c \theta_c \quad \mathcal{L}_a \theta_{bc} = f_{ab}^d \theta_{dc} + f_{ac}^d \theta_{bd} \quad \dots$$



- splitting $d_C \mathcal{H}_C = 0$ into types:

$$\iota_a H = d\theta_a \quad \iota_a \theta_b + \iota_b \theta_a = d\theta_{ab} \quad \dots$$

which are the first two Hull–Spence obstructions

- overcoming these obstructions yields \mathcal{H}_C and minimal coupling yields \mathcal{H} and the gauged WZ term

$$I_{\text{gWZ}} = \int_M \mathcal{H}$$



The two-dimensional case

- $\mathcal{H}_C = H + \theta_a \mathcal{F}^a$
- $d_C \mathcal{H}_C = 0$ implies

$$\iota_a H = d\theta_a \quad \text{and} \quad \iota_a \theta_b + \iota_b \theta_a = 0$$

- $\mathcal{L}_a \mathcal{H}_C = 0$ implies

$$\mathcal{L}_a \theta_b = f_{ab}{}^c \theta_c$$



- the gauged WZ term is

$$I_{\text{gWZ}} = \int_M H + \int_{\Sigma} (\varphi^* \theta_a \wedge A^a + \frac{1}{2} \varphi^* (\iota_a \theta_b) A^a \wedge A^b)$$

[Hull & Spence; Jack, Jones, Mohammadi & Osborn]

- to this action we can add a Yang–Mills term $\int_{\Sigma} \frac{1}{4} |F|^2$
- or other topological terms, corresponding to cocycles in $\Omega_{\mathfrak{g}}^3(X)$



Example: the gauged WZW model

- we try to gauge $\mathfrak{g} \subset \mathfrak{x} \oplus \mathfrak{x}$ defined by homomorphisms

$$\ell : \mathfrak{g} \rightarrow \mathfrak{x} \quad \text{and} \quad r : \mathfrak{g} \rightarrow \mathfrak{x}$$

- the only obstruction is

$$\ell^* \langle -, - \rangle = r^* \langle -, - \rangle$$

with $\langle -, - \rangle$ is the ad -invariant scalar product on \mathfrak{x}



- typical “anomaly-free” \mathfrak{g} :
 - ★ diagonal: $\ell = r$
 - ★ twisted diagonal: $\ell = r \circ \tau$ for some isometry $\tau \in \text{Aut}(\mathfrak{x})$
 - ★ chiral: $r = 0$ and $\mathfrak{g} \subset \mathfrak{x}$ an isotropic subalgebra



The twisted Courant algebroid on $T \oplus \Lambda^{d-1}T^*$

- $TX \oplus \Lambda^{d-1}T^*X$ has a $\Lambda^{d-2}T^*X$ -valued bilinear

$$\langle v + \alpha, w + \beta \rangle = \iota_v \beta + \iota_w \alpha \in \Lambda^{d-2}T^*X$$

- and also has a Courant bracket:

$$[v + \alpha, w + \beta] = [v, w] + \mathcal{L}_v \beta - \mathcal{L}_w \alpha - \frac{1}{2}d(\iota_v \beta - \iota_w \alpha)$$



- $H \in \Omega_{\text{closed}}^{d+1}(X)$ twists the bracket

$$[v + \alpha, w + \beta]_H = [v + \alpha, w + \beta] - \iota_v \iota_w H$$

- we say $L \subset TX \oplus \Lambda^{d-1}T^*X$ is **isotropic** if for all $v + \alpha \in C^\infty(L)$,
 $\iota_v \alpha \in \Omega_{\text{exact}}^{d-2}(X)$
- an isotropic, involutive L defines a Lie algebroid over X



The Lie algebroid of the gauged WZ term

- let L be the image of $\mathfrak{g} \rightarrow C^\infty(TX \oplus \Lambda^{d-1}T^*X)$ given by

$$Z_a \mapsto v_a + \theta_a$$

- $\iota_a \theta_b + \iota_b \theta_a = d\theta_{ab}$, whence L is isotropic
- $d\theta_a = \iota_a H$ and $\mathcal{L}_a \theta_b = f_{ab}^c \theta_c$ then imply that L is involutive
- so L defines a Lie algebroid isomorphic to \mathfrak{g}

[cf. Alekseev & Strobl ($d = 2$), Bonelli & Zabzine]



Gauging the boundary WZ term

- assume G acts preserving (Y, B)
- gauging I_{bWZ} consists in finding equivariant extensions \mathcal{H} and \mathcal{B} of H and B such that $d\mathcal{H} = 0$ and $i^*\mathcal{H} = d\mathcal{B}$ on Y
- or in the Cartan model finding \mathcal{H}_C and \mathcal{B}_C with $d_C\mathcal{H}_C = 0$ and $i^*\mathcal{H}_C = d_C\mathcal{B}_C$
- the gauged boundary WZ term is then

$$I_{\text{gbWZ}} = \int_M \mathcal{H} - \int_D \mathcal{B}$$



Two-dimensional gauged boundary WZ term

- $\mathcal{B}_C = B + h_a \mathcal{F}^a$, where $h_a \in C^\infty(X)$ obeying

$$\mathcal{L}_a h_b = f_{ab}{}^c h_c$$

- $i^* \mathcal{H}_C = d_C \mathcal{B}_C$ is equivalent to

$$i^* \theta_a + i_a B = dh_a$$



and

$$I_{\text{gbWZ}} = \int_M H - \int_D B + \int_{\Sigma} (\varphi^* \theta_a \wedge A^a + \frac{1}{2} \varphi^* (\iota_a \theta_b) A^a \wedge A^b) + \int_{\partial \Sigma} \varphi^* h_a A^a$$

- one can add topological terms, corresponding to relative cocycles in $\Omega_{\mathfrak{g}}^3(X, Y)$



The boundary Lie algebroid

- the twisted Courant bracket restricts to $TY \oplus \Lambda^{d-1}T^*Y$, but since $i^*H = dB$, it is B -related to the untwisted Courant bracket:

$$[v + \alpha, w + \beta]_{dB} = [e^B(v + \alpha), e^B(w + \beta)]$$

- the image of the map $\mathfrak{g} \rightarrow C^\infty(TY \oplus \Lambda^{d-1}T^*Y)$ defined by

$$Z_a \mapsto e^B(v_a + i^*\theta_a) = v_a + \iota_a B + i^*\theta_a = v_a + dh_a$$

is isotropic and involutive: a Lie algebroid on Y isomorphic to \mathfrak{g}



Example: the gauged boundary WZW model

- possible boundary conditions are G -orbits:
 - ★ (twisted) diagonal gaugings: (twisted) conjugacy classes
 - ★ chiral gaugings: cosets
- boundary offers no new obstructions
- $dh_a = 0$ and $h_a \in [\mathfrak{g}, \mathfrak{g}]^o$
- boundary Lie algebroid is the action Lie algebroid of \mathfrak{g} on Y



Open questions

- are there CFT constructions for the cosets?
- relation with boundary conditions on $X \times X$?
- relation with boundary integrable systems?
- (oidish) interpretation for 'higher' obstructions?



Thank you!

