



## EXAMINATION PAPER

<b>Examination Session:</b> May	<b>Year:</b> 2018	<b>Exam Code:</b> MATH3291-WE01
------------------------------------	----------------------	------------------------------------

<b>Title:</b> Partial Differential Equations III
---

Time Allowed:	3 hours	
Additional Material provided:	None	
Materials Permitted:	None	
Calculators Permitted:	Yes	Models Permitted: Casio fx-83 GTPLUS or Casio fx-85 GTPLUS.
Visiting Students may use dictionaries: No		

Instructions to Candidates:	Credit will be given for: the best <b>FOUR</b> answers from Section A and the best <b>THREE</b> answers from Section B. Questions in Section B carry <b>TWICE</b> as many marks as those in Section A.	
		<b>Revision:</b>

## SECTION A

1. Consider the conservation law

$$\begin{cases} u_t + uu_x = 0 & \text{for } (x, t) \in \mathbb{R} \times (0, T), \\ u(x, 0) = e^{-(x-1)^2} & \text{for } x \in \mathbb{R}. \end{cases} \quad (1)$$

- (a) Find the largest value of  $T \in \mathbb{R}$  for which this conservation law has a classical solution  $u : \mathbb{R} \times [0, T) \rightarrow \mathbb{R}$ .
- (b) Give a sketch of the characteristics of (1) up until time  $T$ .
- (c) Find an implicit equation for  $u$  that does not contain partial derivatives.

2. Consider the conservation law

$$\begin{cases} u_t + (f(u))_x = 0 & \text{for } (x, t) \in \mathbb{R} \times (0, \infty), \\ u(x, 0) = g(x) & \text{for } x \in \mathbb{R}. \end{cases} \quad (2)$$

- (a) Define the Rankine-Hugoniot condition for a shock.
- (b) Define the Lax entropy condition for a shock.
- (c) Let

$$g(x) = \begin{cases} 1 & \text{for } x < 0, \\ 0 & \text{for } x > 0, \end{cases} \quad \text{and} \quad f(u) = \frac{1}{4}u^4.$$

Consider the functions

$$u_1(x, t) = \begin{cases} 1 & \text{for } x < t/2, \\ 0 & \text{for } x > t/2, \end{cases} \quad \text{and} \quad u_2(x, t) = \begin{cases} 1 & \text{for } x < t/4, \\ 0 & \text{for } x > t/4. \end{cases}$$

Which one of these two functions is a weak solution of (2)?

Is it also an admissible solution in the sense of the Lax entropy condition?

3. State the theorem of local existence for first-order quasilinear PDEs

$$\begin{cases} a_1(x, y, u)u_x + a_2(x, y, u)u_y = b(x, y, u) & \text{for } (x, y) \in \Omega, \\ u(x, y) = u_0(x, y) & \text{for } (x, y) \in \Gamma, \end{cases}$$

in the following steps:

- (a) State the assumptions on the Cauchy data  $\Gamma$  and  $u_0$ .
- (b) State the assumptions on the coefficients  $a_1$ ,  $a_2$ ,  $b$ .
- (c) State what is means for the data to be noncharacteristic.
- (d) State the result of the theorem.

4. (a) Let  $B_1(\mathbf{0}) \subset \mathbb{R}^2$  be the unit ball in two dimensions. Suppose that  $u : \overline{B_1(\mathbf{0})} \rightarrow \mathbb{R}$  satisfies

$$\begin{aligned} -\Delta u &= g && \text{in } B_1(\mathbf{0}), \\ \nabla u \cdot \mathbf{n} &= 1 + \alpha && \text{on } \partial B_1(\mathbf{0}), \end{aligned}$$

where

$$g(\mathbf{x}) = \begin{cases} \frac{1}{|\mathbf{x}|} \sin(2\pi|\mathbf{x}|) & \text{if } \mathbf{x} \neq \mathbf{0}, \\ 2\pi & \text{if } \mathbf{x} = \mathbf{0}, \end{cases}$$

and where  $\mathbf{n}$  is the outward-pointing unit normal vector field to  $\partial B_1(\mathbf{0})$ , and  $\alpha \in \mathbb{R}$  is a constant. Find  $\alpha$ .

- (b) State the Poincaré inequality for functions  $f \in C^1([a, b])$ .
5. (a) State the strong maximum principle for harmonic functions.
- (b) Let  $\Omega \subset \mathbb{R}^n$  be open, bounded and connected. Let  $u \in C^2(\Omega) \cap C(\overline{\Omega})$  satisfy

$$\begin{aligned} -\Delta u &= f && \text{in } \Omega, \\ u &= g && \text{on } \partial\Omega, \end{aligned}$$

and let  $v \in C^2(\Omega) \cap C(\overline{\Omega})$  satisfy

$$\begin{aligned} -\Delta v &= f && \text{in } \Omega, \\ v &= h && \text{on } \partial\Omega, \end{aligned}$$

where  $f, g, h : \overline{\Omega} \rightarrow \mathbb{R}$ . Assume that  $g \leq h$  and that there exists  $\mathbf{x}_0 \in \partial\Omega$  such that  $g(\mathbf{x}_0) < h(\mathbf{x}_0)$ . Show that  $u < v$  in  $\Omega$ .

6. Let  $\Omega \subset \mathbb{R}^2$  be open and bounded with smooth boundary. Let

$$V = \{\varphi \in C^1(\overline{\Omega}) : \varphi = 0 \text{ on } \partial\Omega\}.$$

Define  $E : V \rightarrow \mathbb{R}$  by

$$E[v] = \frac{1}{2} \int_{\Omega} \nabla v \cdot (A \nabla v) d\mathbf{x} - \int_{\Omega} f v d\mathbf{x},$$

where  $f : \Omega \rightarrow \mathbb{R}$  is smooth and  $A : \Omega \rightarrow \mathbb{R}^{2 \times 2}$  is a smooth, matrix-valued map such that  $A(\mathbf{x})$  is symmetric for all  $\mathbf{x} \in \Omega$ . Assume that  $u \in C^2(\overline{\Omega}) \cap V$  minimises  $E$ :

$$E[u] = \min_{v \in V} E[v].$$

Show that  $u$  satisfies the elliptic PDE

$$-\operatorname{div}(A \nabla u) = f \quad \text{in } \Omega.$$

## SECTION B

7. Let  $a \in \mathbb{R}$  be a constant. Consider the first order PDE

$$\begin{cases} au_x + xyu_y = xu, & (x, y) \in \mathbb{R}^2, \\ u(0, y) = f(y), & y \in \mathbb{R}, \end{cases} \quad (3)$$

where  $f : \mathbb{R} \rightarrow \mathbb{R}$  is an arbitrary continuous function.

- (a) For which values of  $a$  is the initial condition in (3) noncharacteristic?
- (b) Using the method of characteristics, solve (3) for all values of  $a$  for which the initial condition is noncharacteristic.
- (c) Take  $f(y) = y$ ,  $a = 0$ . Prove that (3) has infinitely many solutions.

8. Consider the scalar conservation law

$$\begin{cases} u_t + (f(u))_x = 0, \\ u(x, 0) = u_0(x), \end{cases}$$

where

$$u_0(x) = \begin{cases} 1 & \text{for } x < 0, \\ 3 & \text{for } x > 0, \end{cases} \quad \text{and} \quad f(u) = \frac{1}{3}u^3.$$

- (a) Write down the equation for the characteristics inside the regions

$$\{x < t\} \quad \text{and} \quad \{x > 9t\}.$$

- (b) Verify that the following are weak solutions:

$$u_1(x, t) = \begin{cases} 1 & \text{for } x < \frac{7}{3}t, \\ 2 & \text{for } \frac{7}{3}t < x < 4t, \\ \left(\frac{x}{t}\right)^{1/2} & \text{for } 4t < x < 9t, \\ 3 & \text{for } x > 9t, \end{cases} \quad u_2(x, t) = \begin{cases} 1 & \text{for } x < t, \\ \left(\frac{x}{t}\right)^{1/2} & \text{for } t < x < 9t, \\ 3 & \text{for } x > 9t. \end{cases}$$

- (c) Which of the solutions in part (b) satisfy the Lax entropy condition? Justify your answer.

9. Let  $f \in C_c^2(\mathbb{R}^3)$  and let

$$\Phi(\mathbf{x}) = \frac{1}{4\pi|\mathbf{x}|}$$

be the fundamental solution of Poisson's equation in  $\mathbb{R}^3$ .

(a) Show that for all  $\mathbf{x} \in \mathbb{R}^3$

$$\lim_{\varepsilon \rightarrow 0} \int_{B_\varepsilon(\mathbf{0})} \Phi(\mathbf{y}) \Delta_{\mathbf{y}} f(\mathbf{x} - \mathbf{y}) d\mathbf{y} = 0.$$

Hint: You may use the following, which you do not need to prove:

$$\|\Phi\|_{L^1(B_\varepsilon(\mathbf{0}))} = \frac{1}{2}\varepsilon^2.$$

(b) Show that for all  $\mathbf{x} \in \mathbb{R}^3$

$$\lim_{\varepsilon \rightarrow 0} \int_{\partial B_\varepsilon(\mathbf{0})} (\nabla_{\mathbf{y}} f(\mathbf{x} - \mathbf{y})) \cdot \mathbf{n}(\mathbf{y}) \Phi(\mathbf{y}) dS(\mathbf{y}) = 0$$

where  $\mathbf{n}$  is the outward-pointing unit normal vector field to  $\partial B_\varepsilon(\mathbf{0})$ .

(c) Let  $\mathbf{y} \in \partial B_\varepsilon(\mathbf{0})$ . Show that

$$\nabla \Phi(\mathbf{y}) \cdot \mathbf{n}(\mathbf{y}) = -\frac{1}{4\pi\varepsilon^2}.$$

(d) Use part (c) to show that for all  $\mathbf{x} \in \mathbb{R}^3$

$$\lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^3 \setminus B_\varepsilon(\mathbf{0})} \nabla \Phi(\mathbf{y}) \cdot \nabla_{\mathbf{y}} f(\mathbf{x} - \mathbf{y}) d\mathbf{y} = f(\mathbf{x}).$$

Hint: You may also use the following, which you do not need to prove:

$$\Delta \Phi(\mathbf{x}) = 0 \text{ for } \mathbf{x} \neq \mathbf{0}.$$

(e) Define  $u : \mathbb{R}^3 \rightarrow \mathbb{R}$  by

$$u(\mathbf{x}) = \int_{\mathbb{R}^3} \Phi(\mathbf{y}) f(\mathbf{x} - \mathbf{y}) d\mathbf{y}.$$

Use parts (a), (b) and (d) to show that for all  $\mathbf{x} \in \mathbb{R}^3$

$$-\Delta u(\mathbf{x}) = f(\mathbf{x}).$$

Hint: You may also use the following, which you do not need to prove:

$$\Delta u(\mathbf{x}) = \int_{\mathbb{R}^3} \Phi(\mathbf{y}) \Delta_{\mathbf{y}} f(\mathbf{x} - \mathbf{y}) d\mathbf{y}.$$

10. (a) Let  $u : [0, 1] \times [0, \infty) \rightarrow \mathbb{R}$  be a smooth solution of

$$\begin{aligned} u_t(x, t) - u_{xx}(x, t) &= 0 \quad \text{for } (x, t) \in (0, 1) \times (0, \infty), \\ u(x, 0) &= x(1 - x) \quad \text{for } x \in (0, 1), \\ u(0, t) &= u(1, t) = 0 \quad \text{for } t \in [0, \infty). \end{aligned}$$

(a1) Define  $v(x, t) = x(1 - x)e^{-\beta t}$ . Find the largest value of  $\beta_0 > 0$  such that for all  $\beta \in (0, \beta_0]$

$$v_t - v_{xx} \geq 0 \quad \forall (x, t) \in (0, 1) \times (0, \infty).$$

(a2) For all  $\beta \in (0, \beta_0]$ , prove that

$$0 \leq u(x, t) \leq x(1 - x)e^{-\beta t} \quad \forall (x, t) \in (0, 1) \times (0, \infty).$$

(b) Let  $\Omega \subset \mathbb{R}^n$  be open and bounded with smooth boundary and  $f : \Omega \rightarrow \mathbb{R}$  be smooth. Consider the semilinear elliptic PDE

$$\begin{aligned} -\Delta u + c(u) &= f \quad \text{in } \Omega, \\ u &= 0 \quad \text{on } \partial\Omega, \end{aligned} \tag{4}$$

where  $c : \mathbb{R} \rightarrow \mathbb{R}$  is a smooth, non-decreasing function, i.e.,  $c(t) \geq c(s)$  if  $t \geq s$ . Prove that (4) has at most one smooth solution.