



## EXAMINATION PAPER

<b>Examination Session:</b> May	<b>Year:</b> 2018	<b>Exam Code:</b> MATH3371-WE01
------------------------------------	----------------------	------------------------------------

<b>Title:</b> Representation Theory III
--

Time Allowed:	3 hours	
Additional Material provided:	None	
Materials Permitted:	None	
Calculators Permitted:	No	Models Permitted: Use of electronic calculators is forbidden.
Visiting Students may use dictionaries: No		

Instructions to Candidates:	Credit will be given for: the best <b>FOUR</b> answers from Section A and the best <b>THREE</b> answers from Section B. Questions in Section B carry <b>TWICE</b> as many marks as those in Section A.	
		<b>Revision:</b>

## SECTION A

1. Let  $G$  be a finite group.
  - (a) Define the regular representation  $(\pi, V)$  of  $G$  and state the theorem about the decomposition of  $V$  into irreducible representations. What is  $\dim \operatorname{Hom}_G(V, V)$ , the dimension of the space of  $G$ -intertwiners of  $V$  with itself?
  - (b) Assume  $G$  is abelian and fix  $g \in G$ . Show that  $\pi(g)$  is a  $G$ -intertwiner.
  - (c) Assume  $G = \mathbb{Z}/3\mathbb{Z}$ . Write down explicitly the matrix  $\pi(g)$  for every  $g \in G$  and explain why these give a basis for  $\operatorname{Hom}_G(V, V)$ .

2. (a) Let  $(\rho, V)$  be a representation of a finite group  $G$ . Define the dual representation  $(\rho^*, V^*)$ . What is the dual representation in terms of matrices? Show that

$$\chi_{V^*}(g) = \overline{\chi_V(g)}.$$

- (b) Explain why every finite-dimensional representation of the dihedral group  $D_n$  is equivalent to its dual representation.
3. (a) Give the character table of the symmetric group  $S_3$ . (No justification required).
  - (b) Let  $f$  be the class function on  $S_3$  defined by  $f(C) = \#C$  for a conjugacy class  $C$  in  $S_3$ . Express  $f$  in terms of the irreducible characters of  $S_3$ . Does  $f$  arise as the character of a representation of  $S_3$ ? Justify your answer.

4. Let  $G$  be a linear Lie group and  $\mathfrak{g}$  be its Lie algebra.

- (a) Give explicit formulas for the adjoint representations  $\operatorname{ad}$  and  $\operatorname{Ad}$ .
  - (b) Show that  $\operatorname{ad}$  is the derived representation of  $\operatorname{Ad}$ .
  - (c) Show directly that  $\operatorname{ad}$  is a Lie algebra homomorphism.

5. Let  $G = \operatorname{SL}_2(\mathbb{C})$  and consider the action of  $G$  on the space of smooth functions on row vectors  $\mathbf{x} \in \mathbb{C}^2$  given by

$$(\pi(g)\varphi)(\mathbf{x}) = \varphi(\mathbf{x}g).$$

- (a) Show that  $\pi$  defines a representation, i.e.,  $\pi(gh) = \pi(g)\pi(h)$  for  $g, h \in G$ .
  - (b) Compute the associated derived Lie algebra action  $D\pi$  for  $X = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \in \mathfrak{sl}_2(\mathbb{C})$ .

6. *There is no question 6 on this paper.*

## SECTION B

7. For  $n \geq 2$ , we let  $\text{Dic}_n$  be the so-called dicyclic group. This is by definition the group generated by two elements  $a$  and  $x$  subject to the relations

$$a^{2n} = e, \quad x^2 = a^n, \quad x^{-1}ax = a^{-1}.$$

You may assume that  $\text{Dic}_n$  is a non-abelian group of order  $4n$  and that every element can be written uniquely in the form  $a^i x^j$  with  $0 \leq i < 2n$  and  $j = 0, 1$ .

- Show that every irreducible representation of  $\text{Dic}_n$  has degree at most 2.
  - Use the ‘method of the abelian subgroup’ to explicitly construct all (equivalence classes of) irreducible representations  $(\rho, V)$  of  $\text{Dic}_n$ . (For convenience handle the cases  $\dim V = 1$  and  $\dim V = 2$  separately). Write down the matrices  $\rho(x)$  and  $\rho(a)$  explicitly. Check that your list is complete.
  - For  $n = 3$ , write down the character table of  $\text{Dic}_n$ .  
(Hint: we have  $\omega + \omega^{-1} = 1$ ;  $\omega^2 + \omega^{-2} = -1$  for  $\omega = e^{\pi i/3}$ . You may initially assume that  $a, a^2, a^3, x, ax$  lie in different conjugacy classes. After completing the table explain why this is indeed the case. Then find the order of these classes using the table.)
8. For an *odd* prime  $p$ , we consider the finite field  $\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z}$ . Recall all its (additive) characters are given by  $\psi_\ell(x) := e^{2\pi i x \ell/p}$  with  $\ell \in \mathbb{F}_p$ . (Here we identify elements in  $\mathbb{F}_p$  with their preimage in  $\mathbb{Z}$ ).

Recall that the Heisenberg group  $H$  over  $\mathbb{F}_p$  is given as the set of triples  $(x, y, z) \in \mathbb{F}_p^3$  with the multiplication defined by

$$(x, y, z) \cdot (x', y', z') = (x + x', y + y', z + z' + xy' - x'y).$$

For example, we have  $(-x, 0, 0) \cdot (0, y, z) \cdot (x, 0, 0) = (0, y, z - 2xy)$  and that the center of  $H$  is  $\{(0, 0, z); z \in \mathbb{F}_p\}$ .

- State Mackey’s irreducibility criterion for induced representations from a *normal* subgroup.
- Let  $K = \{(0, y, z); y, z \in \mathbb{F}_p\}$ , which you may assume is a normal abelian subgroup of  $H$  of index  $p$  with left coset representatives  $\{(x, 0, 0)\}$ . Consider for  $\ell \neq 0$  its one-dimensional representation  $\psi'_\ell$  given by  $\psi'_\ell((0, y, z)) := \psi_\ell(z)$ . Use (a) to show that the induced representation  $\text{Ind}_K^H \psi'_\ell$  is irreducible.
- Let  $(\rho, V)$  be an arbitrary representation of  $H$ . Assume that the restriction  $\text{Res}_K V$  to  $K$  contains the representation  $\psi'_\ell$ . Show  $\langle \text{Ind}_K^H \psi'_\ell, V \rangle_H \geq 1$ .
- Now assume that  $(\rho, V)$  is irreducible with central character  $\psi_\ell$ , i.e.,  $\rho(0, 0, z)v = \psi_\ell(z)v$  for all  $v \in V$ . Note that  $V$  must contain an eigenvector  $w$  under the action of  $K$ : More precisely, (you may assume) there exists a  $k \in \mathbb{F}_p$  such that

$$\rho(0, y, z)w = \psi_k(y)\psi_\ell(z)w \quad \text{for all } (0, y, z) \in K.$$

Compute the action of  $(0, y, z)$  on  $w_x := \rho(x, 0, 0)w$  and show that there exists an  $x \in \mathbb{F}_p$  such that

$$\rho(0, y, z)w_x = \psi_\ell(z)w_x \quad \text{for all } (0, y, z) \in K.$$

In particular,  $\text{Res}_K V$  contains  $\psi'_\ell$ . Use (c) to conclude  $\text{Ind}_K^H \psi'_\ell \simeq V$ .

9. We define elements of  $\mathfrak{sl}_2(\mathbb{C})$ ,

$$\tilde{H} := \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad X_+ := \frac{1}{2} \begin{pmatrix} 1 & i \\ i & -1 \end{pmatrix}, \quad X_- := \frac{1}{2} \begin{pmatrix} 1 & -i \\ -i & -1 \end{pmatrix}.$$

- (a) Show that  $\tilde{H}$ ,  $X_+$ , and  $X_-$  satisfy exactly the same bracket relations as the standard basis  $\{H, X, Y\}$  of  $\mathfrak{sl}_2(\mathbb{C})$ .

Show by an explicit calculation that if in a representation  $(\pi, V)$  of  $\mathfrak{sl}_2(\mathbb{C})$  a vector  $v$  is an eigenvector for  $\tilde{H}$  (“ $\tilde{H}$ -weight vector”) with eigenvalue (“ $\tilde{H}$ -weight”)  $\lambda$ , then  $\pi(X_{\pm})v$  is either zero or also an  $\tilde{H}$ -weight vector with  $\tilde{H}$ -weight  $\lambda \pm 2$ .

- (b) Let  $\mathcal{F} := \mathbb{C}[z]$  be the (infinite-dimensional)  $\mathbb{C}$ -vector space of polynomials in one variable. We define operators on  $\mathcal{F}$  by

$$\omega(\tilde{H}) := z \frac{d}{dz} + \frac{1}{2}, \quad \omega(X_+) := \frac{1}{2} z^2, \quad \omega(X_-) := -\frac{1}{2} \frac{d^2}{dz^2}.$$

(So e.g.,  $\omega(\tilde{H})p(z) = zp'(z) + \frac{1}{2}p(z)$ .) Show that these operators preserve the bracket relations and hence define a Lie algebra representation  $\omega$  of  $\mathfrak{sl}_2(\mathbb{C})$ . (The operator identities  $\frac{d}{dz}z = 1 + z\frac{d}{dz}$ ,  $\frac{d^2}{dz^2}z^2 = 2 + 4z\frac{d}{dz} + z^2\frac{d^2}{dz^2}$  should help).

- (c) Find two linear independent lowest weight vectors ( $\omega(X_-)p = 0$ ) in  $\mathcal{F}$  and their  $\tilde{H}$ -weights. Show that these vectors each generate an (infinite dimensional) subrepresentation, say  $\mathcal{F}_1$  and  $\mathcal{F}_2$  respectively, such that  $\mathcal{F} = \mathcal{F}_1 \oplus \mathcal{F}_2$ .

In particular, describe  $\mathcal{F}_1$  and  $\mathcal{F}_2$  and give their  $\tilde{H}$ -weight structure and the action of  $X_{\pm}$ .

Finally, show that  $\mathcal{F}_1$  and  $\mathcal{F}_2$  are irreducible.

10. *There is no question 10 on this paper.*