



## EXAMINATION PAPER

<b>Examination Session:</b> May	<b>Year:</b> 2019	<b>Exam Code:</b> MATH3201-WE01
------------------------------------	----------------------	------------------------------------

<b>Title:</b> Geometry III
-------------------------------

Time Allowed:	3 hours	
Additional Material provided:	Formula Sheet	
Materials Permitted:	None	
Calculators Permitted:	No	Models Permitted: Use of electronic calculators is forbidden.
Visiting Students may use dictionaries: No		

Instructions to Candidates:	Credit will be given for: the best <b>FOUR</b> answers from Section A and the best <b>THREE</b> answers from Section B. Questions in Section B carry <b>TWICE</b> as many marks as those in Section A.	
		<b>Revision:</b>

## SECTION A

1. (a) Let  $G$  be a group acting on a set  $X$ . Give the definition of the orbit of an element  $x \in X$  under the action of  $G$ .  
 (b) Consider the Euclidean plane  $\mathbb{E}^2$  represented by complex numbers. Let  $G$  be a group acting on  $\mathbb{E}^2$  and generated by elements  $g(z) = z + 1$  and  $h(z) = iz$ . Find the orbit of the point  $1 + i$  under the action of the group  $G$ .
2. (a) Is it true that affine transformations act transitively on quadrilaterals in  $\mathbb{E}^2$ ? Justify your answer.  
 (b) Let  $A_1A_2A_3$  be a triangle in  $\mathbb{E}^2$ . Denote  $A_4 = A_1$ ,  $A_5 = A_2$ . Let  $B_i$ ,  $i = 1, 2, 3$ , be a point on the line  $A_iA_{i+1}$  such that  $|B_iA_i| = \frac{1}{2}|A_iA_{i+1}|$  and  $A_i$  lies between  $B_i$  and  $A_{i+1}$ . Similarly, let  $C_i$ ,  $i = 1, 2, 3$ , be a point on the line  $A_iA_{i+2}$  such that  $|C_iA_i| = \frac{1}{2}|A_iA_{i+2}|$  and  $A_i$  lies between  $C_i$  and  $A_{i+2}$ .  
 Show that the points  $K = B_1B_2 \cap C_1C_3$ ,  $L = B_2B_3 \cap C_2C_3$ ,  $M = B_1B_3 \cap C_1C_2$  are collinear.
3. (a) Let  $ABC$  be a triangle in  $\mathbb{E}^2$ , and let  $M$  and  $N$  be the midpoints of  $AB$  and  $AC$  respectively. Show that  $|MN| = \frac{1}{2}|BC|$ .  
 (b) Let  $ABC$  be a triangle in  $S^2$ , and let  $M$  and  $N$  be the midpoints of  $AB$  and  $AC$  respectively. Show that  $|MN| > \frac{1}{2}|BC|$ .
4. (a) Let  $\mathcal{C}_1, \mathcal{C}_2, \mathcal{C}_3$  be three mutually tangent circles. Is it always true that there exists a fourth circle  $\mathcal{C}_4$  tangent to all three of  $\mathcal{C}_1, \mathcal{C}_2, \mathcal{C}_3$ ? Justify your answer.  
 (b) Show that any four mutually tangent circles  $\mathcal{C}_1, \mathcal{C}_2, \mathcal{C}_3, \mathcal{C}_4$  with  $\mathcal{C}_1 \cap \mathcal{C}_2 \cap \mathcal{C}_3 \cap \mathcal{C}_4 = \emptyset$  can be taken by a Möbius transformation to some three mutually tangent unit circles inscribed into another circle.
5. Let  $XYZ$  be an ideal triangle in  $\mathbb{H}^2$ .  
 (a) Show that  $\triangle XYZ$  has an inscribed circle.  
 (b) Find the hyperbolic cosine of the radius of the circle inscribed into  $\triangle XYZ$ .
6. (a) Define the angle of parallelism in  $\mathbb{H}^2$ .  
 (b) Let  $m$  and  $n$  be two orthogonal lines in  $\mathbb{H}^2$ , denote  $O = m \cap n$ . Let  $l_1$  and  $l_3$  be the two distinct lines orthogonal to  $m$  and intersecting  $m$  at distance  $a$  from  $O$ . Let  $l_2$  and  $l_4$  be the two distinct lines orthogonal to  $n$  and crossing  $n$  at distance  $x$  from  $m$ .  
 Given  $a$ , for which values of  $x$  do the lines  $l_1, l_2, l_3, l_4$  compose a quadrilateral having finite area?

## SECTION B

7. Let  $ABC$  be a triangle in  $\mathbb{H}^2$  (labelled clockwise) with angles  $\alpha, \beta, \gamma$  at  $A, B, C$  respectively. Denote by  $R_{X,\varphi}$  a rotation around point  $X$  through angle  $\varphi$  in the clockwise direction.
- Let  $g = R_{A,2\alpha} \circ R_{B,2\beta}$ . Find all the fixed points of  $g$ .
  - Find  $h = R_{A,2\alpha} \circ R_{B,2\beta} \circ R_{C,2\gamma}$ . Does it have fixed points?
  - Now, consider  $\varphi = R_{A,\alpha} \circ R_{B,\beta} \circ R_{C,\gamma}$ . Show that  $\varphi$  takes the line  $AC$  to itself.
  - How many fixed points has the isometry  $\varphi$  introduced in part (c)? Find the order of  $\varphi$ .
8. Let  $A, B, C$  be points on the unit sphere  $S^2$ . Suppose that  $B \in \text{Pol}(A)$  and  $C \in \text{Pol}(B)$ .
- Find  $\angle CAB$ .
  - Suppose that  $\angle CBA = \beta$ . Find the length  $AC$ .
  - Let  $\triangle A'B'C'$  be a triangle polar to  $\triangle ABC$ . Given that  $\angle CBA = \beta < \pi/2$ , which of the triangles  $\triangle ABC$  and  $\triangle A'B'C'$  has larger area?
  - Let  $l \subset S^2$  be a line,  $D_1, D_2 \in \text{Pol}(l)$  be the two distinct poles to  $l$ . Let  $\varepsilon > 0$ , and let  $P_1, P_2$  be two points such that  $d(P_i, l) < \varepsilon$  for  $i = 1, 2$ . Given a point  $A \in \text{Pol}(P_1 P_2)$ , is it true that for at least one of  $D_i$  we have  $d(A, D_i) < \varepsilon$ ?
9. (a) Which of the following statements are true? Justify your answer.
- Projective transformations of  $\mathbb{R}P^2$  act transitively on pairs of projective lines.
  - Projective transformations of  $\mathbb{R}P^2$  act transitively on triples of projective lines.
- The points  $A_1, A_2, A_3, A_4$  lie on a line  $a$  in the Euclidean plane  $\mathbb{E}^2$ , and the points  $B_1, B_2, B_3, B_4$  lie on a line  $b \subset \mathbb{E}^2$ , where  $a$  is not parallel to  $b$ . Assume that all the four lines  $A_i B_i$ ,  $i = 1, 2, 3, 4$  intersect at one point, denote the intersection point by  $P$ . Let  $Q = A_1 B_2 \cap B_1 A_2$  and  $S = A_3 B_4 \cap A_4 B_3$ . Show that the point  $a \cap b$  lies on the line  $QS$ .
  - Assuming that  $P = (1, 0)$ ,  $B_1 = (0, 0)$ ,  $B_2 = (0, 1)$ ,  $B_3 = (0, 2)$ ,  $B_4 = (0, 3)$ , find the cross-ratio of the lines  $PB_1, PB_2, PB_3, PB_4$ .
  - Formulate the statement dual to the one in part (b).

10. The goal of this problem is to prove that if opposite angles of a hyperbolic quadrilateral are equal then there exists a rotation by  $\pi$  taking the quadrilateral to itself, and hence its opposite sides are also equal.
- (a) Let  $l$  be a line in  $\mathbb{H}^2$ , let  $P_1, P_2 \in l$  be points and let  $X \in l \cap \partial\mathbb{H}^2$  be one of the endpoints of  $l$ . Let  $S_1, S_2$  be points lying in one half-plane with respect to  $l$  such that  $\angle S_1 P_1 X = \angle S_2 P_2 X$ . Show that the lines  $S_1 P_1$  and  $S_2 P_2$  do not intersect.
  - (b) In the assumptions of (a), assume that  $P_1$  lies between  $X$  and  $P_2$  on  $l$ . Assume that the line  $m$  through points  $S_1$  and  $S_2$  is ultra-parallel to  $l$ , denote by  $Y$  the endpoint of  $m$  such that  $S_1$  lies between  $Y$  and  $S_2$ . Show that the angle  $\angle Y S_1 P_1$  is larger than the angle  $\angle Y S_2 P_2$ .
  - (c) Let  $ABCD$  be a quadrilateral in  $\mathbb{H}^2$ . Suppose that  $\angle A = \angle C = \alpha$  and  $\angle B = \angle D = \beta$ . Show that the lines  $AB$  and  $CD$  are ultra-parallel.
  - (d) In the assumptions of part (c), show that there exists a point  $O$  such that a rotation  $R_{O,\pi}$  around  $O$  through  $\pi$  takes the quadrilateral  $ABCD$  to itself.

## Formula sheet

Sine and cosine laws:

	sine law	cosine laws
$S^2$	$\frac{\sin a}{\sin \alpha} = \frac{\sin b}{\sin \beta} = \frac{\sin c}{\sin \gamma}$	$\cos a = \cos b \cos c + \sin b \sin c \cos \alpha$ $\cos \alpha = -\cos \beta \cos \gamma + \sin \beta \sin \gamma \cos a$
$\mathbb{E}^2$	$\frac{a}{\sin \alpha} = \frac{b}{\sin \beta} = \frac{c}{\sin \gamma}$	$a^2 = b^2 + c^2 - 2bc \cos \alpha$
$\mathbb{H}^2$	$\frac{\sinh a}{\sin \alpha} = \frac{\sinh b}{\sin \beta} = \frac{\sinh c}{\sin \gamma}$	$\cosh a = \cosh b \cosh c - \sinh b \sinh c \cos \alpha$ $\cos \alpha = -\cos \beta \cos \gamma + \sin \beta \sin \gamma \cosh a$

Circles:

	$S^2$	$\mathbb{E}^2$	$\mathbb{H}^2$
Circumference of a circle	$2\pi \sin R$	$2\pi R$	$2\pi \sinh R$
Area of a disc	$4\pi \sin^2(\frac{R}{2})$	$\pi R^2$	$4\pi \sinh^2(\frac{R}{2})$

Angle of parallelism in hyperbolic geometry:

For a point on distance  $a$  from the line, the angle of parallelism  $\varphi$  satisfies

$$\sin \varphi = \frac{1}{\cosh a}$$

Distance formula in the upper half-plane model of hyperbolic geometry:

$$\cosh d(u, v) = 1 + \frac{|u - v|^2}{2\operatorname{Im}(u)\operatorname{Im}(v)}$$

Distance formula in the hyperboloid model of hyperbolic geometry:

For  $u, v \in \mathbb{R}^{2,1}$ , let  $Q = |\frac{(u,v)^2}{(u,u)(v,v)}|$ . Then

if  $(u, u) < 0, (v, v) < 0$  then  $Q = \cosh^2 d(pt, pt)$

if  $(u, u) < 0, (v, v) > 0$  then  $Q = \sinh^2 d(pt, line)$

if  $(u, u) > 0, (v, v) > 0$  then  $Q < 1 \Rightarrow$  intersecting lines,  $Q = \cos^2 \alpha$ ;  
 $Q = 1 \Rightarrow$  parallel lines;  
 $Q > 1 \Rightarrow$  ultraparallel lines,  $Q = \cosh^2 d(line, line)$