



EXAMINATION PAPER

Examination Session: May/June	Year: 2025	Exam Code: MATH4171-WE01
---	----------------------	------------------------------------

Title: Riemannian Geometry IV

Time:	3 hours	
Additional Material provided:		
Materials Permitted:		
Calculators Permitted:	No	Models Permitted: Use of electronic calculators is forbidden.

Instructions to Candidates:	<p>Answer all questions.</p> <p>Section A is worth 40% and Section B is worth 60%. Within each section, all questions carry equal marks.</p> <p>Write your answer in the white-covered answer booklet with barcodes.</p> <p>Begin your answer to each question on a new page.</p>
-----------------------------	---

Revision:	
------------------	--

SECTION A

Q1 Prove or disprove the following assertions:

(a) *The subset of \mathbb{R}^3 given by*

$$M = \{(x, y, z) \in \mathbb{R}^3 \mid z = 0\} \cup \{(x, y, z) \in \mathbb{R}^3 \mid x = y = 0\}$$

is a smooth submanifold of \mathbb{R}^3 .

(b) *The subset of \mathbb{R}^3 given by*

$$M = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + 2y^2 + 7z^2 = 1\}$$

is a smooth 2-dimensional submanifold of \mathbb{R}^3 .

Q2 (a) State the definition of a differentiable vector field on a smooth manifold M .

(b) Let

$$X = \frac{\partial}{\partial x} \text{ and } Y = x \frac{\partial}{\partial y}$$

be two differentiable vector fields on \mathbb{R}^2 equipped with the usual chart given by the identity map. Show that XY is not a linear derivation on $C^\infty(\mathbb{R}^2)$ by applying it to the product of the smooth real-valued functions f, g on \mathbb{R}^2 given by $f(x, y) = x$ and $g(x, y) = y$.

(c) Compute the Lie bracket of the vector fields X and Y .

Q3 Prove or disprove the following assertions:

(a) *Let*

$$P = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1 \text{ and } (x, y, z) \neq (0, 0, 1)\} \subset \mathbb{R}^3.$$

Then the Riemannian metric induced on P by the Euclidean metric on \mathbb{R}^3 is geodesically complete.

(b) *There exists a Riemannian metric g on \mathbb{R}^2 such that the Christoffel symbols $\Gamma_{ij}^k(p)$ of its Levi-Civita connection ∇ in local coordinates at some point p are given by*

$$\begin{aligned} \Gamma_{11}^1(p) &= 0, & \Gamma_{11}^2(p) &= 0, \\ \Gamma_{12}^1(p) &= 1, & \Gamma_{12}^2(p) &= 0, \\ \Gamma_{21}^1(p) &= -1, & \Gamma_{21}^2(p) &= 0, \\ \Gamma_{22}^1(p) &= 1, & \Gamma_{22}^2(p) &= 0. \end{aligned}$$

(c) *Every complete Riemannian manifold with positive sectional curvature is compact.*

Q4 Let $P = \{(x, y, z) \mid z = x^2 + y^2\} \subset \mathbb{R}^3$ be a paraboloid and let $\varphi: P \rightarrow \mathbb{R}^2$ be a global chart given by

$$\varphi^{-1}(x, y) = (x, y, x^2 + y^2).$$

Consider the smooth map $f: \mathbb{R}^2 \rightarrow P$ given by

$$f(u, v) = (u, 3v, u^2 + 9v^2).$$

- (a) Compute the matrix of the differential $Df_{(u_o, v_o)}$ at a point $(u_o, v_o) \in \mathbb{R}^2$ in terms of the bases of $T_{(u_o, v_o)}\mathbb{R}^2$ and $T_{f(u_o, v_o)}P$ determined by the coordinate tangent vectors.
- (b) Find a Riemannian metric g on P such that f becomes an isometry with respect to the Euclidean metric on \mathbb{R}^2 and compute its matrix expression (g_{ij}) in the coordinates (x, y) . You may assume, without proof, that f is a diffeomorphism.

SECTION B

Q5 (a) Let $\varphi^{-1}(x, y) = (f(x)\cos(y), f(x)\sin(y), h(x))$ with $f: [a, b] \rightarrow (0, \infty)$ be a coordinate chart of a smooth surface of revolution $S \subset \mathbb{R}^3$. Assume that S carries the Riemannian metric g induced by the Euclidean metric on \mathbb{R}^3 . Assume, furthermore, that

$$(f'(x))^2 + (h'(x))^2 = 1 \quad \text{for all } x \in [a, b].$$

Compute the local expression (g_{ij}) of the Riemannian metric g in the coordinates given by the chart φ and show that

$$\nabla_{\frac{\partial}{\partial x}} \frac{\partial}{\partial x} = 0,$$

$$\nabla_{\frac{\partial}{\partial x}} \frac{\partial}{\partial y} = \nabla_{\frac{\partial}{\partial y}} \frac{\partial}{\partial x} = \frac{f'(x)}{f(x)} \frac{\partial}{\partial y},$$

$$\nabla_{\frac{\partial}{\partial y}} \frac{\partial}{\partial y} = -f(x)f'(x) \frac{\partial}{\partial x}.$$

- (b) Prove or disprove the following statement: *Any two tori T^m and T^n with $m \neq n$ are locally diffeomorphic.*

- Q6** Fix $n \geq 2$ and let $M(n, \mathbb{R})$ denote the set of $n \times n$ matrices with real entries, $\text{Sym}(n) = \{A \in M(n, \mathbb{R}) \mid A^T = A\}$ be the set of $n \times n$ real symmetric matrices, and

$$J = \begin{pmatrix} \text{Id}_{n-1} & 0 \\ 0 & -1 \end{pmatrix},$$

where Id_k denotes the $k \times k$ identity matrix. Let

$$\text{O}(n-1, 1) = \{A \in M(n, \mathbb{R}) \mid AJA^T = J\}.$$

- (a) Show that each matrix in $\text{O}(n-1, 1)$ is invertible and that $\text{O}(n-1, 1)$ is a group under matrix multiplication.
- (b) Define $f: M(n, \mathbb{R}) \rightarrow \text{Sym}(n)$ by $f(A) = AJA^T$. Show that

$$Df_A(B) = AJB^T + BJA^T$$

for any $A, B \in M(n, \mathbb{R})$ and conclude that, for $A \in \text{O}(n-1, 1)$ and $C \in \text{Sym}(n)$,

$$Df_A\left(\frac{1}{2}CJA\right) = C.$$

- (c) Show that $J \in \text{Sym}(n)$ is a regular value of f .
- (d) Show that $\text{O}(n-1, 1)$ is a Lie group and determine its dimension.
- Q7** (a) Let $M = S^2(1) \times S^2(1)$ be the Riemannian product of two copies of the 2-dimensional unit round sphere. Show that M has non-negative sectional curvature. You may use, without proof, results on product Riemannian manifolds seen in the lecture.
- (b) Does there exist a geodesic $\gamma: [a, b] \rightarrow \mathbb{H}^n$ in n -dimensional hyperbolic space and a non-zero Jacobi field J along γ with $J(a) = J(b) = 0$? Justify your answer. You may use, without proof, results on Jacobi fields and curvature seen in the lecture.
- (c) Let M be a smooth manifold and let ∇ be an affine connection on M . The *conjugate connection* ∇^* is defined by

$$\nabla_X^* Y = \nabla_X Y - T(X, Y),$$

where $T(X, Y) = \nabla_X Y - \nabla_Y X - [X, Y]$ is the torsion of ∇ . Show that ∇^* is an affine connection on M and that the torsion of ∇^* is $-T$.

- Q8** (a) Let (M, g) be a Riemannian manifold and $c: [a, b] \rightarrow M$ be a geodesic. Let X be a smooth parallel vector field along c . Show that the map $t \mapsto \langle X(t), c'(t) \rangle$ is constant.
- (b) State the symmetries of the curvature tensor R of a Riemannian manifold (M, g) .
- (c) Let (M, g) be an n -dimensional Riemannian manifold and let R be its Riemannian curvature tensor. The *Ricci tensor* of M is defined as

$$\text{Ric}(X, Y) = \text{Tr}(Z \mapsto R(Z, X)Y),$$

where X, Y, Z are smooth vector fields on M , i.e. Ric is the trace of the endomorphism $Z \mapsto R(Z, X)Y$. In terms of an orthonormal basis $\{e_i\}_{i=1}^n$ of the tangent space, Ric is given by

$$\text{Ric}(X, Y) = \sum_{i=1}^n g(R(e_i, X)Y, e_i).$$

Show that Ric defines a $\binom{2}{0}$ symmetric tensor field on M .