



EXAMINATION PAPER

Examination Session: May/June	Year: 2026	Exam Code: MATH2727-WE01
-----------------------------------------	----------------------	------------------------------------

Title: Topology II

Time:	2 hours	
Additional Material provided:	None	
Materials Permitted:	None	
Calculators Permitted:	No	Models Permitted: Use of electronic calculators is forbidden.

Instructions to Candidates:	<p>Answer all questions.</p> <p>The indicative marks shown in brackets for the main parts of each question are given as a guide to the weighting the markers expect to apply.</p> <p>Write your answer in the white-covered answer booklet with barcodes.</p> <p>Begin your answer to each question on a new page.</p>
-----------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Revision:	
------------------	--

SECTION A

1. Let $X = \{1, 2, 3, 4\}$.

(a) Consider the following topologies on X .

$$\tau_a = \{\emptyset, \{1\}, \{1, 2\}, X\} \quad \tau_b = \{\emptyset, \{1, 2\}, \{3, 4\}, X\}$$

$$\tau_c = \{\emptyset, \{1\}, \{1, 2, 3\}, X\} \quad \tau_d = \{\emptyset, \{2\}, \{2, 4\}, X\}$$

Which of the topological spaces (X, τ_a) , (X, τ_b) , (X, τ_c) and (X, τ_d) are homeomorphic? If a pair are homeomorphic, give a homeomorphism; if they are not, explain why not. [3]

For any set Y and finite collection of subsets $\tau_Y = \{\emptyset, U_1, U_2, \dots, U_k, Y\}$, we say τ_Y is *nested* if $\emptyset \subset U_1 \subset U_2 \subset \dots \subset U_k \subset Y$, where $k \geq 1$. (Note that these inclusions are strict.)

(b) Give two topologies on X , one nested and one not, each containing 5 sets. [2]

(c) Show in general that if a collection τ_Y is nested then it is a topology on Y . [2]

(d) What does it mean for a topological space to be Hausdorff? If τ_Y is nested then is (Y, τ_Y) always, sometimes, or never a Hausdorff space? Explain. [3]

2. In \mathbb{R}^n with the standard topology, we define the closed ball and the sphere:

$$D^n = \{(x_1, \dots, x_n) \mid x_1^2 + \dots + x_n^2 \leq 1\}, \quad S^{n-1} = \{(x_1, \dots, x_n) \mid x_1^2 + \dots + x_n^2 = 1\}.$$

(a) Show that both sets are compact. If you use a result from lectures, state it clearly. [4]

In a lecture, we proved that for $n \geq 1$ the quotient space

$$D^n/S^{n-1} \cong S^n \subseteq \mathbb{R}^{n+1}.$$

Prove this for $n = 1$, as follows:

(b) Draw simple sketches of the three sets involved. Give explicitly a suitable continuous, surjective function $f : D^1 \rightarrow S^1$, and define a function $\bar{f} : D^1/S^0 \rightarrow S^1$ in terms of f . [3]

(c) Show that \bar{f} is a homeomorphism. Explain clearly, stating any results from lectures that you use, but detailed calculations are not required. [3]

SECTION B

3. In this question \mathbb{R} always has the standard topology.
Recall that, for $x \in \mathbb{R}$, $\lfloor x \rfloor$ is the largest integer $\leq x$, so $\lfloor 3.142 \rfloor = 3$. We define

$$f : \mathbb{R} \rightarrow \mathbb{R} \text{ by } f(x) = \lfloor x \rfloor,$$

$$g : \mathbb{R} \rightarrow \mathbb{R} \text{ by } g(x) = x - \lfloor x \rfloor.$$

- (a) Specify the image sets $F = f(\mathbb{R})$ and $G = g(\mathbb{R})$ in \mathbb{R} . [1]
- (b) Write down the definition of continuity for a map between topological spaces, and use it to show that neither f nor g is continuous. [3]
- (c) We now give each of F and G the subspace topology from \mathbb{R} . Describe these topologies τ_F and τ_G (that is, say which subsets of F and G are open) and briefly explain why. [3]
- (d) Next we construct the product space $F \times G$, and give it the product topology from τ_F and τ_G . Show that

$$h : \mathbb{R} \rightarrow F \times G \text{ given by } h(x) = (f(x), g(x))$$

is a bijection, and specify its inverse $h^{-1} : F \times G \rightarrow \mathbb{R}$. [3]

- (e) Show that h is not continuous. [2]
- (f) Explain why h^{-1} is continuous. [3]
-

4. (a) For a topological space X , write down the definitions of
i) connectedness, and ii) path-connectedness. [2]

- (b) Show that if X is path-connected, then it is connected. [3]

We now consider the set $M_2(\mathbb{R})$ of 2×2 matrices of real numbers, given the standard topology on \mathbb{R}^4 as usual.

- (c) Show that $\text{GL}_2(\mathbb{R}) = \{A \in M_2(\mathbb{R}) \mid \det(A) \neq 0\}$ is not connected. [2]

Let

$$S = \left\{ A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(\mathbb{R}) \mid \det(A) = 0; a, b, c, d \text{ all } \geq 0 \right\} \subseteq M_2(\mathbb{R}) \setminus \text{GL}_2(\mathbb{R})$$

$$\text{and } T = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in S \mid a, b, c, d \text{ all } > 0 \right\}.$$

- (d) Show explicitly, by giving a path, that T is path-connected. [3]

- (e) A typical matrix in $S \setminus T$ is $L = \begin{pmatrix} 0 & 0 \\ c & d \end{pmatrix}$ with $c, d > 0$.

Show that L is a limit point for T .

(Hint: Many different metrics induce the standard topology on \mathbb{R}^4 .) [3]

- (f) Given that *any* matrix in $S \setminus T$ is a limit point for T , show that S is connected. [2]
-