

Scaling limits of stochastic processes associated with resistance forms

Markov Processes, Mixing Times and Cutoff
London Mathematical Society EPSRC Durham Symposium
29 July 2017

David Croydon (University of Warwick)
partly joint with
Ben Hambly (Oxford) and Takashi Kumagai (Kyoto)


1. MOTIVATION

E.G. CRITICAL GALTON-WATSON TREES

Let T_n be a Galton-Watson tree with a critical (mean 1), aperiodic, finite variance offspring distribution, conditioned to have n vertices, then

$$n^{-1/2}T_n \rightarrow \mathcal{T},$$

where \mathcal{T} is (up to a constant) the **Brownian continuum random tree (CRT)** [Aldous 93], also [Duquesne/Le Gall 02].



Convergence in Gromov-Hausdorff-Prohorov topology implies

$$\left(n^{-1/2} X_{n^{3/2}t}^{T_n} \right) \rightarrow \left(X_t^{\mathcal{T}} \right)_{t \geq 0},$$


see [Krebs 95], [C. 08] and [Athreya/Löhr/Winter 14].

SOME INTUITION

Suppose T is a graph tree, and X^T is the discrete time simple random walk on T , $\pi(\{x\}) = \deg_T(x)$ its invariant measure. The following two properties are then easy to check:

- **[Scale]** For $x, y, z \in T$,

$$P_z^T(\sigma_x < \sigma_y) = \frac{d_T(b_T(x, y, z), y)}{d_T(x, y)}.$$



- **[Speed]** Expected number of visits to z when started at x and killed at y ,

$$d_T(b_T(x, y, z), y)\pi(\{z\}).$$

Analogous properties hold for limiting diffusion.

cf. One-dimensional convergence results of [Stone 63].

2. STOCHASTIC PROCESSES ASSOCIATED WITH RESISTANCE METRICS

RANDOM WALKS ON GRAPHS

Let $G = (V, E)$ be a finite, connected graph, equipped with (strictly positive, symmetric) edge conductances $(c(x, y))_{\{x, y\} \in E}$. Let μ be a finite measure on V (of full-support).

Let X be the continuous time Markov chain with generator Δ , as defined by:

$$(\Delta f)(x) := \frac{1}{\mu(\{x\})} \sum_{y: y \sim x} c(x, y)(f(y) - f(x)).$$

NB. Common choices for μ are:

- $\mu(\{x\}) := \sum_{y: y \sim x} c(x, y)$, the **constant speed random walk (CSRW)**;
- $\mu(\{x\}) := 1$, the **variable speed random walk (VSRW)**.

DIRICHLET FORM AND RESISTANCE METRIC

Define a quadratic form on G by setting

$$\mathcal{E}(f, g) = \frac{1}{2} \sum_{x, y: x \sim y} c(x, y) (f(x) - f(y)) (g(x) - g(y)).$$

Note that (regardless of the particular choice of μ ,) \mathcal{E} is a **Dirichlet form** on $L^2(\mu)$, and

$$\mathcal{E}(f, g) = - \sum_{x \in V} (\Delta f)(x) g(x) \mu(\{x\}).$$

Suppose we view G as an electrical network with edges assigned conductances according to $(c(x, y))_{\{x, y\} \in E}$. Then the **effective resistance** between x and y is given by

$$R(x, y)^{-1} = \inf \{ \mathcal{E}(f, f) : f(x) = 1, f(y) = 0 \}.$$

R is a metric on V , e.g. [Tetali 91], and characterises the weights (and therefore the Dirichlet form) uniquely [Kigami 95].

SUMMARY

RANDOM WALK X WITH GENERATOR Δ



DIRICHLET FORM \mathcal{E} on $L^2(\mu)$



RESISTANCE METRIC R AND MEASURE μ

RESISTANCE METRIC, e.g. [KIGAMI 01]


Let F be a set. A function $R : F \times F \rightarrow \mathbb{R}$ is a **resistance metric** if, for every finite $V \subseteq F$, one can find a weighted (i.e. equipped with conductances) graph with vertex set V for which $R|_{V \times V}$ is the associated effective resistance.

EXAMPLES

- Effective resistance metric on a graph;
- One-dimensional Euclidean (not true for higher dimensions);
- Any shortest path metric on a tree;
- Resistance metric on a Sierpinski gasket, where for 'vertices' of limiting fractal, we set

$$R(x, y) = (3/5)^n R_n(x, y),$$

then use continuity to extend to whole space.



RESISTANCE AND DIRICHLET FORMS

Theorem (e.g. [Kigami 01]) There is a one-to-one correspondence between resistance metrics and a class of quadratic forms called **resistance forms**.

The relationship between a resistance metric R and resistance form $(\mathcal{E}, \mathcal{F})$ is characterised by

$$R(x, y)^{-1} = \inf \{ \mathcal{E}(f, f) : f \in \mathcal{F}, f(x) = 1, f(y) = 0 \}.$$

Moreover, if (F, R) is compact, then $(\mathcal{E}, \mathcal{F})$ is a regular Dirichlet form on $L^2(\mu)$ for any finite Borel measure μ of full support. (Version of the statement also hold for locally compact spaces.)

**RESISTANCE FORM DEFINITION,
e.g. [KIGAMI 12]**

[RF1] \mathcal{F} is a linear subspace of the collection of functions $\{f : F \rightarrow \mathbb{R}\}$ containing constants, and \mathcal{E} is a non-negative symmetric quadratic form on \mathcal{F} such that $\mathcal{E}(f, f) = 0$ if and only if f is constant on F .

[RF2] Let \sim be the equivalence relation on \mathcal{F} defined by saying $f \sim g$ if and only if $f - g$ is constant on F . Then $(\mathcal{F}/\sim, \mathcal{E})$ is a Hilbert space.

[RF3] If $x \neq y$, then there exists an $f \in \mathcal{F}$ such that $f(x) \neq f(y)$.

[RF4] For any $x, y \in F$,

$$\sup \left\{ \frac{|f(x) - f(y)|^2}{\mathcal{E}(f, f)} : f \in \mathcal{F}, \mathcal{E}(f, f) > 0 \right\} < \infty.$$

[RF5] If $\bar{f} := (f \wedge 1) \vee 0$, then $\bar{f} \in \mathcal{F}$ and $\mathcal{E}(\bar{f}, \bar{f}) \leq \mathcal{E}(f, f)$ for any $f \in \mathcal{F}$.

SUMMARY

RESISTANCE METRIC R AND MEASURE μ



RESISTANCE FORM $(\mathcal{E}, \mathcal{F})$, DIRICHLET FORM on $L^2(\mu)$



STRONG MARKOV PROCESS X WITH GENERATOR Δ ,
where

$$\mathcal{E}(f, g) = - \int_F (\Delta f) g d\mu.$$

A FIRST EXAMPLE

Let $F = [0, 1]$, $R = \text{Euclidean}$, and μ be a finite Borel measure of full support on $[0, 1]$. Define

$$\mathcal{E}(f, g) = \int_0^1 f'(x)g'(x)dx, \quad \forall f, g \in \mathcal{F},$$

where $\mathcal{F} = \{f \in C([0, 1]) : f \text{ is abs. cont. and } f' \in L^2(dx)\}$. Then $(\mathcal{E}, \mathcal{F})$ is the resistance form associated with $([0, 1], R)$. Moreover, $(\mathcal{E}, \mathcal{F})$ is a regular Dirichlet form on $L^2(\mu)$. Note that

$$\mathcal{E}(f, g) = - \int_0^1 (\Delta f)(x)g(x)\mu(dx), \quad \forall f \in \mathcal{D}(\Delta), g \in \mathcal{F},$$

where $\Delta f = \frac{d}{d\mu} \frac{df}{dx}$, and $\mathcal{D}(\Delta)$ contains those f such that: f' exists and df' is abs. cont. w.r.t. μ , $\Delta f \in L^2(\mu)$, and $f'(0) = f'(1) = 0$.

If $\mu(dx) = dx$, then the Markov process naturally associated with Δ is reflected Brownian motion on $[0, 1]$.

3. CONVERGENCE OF RESISTANCE METRICS AND STOCHASTIC PROCESSES

MAIN RESULT [C. 16]

Write \mathbb{F}_c for the space of marked compact resistance metric spaces, equipped with finite Borel measures of full support. Suppose that the sequence $(F_n, R_n, \mu_n, \rho_n)_{n \geq 1}$ in \mathbb{F}_c satisfies

$$(F_n, R_n, \mu_n, \rho_n) \rightarrow (F, R, \mu, \rho)$$

in the (marked) Gromov-Hausdorff-Prohorov topology for some $(F, R, \mu, \rho) \in \mathbb{F}_c$.

It is then possible to isometrically embed $(F_n, R_n)_{n \geq 1}$ and (F, R) into a common metric space (M, d_M) in such a way that

$$P_{\rho_n}^n \left((X_t^n)_{t \geq 0} \in \cdot \right) \rightarrow P_\rho \left((X_t)_{t \geq 0} \in \cdot \right)$$

weakly as probability measures on $D(\mathbb{R}_+, M)$.

Holds for locally compact spaces if $\limsup_{n \rightarrow \infty} R_n(\rho_n, B_{R_n}(\rho_n, r)^c)$ diverges as $r \rightarrow \infty$. (Can also include 'spatial embeddings'.)

PROOF IDEA 1: RESOLVENTS

For $(F, R, \mu, \rho) \in \mathbb{F}_c$, let

$$G_x f(y) = E_y \int_0^{\sigma_x} f(X_s) ds$$

be the resolvent of X killed on hitting x . NB. Processes associated with resistance forms hit points.

We have [Kigami 12] that

$$G_x f(y) = \int_F g_x(y, z) f(z) \mu(dz),$$

where

$$g_x(y, z) = \frac{R(x, y) + R(x, z) - R(y, z)}{2}.$$

Metric measure convergence \Rightarrow resolvent convergence \Rightarrow semi-group convergence \Rightarrow finite dimensional distribution convergence.

PROOF IDEA 2: TIGHTNESS

Using that X has local times $(L_t(x))_{x \in F, t \geq 0}$, and

$$E_y L_{\sigma_A}(z) = g_A(y, z) = \frac{R(y, A) + R(z, A) - R_A(y, z)}{2},$$

can establish via Markov's inequality a general estimate of the form:

$$\sup_{x \in F} P_x \left(\sup_{s \leq t} R(x, X_s) \geq \varepsilon \right) \leq \frac{32N(F, \varepsilon/4)}{\varepsilon} \left(\delta + \frac{t}{\inf_{x \in F} \mu(B_R(x, \delta))} \right),$$

where $N(F, \varepsilon)$ is the minimal size of an ε cover of F .

Metric measure convergence \Rightarrow estimate holds uniformly in $n \Rightarrow$ tightness (application of Aldous' tightness criterion).

Similar estimate also gives non-explosion in locally compact case.

4. APPLICATIONS

TREES


For any sequence of graph trees $(T_n)_{n \geq 1}$ such that

$$(V(T_n), a_n R_n, b_n \mu_n) \rightarrow (\mathcal{T}, R, \mu),$$

it holds that

$$\left(a_n^{-1} X_{t a_n b_n} \right)_{t \geq 0} \rightarrow (X_t)_{t \geq 0}.$$


- Critical Galton-Watson trees with finite variance conditioned on size, $a_n = n^{1/2}$, $b_n = n$.
- Uniform spanning tree in two dimensions, $a_n = n^{5/4}$, $b_n = n^2$, e.g. after 5,000 and 50,000 steps (picture: Sunil Chhita).



- Many other interesting models...

CONJECTURE FOR CRITICAL PERCOLATION

Bond percolation on integer lattice \mathbb{Z}^d :




At criticality $p = p_c(d)$ in high dimensions, incipient infinite cluster (IIC) conjectured to have same scaling limit as Galton-Watson tree, e.g. [Hara/Slade]. So, expect

$$\left(\text{IIC}, n^{-2} R_{\text{IIC}}, n^{-4} \mu_{\text{IIC}} \right)$$

to converge, and thus obtain scaling limit for random walks. cf. recent work of [Ben Arous, Fribergh, Cabezas 16] for branching random walk.

RANDOM WALK SCALING ON CRITICAL RANDOM GRAPH

Consider largest connected component C_1^n of $G(n, 1/n)$:




It holds that:

$$\left(C_1^n, n^{-1/3}R_n, n^{-2/3}\mu_n\right) \rightarrow (F, R, \mu),$$

cf. [Addario-Berry, Broutin, Goldschmidt 12]. Hence, as in [C. 12],

$$\left(n^{-1/3}X_{tn}^n\right)_{t \geq 0} \rightarrow (X_t)_{t \geq 0}.$$

HEAVY-TAILED RCM ON FRACTALS #1



Suppose that $P(c(x, y) \geq u) = u^{-\alpha}$ for $u \geq 1$ and some $\alpha \in (0, 1)$. For gaskets, can then check that resistance homogenises [C., Hambly, Kumagai 16]

$$\left(V_n, (3/5)^n R_n, 3^{-n} \mu_n \right) \rightarrow (F, R, \mu),$$

where:

- (up to a deterministic constant) R is the standard resistance,
- μ is a Hausdorff measure on fractal.

Hence VSRW converges to Brownian motion (spatial scaling assumes graphs already embedded into limiting fractal):

$$(X_{t5^n}^n)_{t \geq 0} \rightarrow (X_t)_{t \geq 0}.$$

HEAVY-TAILED RCM ON FRACTALS #2

It further holds that

$$\nu_n := 3^{-n/\alpha} \sum_{x \in V_n} c(x) \delta_x \rightarrow \nu = \sum_i v_i \delta_{x_i},$$

in distribution, where $\{(v_i, x_i)\}$ is a Poisson point process with intensity $cv^{-1-\alpha} dv \mu(dx)$. Hence CSRW (and discrete time random walk) converges:

$$\left(X_{t(5/3)^n 3^{n/\alpha}}^{n, \nu_n} \right)_{t \geq 0} \rightarrow (X_t^\nu)_{t \geq 0},$$

where the limiting process X^ν is the **Fontes-Isopi-Newman (FIN)** diffusion on the limiting fractal.

Similarly scaling result for heavy-tailed Bouchaud trap model.