

PARTIAL DIFFERENTIAL EQUATIONS III & V
PROBLEM CLASS 6

In all our exercises in this sheet we will assume that Ω is an open, bounded, and connected set with smooth boundary.

Exercise 1 (*Lower bound for Dirichlet's Energy*). In this problem we will consider the Dirichlet Energy associated to the PDE

$$(1) \quad \begin{cases} -\Delta u = f & x \in \Omega \\ u = 0 & x \in \partial\Omega \end{cases}$$

$$E[u] = \frac{1}{2} \int_{\Omega} |\nabla u(x)|^2 dx - \int_{\Omega} f(x)u(x) dx.$$

- (i) Young's inequality states that for any $a, b \in \mathbb{R}$ and any Hölder conjugate numbers $p, q \in (1, \infty)$ we have that

$$|ab| \leq \frac{|a|^p}{p} + \frac{|b|^q}{q}.$$

Consequently, for any $\varepsilon > 0$ we find that by replacing a with $(p\varepsilon)^{\frac{1}{p}} a$ and b with $\frac{b}{(p\varepsilon)^{\frac{1}{q}}}$ we get that

$$|ab| \leq \varepsilon |a|^p + \frac{|b|^q}{qp^{q-1}\varepsilon^{q-1}}$$

and in particular that for any $\varepsilon > 0$ choosing $p = q = 2$ yields

$$|ab| \leq \varepsilon a^2 + \frac{b^2}{4\varepsilon}.$$

Show that for any $\varepsilon > 0$ and any

$$u \in V = \left\{ v \in C^1(\overline{\Omega}) \mid v = 0 \text{ on } \partial\Omega \right\}$$

we have that

$$E[u] \geq \left(\frac{1}{2} - \varepsilon C_P(\Omega)^2 \right) \|u\|_{H_0^1(\Omega)}^2 - \frac{1}{4\varepsilon} \|f\|_{L^2(\Omega)}^2$$

where $C_P(\Omega)$ is the Poincaré constant associated to the domain Ω .

- (ii) Conclude that there exists a constant $C > 0$ such that

$$\inf_{u \in V} E[u] \geq -C.$$

Exercise 2 (*Uniqueness of weak solutions*). Show that if u and v are weak solutions, in the sense defined in class, for

$$\begin{cases} -\Delta u = f & x \in \Omega \\ u = 0 & x \in \partial\Omega \end{cases}$$

then $u = v$.

Exercise 3 (*Uniqueness for a more general elliptic problem*). Consider the linear, second-order, elliptic PDE

$$(2) \quad \begin{aligned} -\operatorname{div}(A \nabla u) + \mathbf{b} \cdot \nabla u + cu &= f \quad \text{in } \Omega, \\ u &= g \quad \text{on } \partial\Omega, \end{aligned}$$

where $\Omega \subset \mathbb{R}^n$ is open and bounded with smooth boundary, $A \in C^1(\overline{\Omega}; \mathbb{R}^{n \times n})$, $\mathbf{b} \in C^1(\overline{\Omega}; \mathbb{R}^n)$, and $c, f, g \in C(\overline{\Omega})$. Assume that c is non-negative, $\operatorname{div} \mathbf{b} = 0$, and A is uniformly positive definite, i.e., there exists a constant $\alpha > 0$ such that $y^T A(x) y \geq \alpha |y|^2$ for all $y \in \mathbb{R}^n$, $x \in \Omega$. Prove that (2) has at most one solution $u \in C^2(\overline{\Omega})$.

Exercise 4. Consider the space

$$V = \{\varphi \in C^1(\overline{\Omega}) : \varphi = 0 \text{ on } \partial\Omega, \varphi \neq 0\}$$

and the functional $E : V \rightarrow \mathbb{R}$ defined by

$$E[v] = \frac{\int_{\Omega} |\nabla v(x)|^2 dx}{\int_{\Omega} |v(x)|^2 dx}.$$

Suppose that $u \in C^2(\overline{\Omega}) \cap V$ minimises E and show that

$$-\Delta u = \lambda u \quad \text{in } \Omega,$$

$$u = 0 \quad \text{on } \partial\Omega,$$

where $\lambda = E[u]$.

