

## Additional Exercise - Wave Equation

**Exercise.** Consider the wave equation

$$(1) \quad \begin{cases} u_{tt}(x, t) = c^2 u_{xx}(x, t), & (x, t) \in (0, 2\pi) \times (0, \infty), \\ u(0, t) = u(2\pi, t), & t \in (0, \infty), \\ u(x, 0) = f(x), & t \in (0, 2\pi), \\ u_t(x, 0) = 0, & x \in (0, 2\pi), \end{cases}$$

where  $f$  is a smooth function on  $[0, 2\pi]$  with  $f(0) = f(2\pi) = 0$ . We will find a solution to the (1) by utilising the Fourier series (like the heat equation). We assume that we can write

$$u(x, t) = \sum_{n \in \mathbb{Z}} a_n(t) e^{inx}$$

with

$$a_n(t) = \frac{1}{2\pi} \int_0^{2\pi} u(x, t) e^{-inx} dx$$

and that the convergence is such that all interchanging of differentiation and integration is allowed (this will happen, for instance, if we seek a solution that is  $C^4$  on the periodic domain).

(i) Show that  $a_n(t)$  satisfies the equation

$$a_n''(t) + n^2 c^2 a_n(t) = 0.$$

(ii) using the boundary conditions show that

$$a_n(t) = A_n \cos(ncy)$$

and find an explicit expression for  $\{A_n\}_{n \in \mathbb{Z}}$  which depends on  $f$ .

(iii) Write a solution to (1)