1) Consider a Lorentz vector with components  $x^{\mu}$ , which transforms under Lorentz transformations as

$$x^{\mu} \to x'^{\mu} = \Lambda^{\mu}_{\ \nu} x^{\nu}$$

Note that throughout this problem we are using summation convention.

- a) Let  $f^{\mu\nu} \equiv x^{\mu}x^{\nu}$ . Find the transformation behavior of  $f^{\mu\nu}$ ,  $f^{\mu}{}_{\nu} = x^{\mu}x_{\nu}$ and  $f_{\mu\nu} = x_{\mu}x_{\nu}$  under Lorentz transformations.
- b) For another Lorentz vector  $y^{\mu}$ , find the transformation behavior of  $f^{\mu\nu}y_{\mu}$  under Lorentz transformations.
- c) Compute

$$\sum_{\mu} \frac{\partial}{\partial x^{\mu}} x^{\mu} \, .$$

d) Work out the transformation behavior of

$$\frac{\partial}{\partial x^{\mu}}$$

under Lorentz transformations. Use c) to argue for the same result.

## solution:

(a) We can infer the transformation of  $f^{\mu\nu}$ ,  $f^{\mu}_{\nu}$ ,  $f_{\mu\nu}$  from that of  $x^{\mu}$  and  $x_{\mu}$ 

$$\begin{aligned} f^{\mu\nu} &\to \Lambda^{\mu}_{\ \mu'} \Lambda^{\nu}_{\ \nu'} f^{\mu'\nu'} \\ f^{\mu}_{\ \nu} &\to \Lambda^{\mu}_{\ \mu'} f^{\mu'}_{\ \nu'} (\Lambda^{-1})^{\nu'}_{\ \nu} \\ f_{\mu\nu} &\to f_{\mu'\nu'} (\Lambda^{-1})^{\mu'}_{\ \mu} (\Lambda^{-1})^{\nu'}_{\ \nu} \end{aligned} (0.1)$$

Here the ordering of things I used on the rhs is not really important, I have written things in such a way that serves the slogan **upper indices** transform with  $\Lambda$  and lower indices transform with  $\Lambda^{-1}$  from the right.

(b) Using the result of a) and the fact that  $y_{\mu}$  transforms with a  $\Lambda^{-1}$  we immediately see that

$$f^{\mu\nu}y_{\mu} \to \Lambda^{\nu}{}_{\nu'}f^{\mu\nu'}y_{\mu} \tag{0.2}$$

I.e.  $\mu$  is a contracted dummy index and the only non-trivial transformation is coming from  $\nu$ .

(c) This is simply

$$\sum_{\mu} \frac{\partial}{\partial x^{\mu}} x^{\mu} = \frac{\partial x^{0}}{\partial x^{0}} + \frac{\partial x^{1}}{\partial x^{1}} + \frac{\partial x^{2}}{\partial x^{2}} + \frac{\partial x^{3}}{\partial x^{3}} = 4$$
(0.3)

(d) Writing

$$x^{\prime\mu} = \Lambda^{\mu}_{\ \nu} x^{\nu} \tag{0.4}$$

implies that

$$\frac{\partial}{\partial x'^{\mu}} = \frac{\partial}{\partial x^{\nu}} \frac{\partial x^{\nu}}{\partial x'^{\mu}} = \frac{\partial}{\partial x^{\nu}} (\Lambda^{-1})^{\nu}{}_{\mu} \tag{0.5}$$

The derivative with respect to a Lorentz vector hence transforms like a Lorentz covector. For this reason people usually write  $\frac{\partial}{\partial x^{\mu}} \equiv \partial_{\mu}$ . This result can also be seen from part c): The number 4 is a Lorentz scalar and as  $x^{\mu}$  is a Lorentz vector  $\frac{\partial}{\partial x^{\mu}}$  must be a covector to get something invariant.

2) Write a 4-vector  $(x^0, x^1, x^2, x^3)$  as a matrix  $M_x$  with  $M_x^{\dagger} = M_x$ :

$$M_x := \begin{pmatrix} x^0 + x^3 & x^1 - ix^2 \\ x^1 + ix^2 & x^0 - x^3 \end{pmatrix}.$$
 (0.6)

For  $g \in SL(2, \mathbb{C})$  define an action F(g) on  $\mathbb{R}^4$  by

$$g \to F(g)$$
  $F(g)M_x := gM_x g^{\dagger}$ . (0.7)

- a) Show that F is a homomorphism from  $SL(2, \mathbb{C})$  to L.
- b) For a rotation in the  $x^1, x^2$ -plane, find the element  $g \in SL(2, \mathbb{C})$  that is mapped to it by F. Repeat the same for a boost along the  $x^1$  direction.

## solution:

a) First note that  $M_x$  is the most general  $2 \times 2$  matrix with the property  $M_x = M_x^{\dagger}$ . This property is preserved by F(g) as

$$(gM_xg^{\dagger})^{\dagger} = g^{\dagger\dagger}M_x^{\dagger}g^{\dagger} = gM_xg^{\dagger}.$$
 (0.8)

Furthermore F(g) acts as a linear map on  $\mathbb{R}^{1,3}$  which preserves det  $M_x$ :

$$\det M_x \to \det(gM_xg^{\dagger}) = \det g \det M_x \det g^{\dagger} = \det M_x \,. \tag{0.9}$$

As  $x_{\mu}x^{\mu} = -(x^0)^2 + (x^1)^2 + (x^2)^2 + (x^3)^2 = -\det M_x$ , the linear map F(g) preserves the length of vectors in  $\mathbb{R}^{1,3}$  and is hence in L.

In fact, we can argue that F(g) is contained in  $L^{\uparrow}_{+}$  (you weren't asked this for the assignment, but it is good to know). The group  $SL(2, \mathbb{C})$  is connected as the following argument shows: by a standard result from linear algebra, we can write any matrix in  $SL(2, \mathbb{C})$  as

$$g = B \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix} B^{-1} \tag{0.10}$$

We can now simply let b go to zero and a go to 1 continuously to connect any element in  $SL(2, \mathbb{C})$  continuously to the identity. As  $SL(2, \mathbb{C})$  is connected and L has four connected components, F can only map to one of them (it is a continuous map). Using g = 1 we see that Fmaps to  $L_{+}^{\uparrow}$ , the component containing the identity. This map is not injective, as g and -g are mapped to the same F(g).

b) Now let us consider how different matrices in  $SL(2, \mathbb{C})$  act on  $M_x$ . First we investigate elements of  $SL(2, \mathbb{C})$  that are in SU(2). For  $\theta \in \mathbb{R}$  set

$$g_{3}(\theta) := e^{i\theta\sigma_{3}} = \begin{pmatrix} e^{i\theta} & 0\\ 0 & e^{-i\theta} \end{pmatrix}$$

$$g_{2}(\theta) := e^{i\theta\sigma_{2}} = \begin{pmatrix} \cos(\theta) & \sin(\theta)\\ -\sin(\theta) & \cos(\theta) \end{pmatrix}$$

$$g_{1}(\theta) := e^{i\theta\sigma_{1}} = \begin{pmatrix} \cos(\theta) & i\sin(\theta)\\ i\sin(\theta) & \cos(\theta) \end{pmatrix}$$

$$(0.11)$$

Their action on  $M_x$  is

$$g_{3}M_{x}g_{3}^{\dagger} = \begin{pmatrix} x^{0} + x^{3} & e^{2i\theta}(x^{1} - ix^{2}) \\ e^{-2i\theta}(x^{1} + ix^{2}) & x^{0} - x^{3} \end{pmatrix}$$

$$g_{2}M_{x}g_{2}^{\dagger} = \begin{pmatrix} x^{0} + x^{1}\sin(2\theta) + x^{3}\cos(2\theta) & x^{1}\cos(2\theta) - ix^{2} - x^{3}\sin(2\theta) \\ x^{1}\cos(2\theta) + ix^{2} - x^{3}\sin(2\theta) & x^{0} - x^{1}\sin(2\theta) - x^{3}\cos(2\theta) \end{pmatrix}$$

$$g_{1}M_{x}g_{1}^{\dagger} = \begin{pmatrix} x^{0} - x^{2}\sin(2\theta) + x^{3}\cos(2\theta) & x^{1} - ix^{2}\cos(2\theta) - ix^{3}\sin(2\theta) \\ x^{1} + ix^{2}\cos(2\theta) + ix^{3}\sin(2\theta) & x^{0} + x^{2}\sin(2\theta) - x^{3}\cos(2\theta) \end{pmatrix}$$

$$(0.12)$$

This is the same we observed when we studied the same action of SU(2)on  $\mathbb{R}^3$ :  $g_i$  defines a rotation by angle of magnitude  $2\theta$  around the  $x_i$ axis is  $\mathbb{R}^3$  with coordinates  $x^1, x^2, x^3$ . Similarly, one can parametrize rotations around arbitrary axis. As we can write any matrix in SO(3)as product of such elementary rotations (see again problem class 1), the map from  $SU(2) \subset SL(2, \mathbb{C})$  is surjective onto  $SO(3) \subset L_+^{\uparrow}$ . We can realize three other independent elements of  $SL(2, \mathbb{C})$  as

$$h_{3}(\theta) := e^{\theta \sigma_{3}} = \begin{pmatrix} e^{\theta} & 0\\ 0 & e^{-\theta} \end{pmatrix}$$

$$h_{2}(\theta) := e^{\theta \sigma_{2}} = \begin{pmatrix} \cosh(\theta) & -i\sinh(\theta)\\ i\sinh(\theta) & \cosh(\theta) \end{pmatrix}$$

$$h_{1}(\theta) := e^{\theta \sigma_{1}} = \begin{pmatrix} \cosh(\theta) & \sinh(\theta)\\ \sinh(\theta) & \cosh(\theta) \end{pmatrix}$$
(0.13)

where again  $\theta \in \mathbb{R}$ . Note that now  $h_j^{\dagger} \neq h_j^{-1}$  but still det  $h_j = 1$ . We now have

$$h_3 M_x h_3^{\dagger} = \begin{pmatrix} e^{2\theta} (x^0 + x^3) & x^1 - ix^2 \\ x^1 + ix^2 & e^{-2\theta} (x^0 - x^3) \end{pmatrix}$$
(0.14)

As  $e^{\theta} = \cosh \theta + \sinh \theta$  for real  $\theta$  we can summarize this as

$$\begin{pmatrix} x^{0} \\ x^{1} \\ x^{2} \\ x^{3} \end{pmatrix} \to \begin{pmatrix} \cosh 2\theta & 0 & 0 & \sinh 2\theta \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ \sinh 2\theta & 0 & 0 & \cosh 2\theta \end{pmatrix} \begin{pmatrix} x^{0} \\ x^{1} \\ x^{2} \\ x^{3} \end{pmatrix}$$
(0.15)

i.e. this exactly a boost in (minus) the  $x^3$  direction. Boosts along the  $x^2$  and  $x^1$  axis are likewise realized by  $h_2$  and  $h_1$  and we can again find a boost along an arbitrary direction by exponentiating appropriate real linear combinations of the  $\sigma_j$ . In particular taking

$$h_1 = \begin{pmatrix} \cosh \theta & \sinh \theta \\ \sinh \theta & \cosh \theta \end{pmatrix} \tag{0.16}$$

gives

$$h_1 M h_1^{\dagger} = \begin{pmatrix} x_3 + x_0 \cosh 2\theta + x_1 \sinh 2\theta & -ix_2 + x_1 \cosh 2\theta + x_2 \sinh 2\theta \\ ix_2 + x_1 \cosh 2\theta + x_2 \sinh 2\theta & -x_3 + x_0 \cosh 2\theta + x_1 \sinh 2\theta \end{pmatrix}$$
(0.17)

which is a boost in the  $x_1$  direction.

You weren't asked to do this for the assignment, but we can now comment on how we should show that F(g) is a surjective homomorphism. Given the above, it should be clear that we can write any rotation and any boost in  $L_+^{\uparrow}$  as the image of an element of  $SL(2, \mathbb{C})$  under F(g). In the lectures we have stated a thoerem that every element of  $L_+^{\uparrow}$  can be written as the product of an element of SO(3) and a boost, hence every element in  $L_+^{\uparrow}$  is in the image of F. Here are some things to ponder:

- 1. How is the Lorent group defined? Why is it defined that way?
- 2. What's the point about upper/lower indices?