

- 1) Let  $\mathbb{C}$  be the complex numbers and  $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$ . Which of these is a group under addition? Which of these is a group under multiplication?

**Solution:**

$\mathbb{C}$  satisfies the definition of a group under addition, here the inverse of  $z$  is  $-z$  and the neutral element is 0. Under multiplication, 0 does not have an inverse, so it is not a group under multiplication.

$\mathbb{C}^*$  satisfies the definition of a group under multiplication, here the inverse of  $z$  is  $z^{-1}$  and the neutral element is 1. It is not a group under addition as it does not contain a neutral element for that.

- 2) Consider the set  $V$  of real  $n \times n$  matrices.
- a) Show that  $V$  is a (real) vector space.
  - b) Let  $U \subset V$  be the set of matrices with determinant 1. Is  $U$  a vector space as well?
  - c) For any matrix  $Q$  in  $V$  define a map

$$g_M : Q \rightarrow M^{-1}QM$$

where  $M$  is a fixed invertible matrix. Show that  $g_M$  is a linear map on  $V$ .

**Solution:**

- a) We need to show that for any vectors  $v, v' \in V$ ,  $v + v'$  and  $av$  for  $a \in \mathbb{R}$  is also in  $V$ . For arbitrary real  $n \times n$  matrices  $S, S'$ , both the sum  $S + S'$  and a rescaled  $aS$  is another real  $n \times n$  matrix. Hence this is a vector space.
- b) Here we are in trouble: for  $S \in U$  we have  $\det S = 1$ . But to be a vector space,  $U$  also has to contain  $aS$  for all  $a \in \mathbb{R}$ . But  $\det aS = a^n$ , so  $aS$  is not in  $U$ . Hence  $U$  is not a vector space.
- c) First note that for  $Q \in V$ ,  $g_M(Q)$  is another real  $n \times n$  matrix, the map hence maps elements in  $V$  to elements in  $V$  again. To check linearity we work out that for  $Q, P \in V$  we have

$$g_M(Q + P) = M^{-1}(Q + P)M = M^{-1}QM + M^{-1}PM = g_M(Q) + g_M(P) \quad (0.1)$$

so  $g_M$  is a linear map acting on  $V$ .

- 3) a) By working out the derivative of

$$\lim_{n \rightarrow \infty} (1 + i\phi/n)^n$$

with respect to  $\phi$ , show that this expression satisfies the same differential equation as  $e^{i\phi}$ . You may assume that you can swap the order of the limit and taking the derivative.

As both functions have the same value at  $\phi = 0$  this implies that they are equal by the uniqueness of solutions of ordinary differential equations.

- b) Consider a square matrix  $A$  and let  $g = e^{iA}$ , which is defined via the Taylor series of the exponential. Show that

$$\lim_{n \rightarrow \infty} (1 + iA/n)^n = e^{iA}.$$

**Solution:**

- a) The differential equation of the exponential  $g = e^{i\phi}$  is

$$\frac{\partial}{\partial \phi} g = ig. \quad (0.2)$$

We now work out

$$\frac{\partial}{\partial \phi} (1 + i\phi/n)^n = n(1 + i\phi/n)^{n-1} \frac{i}{n} = \frac{i(1 + i\phi/n)^n}{1 + i\phi/n}. \quad (0.3)$$

In the limit  $n \rightarrow \infty$ , the rhs converges to  $i(1 + i\phi/n)^n$ . Assuming that differentiation and taking the limit commute (which we can do by using uniform convergence), we have hence shown that

$$\frac{\partial}{\partial \phi} \lim_{n \rightarrow \infty} (1 + i\phi/n)^n = i \lim_{n \rightarrow \infty} (1 + i\phi/n)^n. \quad (0.4)$$

Letting  $e(\phi) \equiv \lim_{n \rightarrow \infty} (1 + i\phi/n)^n$ , we have just shown that

$$\frac{\partial}{\partial \phi} e(\phi) = ie(\phi) \quad (0.5)$$

and can conclude that  $e(\phi) = ce^{i\phi}$  using uniqueness of solutions of ordinary differential equations. Setting  $\phi = 0$  the constant of proportionality  $c$  is seen to be 1 and we are done.

- b) We can repeat the proof in the lecture by replacing  $\phi$  with  $A$  in every step.

Here are some things to ponder:

1. What is a group? Why are symmetries groups?
2. What does the  $U$  in  $U(1)$  stand for ? What might somebody mean when they say  $U(n)$  ?
3. How many ways to define the exponential function can you come up with ?
4. Tangents are linear approximations.