

Homework 11-12
Starred problems due on Thursday, 1 February.

Isometries and conformal maps - 1

11.1. Let $a > 0$. Construct explicitly a local isometry from the plane $P = \{(u, v, 0) \in \mathbb{R}^3 \mid u, v \in \mathbb{R}\}$ onto the cylinder $S = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 = a^2\}$.

11.2. (*) Let b be a positive number such that $\sqrt{1 + b^2}$ is an integer n . Let S be the circular cone obtained by rotating the curve given by $\alpha(v) = (v, 0, bv)$, $v > 0$, about the z -axis. Let the coordinate xy -plane P be parametrized by polar coordinates (r, ϑ) :

$$\mathbf{x}: U = (0, \infty) \times (0, 2\pi) \longrightarrow P, \quad \mathbf{x}(r, \vartheta) = (r \cos \vartheta, r \sin \vartheta, 0).$$

Show that the map $f: P \setminus \{(0, 0, 0)\} \longrightarrow S$ defined on $\mathbf{x}(U)$ by

$$f(\mathbf{x}(r, \vartheta)) = \frac{1}{n}(r \cos n\vartheta, r \sin n\vartheta, br)$$

is a local isometry on $\mathbf{x}(U)$.

11.3. Let S_1, S_2, S_3 be regular surfaces.

- (a) Suppose that $f: S_1 \longrightarrow S_2$ and $g: S_2 \longrightarrow S_3$ are local isometries. Prove that $g \circ f: S_1 \longrightarrow S_3$ is a local isometry.
- (b) Suppose that $f: S_1 \longrightarrow S_2$ and $g: S_2 \longrightarrow S_3$ are conformal maps with conformal factors $\lambda: S_1 \longrightarrow (0, \infty)$ and $\mu: S_2 \longrightarrow (0, \infty)$, respectively. Prove that $g \circ f: S_1 \longrightarrow S_3$ is a conformal map and calculate its conformal factor. (The conformal factor of a conformal map is the function appearing as factor in front of the inner product in the definition.)
- (c) Let f and g be conformal maps with conformal factors λ and μ as in the previous part. Find a condition on λ and μ such that $g \circ f$ is a (local) isometry.

11.4. Let S be the surface of revolution parametrized by

$$\mathbf{x}(u, v) = \left(\cos v \cos u, \cos v \sin u, -\sin v + \log \tan \left(\frac{\pi}{4} + \frac{v}{2} \right) \right),$$

where $0 < u < 2\pi, 0 < v < \pi/2$. Let S_1 be the region

$$S_1 = \{ \vec{x}(u, v) \mid 0 < u < \pi, 0 < v < \pi/2 \}$$

and let S_2 be the region

$$S_2 = \{ \vec{x}(u, v) \mid 0 < u < 2\pi, \pi/3 < v < \pi/2 \}.$$

Show that the map

$$\mathbf{x}(u, v) \mapsto \mathbf{x} \left(2u, \arccos \left(\frac{1}{2} \cos v \right) \right)$$

is an isometry from S_1 onto S_2 .

Isometries and conformal maps - 2

12.1. (*) Let S be a surface of revolution. Prove that any rotation about the axis of revolution is an isometry of S .

12.2. The disc model of the hyperbolic plane.

Let \mathbb{D} denote the unit disc $\{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 < 1\}$ with first fundamental form

$$\tilde{E} = \tilde{G} = \frac{4}{(1 - x^2 - y^2)^2}, \quad \tilde{F} = 0.$$

Let \mathbb{H} be the hyperbolic plane with coordinates $(u, v) \in \mathbb{R} \times (0, \infty)$ and first fundamental form

$$E = G = \frac{1}{v^2}, \quad F = 0.$$

Show that the map $f: \mathbb{H} \rightarrow \mathbb{D}$ given by

$$f(z) = \frac{z - i}{z + i}, \quad z = u + iv \in \mathbb{H},$$

is an isometry.

12.3. Hyperboloid model of the hyperbolic plane.

Let $Q: \mathbb{R}^3 \rightarrow \mathbb{R}$ be the quadratic form defined by

$$Q(x_1, x_2, x_3) = x_1^2 + x_2^2 - x_3^2, \quad (x_1, x_2, x_3) \in \mathbb{R}^3$$

(the quadratic space (\mathbb{R}^3, Q) is usually denoted by $\mathbb{R}^{2,1}$). Let

$$S = \{(x_1, x_2, x_3) \in \mathbb{R}^3 \mid Q(x_1, x_2, x_3) = -1\}$$

(i.e. S is a hyperboloid of two sheets).

Recall that the *induced quadratic form* $I_{\mathbf{p}}$ on each tangent plane $T_{\mathbf{p}}S$ is defined by $I_{\mathbf{p}}(\mathbf{w}) = Q(\mathbf{w})$ for every $\mathbf{w} \in T_{\mathbf{p}}(S)$. Show that $I_{\mathbf{p}}$ is positive definite and that the map $f: \mathbb{D} \rightarrow S$ from the disc model of the hyperbolic plane (see the previous exercise) defined by

$$f(x, y) = \frac{1}{1 - x^2 - y^2} (2x, 2y, 1 + x^2 + y^2), \quad (x, y) \in \mathbb{D},$$

maps \mathbb{D} isometrically onto the component of S for which $x_3 > 0$.