

Homework 9-10

First fundamental form

9.1. Find the coefficients of the first fundamental forms of:

(a) the *catenoid* parametrized by

$$\mathbf{x}(u, v) = (\cosh v \cos u, \cosh v \sin u, v), \quad (u, v) \in U := (0, 2\pi) \times \mathbb{R};$$

(b) the *helicoid* parametrized by

$$\tilde{\mathbf{x}}(u, v) = (-\sinh v \sin u, \sinh v \cos u, -u), \quad (u, v) \in U;$$

(c) the surface S_ϑ (for some $\vartheta \in \mathbb{R}$) parametrized by

$$\mathbf{y}_\vartheta(u, v) = (\cos \vartheta)\mathbf{x}(u, v) + (\sin \vartheta)\tilde{\mathbf{x}}(u, v), \quad (u, v) \in U.$$

9.2. Find the coefficients of the first fundamental form of $S^2(1)$ with respect to the local parametrization \mathbf{x} defined in Exercise 6.2.

9.3. Let $U = \mathbb{R} \times (0, \infty)$, and let $\mathbf{x} : U \rightarrow \mathbb{R}^n$ be a parametrization of a surface \mathbb{H} in \mathbb{R}^2 with corresponding coefficients of the first fundamental form $E(u, v) = G(u, v) = 1/v^2$ and $F(u, v) = 0$ for all $(u, v) \in U$. Then \mathbb{H} is called the *hyperbolic plane*. For $r > 0$ denote by $\boldsymbol{\alpha} : (0, \pi) \rightarrow \mathbb{H}$ the curve given by

$$\boldsymbol{\alpha}(t) = \mathbf{x}(r \cos t, r \sin t).$$

Show that the length of $\boldsymbol{\alpha}$ in \mathbb{H} from $\boldsymbol{\alpha}(\pi/6)$ to $\boldsymbol{\alpha}(5\pi/6)$ is equal to

$$\int_{\pi/6}^{5\pi/6} \frac{1}{\sin t} dt.$$

(In fact, $\boldsymbol{\alpha}$ is the curve of shortest length between its endpoints.) Now take $r = \sqrt{2}$ and find the angle of intersection of $\boldsymbol{\alpha}$ with the curve $\boldsymbol{\beta}(s) = \mathbf{x}(1, s)$ at their point of intersection.

9.4. Let S be a surface parametrized by

$$\mathbf{x}(u, v) = (u \cos v, u \sin v, \log \cos v + u), \quad (u, v) \in U := \mathbb{R} \times \left(-\frac{\pi}{2}, \frac{\pi}{2}\right).$$

For $c \in (-\pi/2, \pi/2)$, let $\boldsymbol{\alpha}_c$ be the curve given by $\boldsymbol{\alpha}_c(u) = \mathbf{x}(u, c)$. Show that the length of $\boldsymbol{\alpha}_c$ from $u = u_0$ to $u = u_1$ does not depend on c .

Coordinate curves, angles and area

- 10.1.** Let $\mathbf{x} : U \rightarrow S$ be a local parametrization of a regular surface S , and denote by E, F, G the coefficients of the first fundamental form in this parametrization. Show that the tangent vector $a\partial_u\mathbf{x} + b\partial_v\mathbf{x}$ bisects the angle between the coordinate curves if and only if

$$\sqrt{G}(aE + bF) = \sqrt{E}(aF + bG).$$

Further, if

$$\mathbf{x}(u, v) = (u, v, u^2 - v^2),$$

find a vector tangential to S which bisects the angle between the coordinate curves at the point $(1, 1, 0) \in S$.

- 10.2.** Find two families of curves on the helicoid parametrized by

$$\mathbf{x}(u, v) = (v \cos u, v \sin u, u)$$

which, at each point, bisect the angles between the coordinate curves.

(Show that they are given by $u \pm \sinh^{-1} v = c$, where c is a constant on each curve in the family.)

- 10.3.** The coordinate curves of a parametrization $\mathbf{x}(u, v)$ constitute a *Chebyshev net* if the lengths of the opposite sides of any quadrilateral formed by them are equal.

(a) Show that a necessary and sufficient condition for this is

$$\frac{\partial E}{\partial v} = \frac{\partial G}{\partial u} = 0$$

(b) Show that if coordinate curves constitute a Chebyshev net, then it is possible to reparametrize the coordinate neighborhood in such a way that the new coefficients of the first fundamental form are

$$E = 1, \quad F = \cos \vartheta, \quad G = 1,$$

where ϑ is the angle between coordinate curves.

- 10.4.** Show that a surface of revolution can always be parametrized so that

$$E = E(v), \quad F = 0, \quad G = 1$$

- 10.5.** Let S be the surface $\{(x, y, z) \in \mathbb{R}^3 \mid z = x^2 - y^2\}$ and let \mathcal{F} be the family of curves on S obtained as the intersection of S with the planes $z = \text{const}$. Find the family of curves on S which meet \mathcal{F} orthogonally and show that they are the intersections of S with the family of hyperbolic cylinders $xy = \text{const}$.

- 10.6.** Using the notation of Exercise 10.2, show that the family of curves orthogonal to the family

$$v \cos u = \text{const}$$

is the family defined by $(1 + v^2) \sin^2 u = \text{const}$.