

Homework 1-2

Starred problems due on Thursday, 24 October.

Plane curves - 1

1.1. Sketch the trace of the smooth curve given by $\alpha(u) = (u^5, u^2 - 1)$, and mark the singular points.

1.2. Let $\alpha : I \rightarrow \mathbb{R}^2$ be a smooth curve, and let $[a, b] \subset I$ be a closed interval. For every partition $a = u_0 < u_1 < \dots < u_n = b$ consider the sum

$$\ell_{\alpha, P} := \sum_{i=1}^n \|\alpha(u_i) - \alpha(u_{i-1})\|$$

where P stands for the given partition. Give a geometric interpretation of $\ell_{\alpha, P}$. What length does $\ell_{\alpha, P}$ measure? Now assume that the partition becomes *finer*, i.e., $\|P\| := \max_{i=1, \dots, n} |u_i - u_{i-1}|$ becomes smaller. What is the limit of $\ell_{\alpha, P}$ as $\|P\| \rightarrow 0$?

1.3. (*) An *epicycloid* α is obtained as the locus of a point on the circumference of a circle of radius r which rolls without slipping on a circle of the same radius.

(a) Sketch α .

(b) Show that the epicycloid can be parametrized by

$$\alpha(u) = (2r \sin u - r \sin 2u, 2r \cos u - r \cos 2u), \quad u \in \mathbb{R}.$$

Find the length of α between the singular points at $u = 0$ and $u = 2\pi$.

1.4. (*) (a) Let $\alpha(u)$ and $\beta(u)$ be two smooth plane curves. Show that

$$\frac{d}{du}(\alpha(u) \cdot \beta(u)) = \alpha'(u) \cdot \beta(u) + \alpha(u) \cdot \beta'(u),$$

where $\alpha(u) \cdot \beta(u)$ denotes a Euclidean dot product of vectors $\alpha(u)$ and $\beta(u)$.

Hint: write $\alpha(u) = (\alpha_1(u), \alpha_2(u))$, $\beta(u) = (\beta_1(u), \beta_2(u))$ and compute everything in coordinates.

(b) Let $\alpha(u) : I \rightarrow \mathbb{R}^2$ be a smooth curve which does not pass through the origin. Suppose there exists $u_0 \in I$ such that the point $\alpha(u_0)$ is the closest to the origin amongst all the points of the trace of α . Show that $\alpha(u_0)$ is orthogonal to $\alpha'(u_0)$.

1.5. The second derivative $\alpha''(u)$ of a smooth plane curve $\alpha(u)$ is identically zero. What can be said about α ?

1.6. Let $\alpha : (0, \pi) \rightarrow \mathbb{R}^2$ be a curve defined by

$$\alpha(u) = \left(\sin u, \cos u + \log \tan \frac{u}{2} \right)$$

The trace of α is called a *tractrix*.

(a) Sketch α .

(b) Show that a tangent vector at $\alpha(u_0)$ can be written as

$$\alpha'(u_0) = \left(\cos u_0, -\sin u_0 + \frac{1}{\sin u_0} \right)$$

Show that $\alpha(u)$ is smooth, and it is regular everywhere except $u = \pi/2$.

(c) Write down the equation of a tangent line l_{u_0} to the trace of α at $\alpha(u_0)$.

(d) Show that the distance between $\alpha(u_0)$ and the intersection of l_{u_0} with y -axis is constantly equal to 1.

Plane curves - 2

2.1. The *catenary* is the plane curve $\alpha : \mathbb{R} \rightarrow \mathbb{R}^2$ given by $\alpha(u) = (u, \cosh u)$. It is the curve assumed by a uniform chain hanging under the action of gravity. Sketch the curve. Find its curvature.

2.2. Suppose that $\alpha : I \rightarrow \mathbb{R}^2$ is a regular curve, but not necessarily unit speed. Write $\alpha(u) = (x(u), y(u))$. Find the formula for the curvature $\kappa(u)$ at the parameter value u in terms of the functions x and y (and their derivatives) at u .

Hint: consider the corresponding curve $\tilde{\alpha}$ parametrised by arc length. The curvature $\tilde{\kappa}$ of $\tilde{\alpha}$ is then $\tilde{\kappa}(s) = \tilde{\mathbf{n}}(s) \cdot \tilde{\mathbf{t}}'(s)$, where $\tilde{\mathbf{t}}$ and $\tilde{\mathbf{n}}$ are the unit tangent and unit normal vector of $\tilde{\alpha}$. Use the relation $\tilde{\alpha}(s) = \alpha(\ell^{-1}(s))$, where $s = \ell(u)$ is the arc length, together with the chain rule.

2.3. (*) Compute the curvature of tractrix (see Exercise 1.6) at $\alpha(u)$.

2.4. Let $\alpha : I \rightarrow \mathbb{R}^2$ be a smooth regular plane curve.

(a) Assume that for some $u_0 \in I$ the normal line to α at $\alpha(u_0)$ passes through the origin. Show that for some $\epsilon > 0$ the trace $\alpha(u_0 - \epsilon, u_0 + \epsilon)$ can be written in polar coordinates as

$$\beta(\vartheta) = (\rho(\vartheta) \cos \vartheta, \rho(\vartheta) \sin \vartheta)$$

for an appropriate smooth function $\rho(\vartheta)$, where $\vartheta \in J$ for some interval J .

(b) Assume that all normal lines to α pass through the origin. Show that the trace of α is contained in a circle.

(c) Let $\alpha : I \rightarrow \mathbb{R}^2$ be given in polar coordinates by

$$\alpha(\vartheta) = (\rho(\vartheta) \cos \vartheta, \rho(\vartheta) \sin \vartheta), \quad \vartheta \in [a, b]$$

Show that the length of α is

$$\int_a^b \sqrt{\rho^2 + (\rho')^2} d\vartheta$$

(d) In the assumptions of (c), show that the curvature of α is

$$\kappa(\vartheta) = \frac{2(\rho')^2 - \rho\rho'' + \rho^2}{[\rho^2 + (\rho')^2]^{3/2}}$$

2.5. Find an arc length parameter for the graphs of the following functions $f, g : (0, \infty) \rightarrow \mathbb{R}$:

(a) $f(x) = ax + b$, $a, b \in \mathbb{R}$;

(b)(*) $g(x) = \frac{8}{27}x^{3/2}$.