

Solutions 7-8

- 7.1. (a) For an affine transformation $f = A\mathbf{v} + \mathbf{b}$ in affine 2-dimensional space, find f^{-1} .
 (b) Find an affine transformation $f(\mathbf{v}) = A\mathbf{v} + \mathbf{b}$ which maps the points $(0, 0)$, $(1, 0)$, $(0, 1)$ to the points $(0, 1)$, $(2, 4)$, $(4, 4)$ respectively.
 (c) Find an affine transformation $g(\mathbf{v}) = C\mathbf{v} + \mathbf{d}$ which maps the points $(4, 7)$, $(9, 6)$, $(-2, 8)$ to the points $(0, 0)$, $(1, 0)$, $(0, 1)$ respectively.
 (d) Use (b) and (c) to find an affine transformation $h(\mathbf{v}) = E\mathbf{v} + \mathbf{q}$ which maps the points $(4, 7)$, $(9, 6)$, $(-2, 8)$ to the points $(0, 1)$, $(2, 4)$, $(4, 4)$ respectively.

Solution. The question (d) may be solved by a direct computation (finding 6 variables from 6 linear equations). The sequence of subquestions (a)-(d) presents the easier and more reliable way to do that.

(a) f is a linear transformation followed by a translation, so, f^{-1} is a translation (by $-\mathbf{b}$) followed by the inverse linear transformation, $f^{-1}(\mathbf{v}) = A^{-1}(\mathbf{v} - \mathbf{b})$.

(b) Looking at the image of $(0, 0)$ we find $\mathbf{b} = (0, 1)$. Then looking at the image of $(1, 0)$ we find the first column of A , and from the image of $(0, 1)$ we find the second column of A , we get

$$f(\mathbf{v}) = \begin{pmatrix} 2 & 4 \\ 3 & 3 \end{pmatrix} \mathbf{v} + \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

(c) Similarly to (b) we first find the inverse transformation g^{-1} (takes the points $(0, 0)$, $(1, 0)$, $(0, 1)$ to the points $(4, 7)$, $(9, 1)$, $(3, 8)$):

$$g^{-1}(\mathbf{v}) = \begin{pmatrix} 5 & -6 \\ -1 & 1 \end{pmatrix} \mathbf{v} + \begin{pmatrix} 4 \\ 7 \end{pmatrix}.$$

Then, using the result of (a) and the equation $\begin{pmatrix} a & b \\ c & d \end{pmatrix}^{-1} = \frac{1}{ad-bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$ we get

$$g(\mathbf{v}) = \begin{pmatrix} -1 & -6 \\ -1 & -5 \end{pmatrix} \left(\mathbf{v} - \begin{pmatrix} 4 \\ 7 \end{pmatrix} \right) = \begin{pmatrix} -1 & -6 \\ -1 & -5 \end{pmatrix} \mathbf{v} + \begin{pmatrix} 46 \\ 39 \end{pmatrix}.$$

(d) We find

$$h(\mathbf{v}) = f \circ g(\mathbf{v}) = \begin{pmatrix} 2 & 4 \\ 3 & 3 \end{pmatrix} \left(\begin{pmatrix} -1 & -6 \\ -1 & -5 \end{pmatrix} \mathbf{v} + \begin{pmatrix} 46 \\ 39 \end{pmatrix} \right) + \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} -6 & -32 \\ -6 & -33 \end{pmatrix} \mathbf{v} + \begin{pmatrix} 248 \\ 256 \end{pmatrix}.$$

- 7.2. (*) Through each vertex of a triangle, two lines dividing the opposite side into three equal parts are drawn. Let P be the hexagon bounded by these six lines. Prove that the diagonals joining the opposite vertices of P are concurrent.

Solution. Consider an affine transformation f which takes the given triangle ABC to a regular triangle $A'B'C'$. Notice that as affine transformations preserve collinearity and the ratios of lengths on every line, the six lines in ABC will map to the lines dividing the sides of $A'B'C'$ in proportion 1:2 and 2:1. So, the image $f(P)$ is symmetric with respect to every median on $A'B'C'$. In particular, the diagonals of $f(P)$ lie on the medians of $A'B'C'$. As the medians of $A'B'C'$ are concurrent, the diagonals of $f(P)$ also intersect in one point (since $f(P)$ preserves concurrency). As f (being an affine transformation) preserves concurrency, we conclude that the diagonals of P are also concurrent.

- 7.3. Prove that an arbitrary convex pentagon $ABCDE$ with sides parallel to its diagonals (i.e. such that $AB \parallel CE, BC \parallel DA$, etc) can be affinely transformed into a regular pentagon.

Solution. Let $A'B'C'D'E'$ be a regular pentagon. Let f be an affine map which takes the points A, B, C to $A'B'C'$. As affine maps preserve the parallelism, $f(D)$ lies on the line parallel to $B'C'$ through A' , i.e., $f(D)$ lies on the line $A'D'$. Similarly, $f(E)$ lies on $C'E'$. Furthermore, the side $f(DE)$ is parallel to $A'C' = f(AC)$. For each line l parallel to $A'C'$ denote $D_l = l \cap A'D'$, $E_l = l \cap C'E'$.

Let us look at the line l sliding parallelly to $A'C'$ and prove that there is a unique position of this line such that $A'E_l$ is parallel to $B'D_l$. The existence side is easy: it is just the line $D'E'$ side of the regular pentagon. If we could prove that for every other position of l the lines $A'E_l$ and $B'D_l$ do intersect, we know that f takes $ABCDE$ to the regular pentagon $A'B'C'D'E'$ (as f should also preserve parallelism of AE and BD).

To show that the position of l such that $A'E_l$ is parallel to $B'D_l$ is unique, let us move the line l parallel to $A'C'$ starting from the line through $Y = A'D' \cap C'E'$ and moving away from AC . As we move, the line $A'E_l$ rotates about A' in one direction and the line $B'D_l$ rotates about B' in the other direction, and it is easy to see that $A'E_l$ does intersect $B'D_l$ when l is closer to Y than $D'E'$ is, as well as $A'E_l$ does intersect $B'D_l$ (on the other side with respect to l) when l lies further from Y than $D'E'$ does.

- 7.4. (*) (a) Use similarity of triangles (or any other arguments of affine geometry) to prove Theorem of Menelaus:

Given a triangle ABC , and a transversal line that crosses BC , AC and AB at points D , E and F respectively, with D , E , and F distinct from A , B and C , then

$$\frac{AF}{FB} \cdot \frac{BD}{DC} \cdot \frac{CE}{EA} = 1.$$

(Note that at least one of the sides will have to be extended to get the intersection point).

- (b) The theorem was known before Menelaus. Menelaus proved the spherical version of the theorem:

$$\frac{\sin AF}{\sin FB} \cdot \frac{\sin BD}{\sin DC} \cdot \frac{\sin CE}{\sin EA} = 1.$$

Use sine law to prove the spherical version of of Theorem of Menelaus.

- (a) **Solution.** Let l be the transversal line and let A' , B' and C' be the orthogonal projections of the points A, B, C to l . Then from three pairs of similar triangles we get

$$\frac{AF}{BF} = \frac{AA'}{BB'} \quad \frac{BD}{DC} = \frac{BB'}{CC'} \quad \frac{CE}{AE} = \frac{CC'}{AA'}.$$

Taking the product of the three values we get

$$\frac{AF}{BF} \cdot \frac{BD}{DC} \cdot \frac{CE}{AE} = \frac{AA'}{BB'} \cdot \frac{BB'}{CC'} \cdot \frac{CC'}{AA'} = 1.$$

Another solution: Consider the three homothecies f, g, h with centers D, E, F that respectively send B to C , C to A , and A to B . The composition $h \circ g \circ f$ then is an element of the group of homothecy-translations that fixes B , so it is a homothecy with center B , possibly with ratio 1 (in which case it is the identity). This composition $h \circ g \circ f$ fixes the line DE if and only if F is collinear to D and E (since the first two homothecies certainly fix DE , and the third does so only if F lies on DE). Therefore, D, E, F are collinear if and only if this composition is the identity, which means that the product of the following three ratios is 1:

$$\frac{DC}{DB} = \frac{EA}{AC} = \frac{FB}{FA} = 1.$$

- (b) Using sine law for triangles AFE, BFD and CED respectively we get

$$\frac{\sin AF}{\sin \angle E} = \frac{\sin AE}{\sin \angle F}, \quad \frac{\sin BF}{\sin \angle D} = \frac{\sin BD}{\sin \angle F}, \quad \frac{\sin CE}{\sin \angle D} = \frac{\sin CD}{\sin \angle E},$$

(here we use that $\sin AEF = \sin DEC$). Rewriting the three equations we get

$$\frac{\sin AF}{\sin AE} = \frac{\sin \angle E}{\sin \angle F}, \quad \frac{\sin BD}{\sin BF} = \frac{\sin \angle F}{\sin \angle D}, \quad \frac{\sin CE}{\sin CD} = \frac{\sin \angle D}{\sin \angle E}.$$

Multiplying the three equations we get the required identity.

- 7.5. Three pegs on a plane form an isosceles right triangle with a leg of length 3. The pegs may move to an arbitrary distance but on a line parallel to the line formed by the other two. Is it possible to eventually get the three pegs at the vertices of a right triangle with legs 2 and 4?

Solution. The admissible transformation on the triple of pegs preserves the area of the triangle. As $3 \cdot 3/2 \neq 2 \cdot 4/2$, no composition of these transformation will take the initial triangle to the right triangle with legs 2 and 4.

- 7.6. (a) Find a projective transformation f which takes the points $0, 1, \infty$ of the projective line to the points $3, 4, 0$ respectively.
 (b) Find the image of the point 2 under the transformation f in part (a) and use it to check that f preserves the cross-ratio of the points $0, 1, 2, \infty$.

Solution. (a) We will find f as a linear-fractional transformation $f = \frac{ax+b}{cx+d}$, $a, b, c, d \in \mathbb{R}$:

Since $f(\infty) = 0$, we have $a = 0$.

Since $f(0) = 3$, we have $b/d = 3$.

Since $f(1) = 4$, we have $b/(c+d) = 4$. We can choose $b = 12$, then $d = 4$, $c = -1$.

So $f(x) = \frac{12}{-x+4}$.

(b) $f(2) = 6$, $[0, 1, 2, \infty] = \frac{2}{1/\infty} = 2$; $[f(0), f(1), f(2), f(\infty)] = [3, 4, 6, 0] = \frac{3}{2}/\frac{-3}{-4} = 2$.

- 7.7. Let $A, B, C, D \in \mathbb{R}^2$ be four collinear points and O be a point not on the same line. Suppose that OB is a median of AOC and OC is a median of BOD . Find the cross-ratio of the lines OA, OB, OC and OD .

Solution. The cross-ratio of four lines is the cross-ratio of their intersection points with any given line l , in particular, it coincides with $[A, B, C, D] = \frac{CA}{CB}/\frac{DA}{DB} = \frac{2}{1}/\frac{3}{2} = 4/3$.

- 7.8. (*) (a) Show that $[A, B, C, D] = [C, D, A, B] = [B, A, D, C] = [D, C, B, A]$.
 (b) Given $[A, B, C, D] = \lambda$ find $[A, B, D, C]$ and $[A, C, B, D]$.
 (c) Assuming that $[A, B, C, D] = \lambda$, find all other possible values for $[X_1, X_2, X_3, X_4]$, where (X_1, X_2, X_3, X_4) is a permutation of (A, B, C, D) .

Solution. (a) The first two equalities may be checked directly (you get the same multiples in the other order). The third follows from the first two.

(b) $[A, B, D, C] = \frac{d-a}{d-b}/\frac{c-a}{c-b} = \frac{1}{[A, B, C, D]} = 1/\lambda$.

$$\begin{aligned} [A, C, B, D] &= \frac{b-a}{b-c}/\frac{d-a}{d-c} = \frac{(b-a)(d-c)}{(b-c)(d-a)} = \frac{((b-c) - (a-c))((d-b) - (c-b))}{(b-c)(d-a)} = \\ &= -\frac{(a-c)(d-b)}{(b-c)(d-a)} + \frac{(b-c)((d-b) - (c-b) - (a-c))}{(b-c)(d-a)} = -\lambda + 1, \end{aligned}$$

since $(d-b) - (c-b) - (a-c) = (d-a)$.

(c) In (b) we have seen that together with λ one gets $1/\lambda$ and $1 - \lambda$ (we will denote these operations by “ $-$ ” and “ \div ”). We may apply these transformations repeatedly (but it does not make sense to apply the same transformation twice in a row - we get back to λ). So, applying these transformations one after another we get

$$\lambda \xleftrightarrow{-} 1 - \lambda \xleftrightarrow{\div} \frac{1}{1 - \lambda} \xleftrightarrow{-} \frac{-\lambda}{1 - \lambda} \xleftrightarrow{\div} \frac{1 - \lambda}{-\lambda} \xleftrightarrow{-} \frac{1}{\lambda} \xleftrightarrow{\div} \lambda,$$

which gives 6 different values. Taking in account that each value works for 4 permutations (see (a)) and that 24 is the number of all permutations of 4 objects, we see that the six values is the complete list.

So, all possible values are $\lambda, 1 - \lambda, \frac{1}{1-\lambda}, \frac{\lambda}{1-\lambda}, \frac{1-\lambda}{\lambda}, \frac{1}{\lambda}$.

- 8.1. Find the projective transformation of \mathbb{RP}^2 sending the points $(1 : 0 : 0), (0 : 1 : 0), (1 : 0 : 1)$ and $(3 : -1 : 2)$ to the points $(2 : -1 : 1), (0 : 1 : 1), (1 : 0 : 2), (0 : 0 : 1)$.

Solution. First, we will find a projective transformation f mapping the points $(1 : 0 : 0), (0 : 1 : 0), (1 : 0 : 1)$ and $(3 : -1 : 2)$ to $(1 : 0 : 0), (0 : 1 : 0), (0 : 0 : 1), (1 : 1 : 1)$.

Let $f(v) = Av$, where $A \in GL(3, \mathbb{R})$, $v = (v, y, x)$. Since

$$\begin{pmatrix} a & b & c \\ d & e & f \\ g & h & j \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} a \\ d \\ g \end{pmatrix},$$

we see that $d = g = 0$. Similarly applying A to $(0, 1, 0)$ we see that $b = h = 0$.

Applying A to $(1, 0, 1)$ we get $(a + c, f, j)$ which should be proportional to $(0, 0, 1)$, so $c = -a$, $f = 0$.

Next, Applying A to $(3, -1, 2)$ we get $(a, -e, 2j)$, which should be proportional to $(1, 1, 1)$. So, $a = -e = 2j$ and A may be written as

$$A = \begin{pmatrix} 2 & 0 & -2 \\ 0 & -2 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Next, we will find a projective transformation g mapping the points $(1 : 0 : 0), (0 : 1 : 0), (0 : 0 : 1), (1 : 1 : 1)$ to the points $(2 : -1 : 1), (0 : 1 : 1), (1 : 0 : 2), (0 : 0 : 1)$.

Let $g(v) = Bv$, $B \in GL(3, \mathbb{R})$. Applying B to the first three vectors, it is easy to see that

$$B = \begin{pmatrix} 2 \cdot k & 0 \cdot l & 1 \cdot n \\ -1 \cdot k & 1 \cdot l & 0 \cdot n \\ 1 \cdot k & 1 \cdot l & 2 \cdot n \end{pmatrix}$$

for some $k, l, n \in \mathbb{R}$ (columns proportional to the vectors we want to get).

Also, applying B to the vector $(1, 1, 1)$ we want to get a vector proportional to $(0, 0, 1)$, which gives $k = l = -n/2$ and B may be written as

$$B = \begin{pmatrix} 2 & 0 & -2 \\ -1 & 1 & 0 \\ 1 & 1 & -4 \end{pmatrix}.$$

Finally, we obtain the transformation as

$$BA = \begin{pmatrix} 2 & 0 & -2 \\ -1 & 1 & 0 \\ 1 & 1 & -4 \end{pmatrix} \begin{pmatrix} 2 & 0 & -2 \\ 0 & -2 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 4 & 0 & -6 \\ -2 & -2 & 2 \\ 2 & -2 & -6 \end{pmatrix}.$$

Remark. While searching for the matrix A we have actually used that the set of initial vectors was quite similar to what we wanted to obtain (so that the matrix A is upper triangular, with many zeros). In general, one should first search for A^{-1} , matrix taking $(1 : 0 : 0), (0 : 1 : 0), (0 : 0 : 1), (1 : 1 : 1)$ to the initial vectors (this is easy, its columns are proportional to the first three vectors), and then to find A as the inverse to A^{-1} .

- 8.2. How many projective transformations send a quadrilateral to itself?

Solution. Let $ABCD$ be a quadrilateral. For each permutation σ of the vertices A, B, C, D there exists a unique projective transformation which sends $ABCD$ to $\sigma(A)\sigma(B)\sigma(C)\sigma(D)$. Since there are 24 permutations on the 4-element set, there are 24 projective maps taking the quadrilateral to itself.

- 8.3. (*) Calculate the cross-ratio of the following four points lying on the infinite line: $(1; 2; 0)$, $(2; 3; 0)$, $(3; 4; 0)$, $(4; 1; 0)$.

Solution. The cross-ratio of four collinear points in \mathbb{RP}^2 is the cross-ratio of the corresponding four (coplanar) lines in \mathbb{R}^3 (the point $(x; y; z)$ corresponds to the line in \mathbb{R}^3 passing through the origin and (x, y, z)).

To find the cross-ratio of the four lines (all lying in the plane $z = 0$) we find the intersections of these lines with some line. Let l be the line given by $z = 0, y = 12$. Then the four lines intersect l at points $x = 6, 8, 9, 48$ respectively.

So, the cross-ratio is $\frac{3}{1} / \frac{42}{40} = 20/7$.