

Denote by \mathcal{F} the space of all functions from phase space \mathcal{P} to \mathbb{R} . Given any function $f \in \mathcal{F}$, we can define an operator Φ_f that generates infinitesimal transformations on \mathcal{F} using the Poisson bracket.

Definition 5.2.8. The *Hamiltonian flow* defined by $f: \mathcal{P} \rightarrow \mathbb{R}$ is the infinitesimal transformation on \mathcal{F} defined by

$$\begin{aligned}\Phi_f^{(\epsilon)}: \mathcal{F} &\rightarrow \mathcal{F} \\ \Phi_f^{(\epsilon)}(g) &= g + \epsilon\{g, f\} + \mathcal{O}(\epsilon^2).\end{aligned}$$

Remark 5.2.9. I am taking a small liberty with the language here to avoid having to introduce some additional formalism: what I have just introduced is the infinitesimal version of what is commonly known as “Hamiltonian flow” in the literature, which is typically defined for finite (that is, non-infinitesimal) transformations. The finite version of the transformation is obtained by exponentiation:

$$\Phi_f^{(a)}(g) := e^{a\{\cdot, f\}}g = g + a\{g, f\} + \frac{a^2}{2!}\{\{g, f\}, f\} + \frac{a^3}{3!}\{\{\{g, f\}, f\}, f\} + \dots$$

Remark 5.2.10. By studying the action of $\Phi_f^{(\epsilon)}$ on the coordinates \mathbf{q}, \mathbf{p} of phase space, we can also understand $\Phi_f^{(\epsilon)}$ as the generator of a map from phase space to itself. We have

$$\begin{aligned}\Phi_f^{(\epsilon)}(q_i) &= q_i + \epsilon\{q_i, f\} + \mathcal{O}(\epsilon^2) = q_i + \epsilon\frac{\partial f}{\partial p_i} + \mathcal{O}(\epsilon^2) \\ \Phi_f^{(\epsilon)}(p_i) &= p_i + \epsilon\{p_i, f\} + \mathcal{O}(\epsilon^2) = p_i - \epsilon\frac{\partial f}{\partial q_i} + \mathcal{O}(\epsilon^2).\end{aligned}$$

The two definitions are compatible:


$$\begin{aligned}\Phi_f^{(\epsilon)}(g) &= g(q_1 + \epsilon\{q_1, f\}, \dots, q_n + \epsilon\{q_n, f\}, p_1 + \epsilon\{p_1, f\}, \dots, p_n + \epsilon\{p_n, f\}) \\ &= g(q_1, \dots, q_n, p_1, \dots, p_n) + \epsilon \sum_{i=1}^n \left(\frac{\partial g}{\partial q_i} \{q_i, f\} + \frac{\partial g}{\partial p_i} \{p_i, f\} \right) \\ &= g(q_1, \dots, q_n, p_1, \dots, p_n) + \epsilon \sum_{i=1}^n \left(\frac{\partial g}{\partial q_i} \frac{\partial f}{\partial p_i} - \frac{\partial g}{\partial p_i} \frac{\partial f}{\partial q_i} \right) \\ &= g + \epsilon\{g, f\}\end{aligned}$$

where in the second line we have done a Taylor expansion, and we have omitted higher order terms in ϵ throughout for notational simplicity.

Example 5.2.11. As a simple example, consider a particle moving in one dimension. The Hamiltonian flow Φ_p associated to the canonical momentum p acts on phase space functions as:

$$\Phi_p^{(\epsilon)}(g(q, p)) = g(q, p) + \epsilon\frac{\partial g}{\partial q} + \mathcal{O}(\epsilon^2).$$

Alternatively, $\Phi_p^{(\epsilon)}$ acts on the coordinate q as $q \rightarrow q + \epsilon$, so the effect of $\Phi_p^{(\epsilon)}$ on phase space is a uniform shift in the q direction:



We can reproduce the effect on arbitrary functions of q from this viewpoint by doing a Taylor expansion:

$$g(q + \epsilon, p) = g(q, p) + \epsilon \frac{\partial g}{\partial q} + \mathcal{O}(\epsilon^2).$$

(You might also find it interesting to reproduce the full form of the Taylor expansion of $f(x + a)$ around x using the exponentiated version in remark 5.2.9.)

Example 5.2.12. As a second example, consider a particle of unit mass moving in two dimensions, expressed in Cartesian coordinates, which we call q_1 and q_2 . We choose the Lagrangian to be of the form

$$L = \frac{1}{2}(\dot{q}_1^2 + \dot{q}_2^2) - V(q_1, q_2).$$


For the function generating the flow we will choose $J = q_1\dot{q}_2 - q_2\dot{q}_1$. (Recall from example 4.1.16 that this function is angular momentum, which Noether's theorem associated with rotations around the origin.) From the Lagrangian we have $p_1 = \dot{q}_1$ and $p_2 = \dot{q}_2$, so in terms of standard (\mathbf{q}, \mathbf{p}) coordinates of phase space we have $J(\mathbf{q}, \mathbf{p}) = q_1p_2 - q_2p_1$. The Hamiltonian flow $\Phi_J^{(\epsilon)}$ then acts on phase space as

$$\begin{aligned} \Phi_J^{(\epsilon)}(q_1) &= q_1 + \epsilon\{q_1, J\} = q_1 + \epsilon \frac{\partial J}{\partial p_1} = q_1 - \epsilon q_2, \\ \Phi_J^{(\epsilon)}(q_2) &= q_2 + \epsilon\{q_2, J\} = q_2 + \epsilon \frac{\partial J}{\partial p_2} = q_2 + \epsilon q_1, \\ \Phi_J^{(\epsilon)}(p_1) &= p_1 + \epsilon\{p_1, J\} = p_1 - \epsilon \frac{\partial J}{\partial q_1} = p_1 - \epsilon p_2, \\ \Phi_J^{(\epsilon)}(p_2) &= p_2 + \epsilon\{p_2, J\} = p_2 - \epsilon \frac{\partial J}{\partial q_2} = p_2 + \epsilon p_1. \end{aligned}$$

omitting higher orders in ϵ . So the effect of J on the coordinates can be written as an infinitesimal rotation on the \mathbf{q} and the \mathbf{p} (independently)

$$\begin{aligned} \Phi_J^{(\epsilon)} \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} &= \begin{pmatrix} 1 & -\epsilon \\ \epsilon & 1 \end{pmatrix} \begin{pmatrix} q_1 \\ q_2 \end{pmatrix}, \\ \Phi_J^{(\epsilon)} \begin{pmatrix} p_1 \\ p_2 \end{pmatrix} &= \begin{pmatrix} 1 & -\epsilon \\ \epsilon & 1 \end{pmatrix} \begin{pmatrix} p_1 \\ p_2 \end{pmatrix}. \end{aligned}$$

For instance, the action of $\Phi_J^{(\epsilon)}$ on the (q_1, q_2) slice of phase space (which in this case has four dimensions) is as in the following picture:



§5.2.1 Flows for conserved charges

We have just seen that linear momentum p generates spatial translations, and angular momentum generates rotations. This is in fact general: assume that we have a transformation acting as $q_i \rightarrow q_i + \epsilon a_i(\mathbf{q}) + O(\epsilon^2)$ on the generalised coordinates. Noether's theorem assigns a charge to this transformation given, in the Lagrangian framework, by

$$Q(\mathbf{q}, \dot{\mathbf{q}}, t) = \left(\sum_{i=1}^n a_i(\mathbf{q}) \frac{\partial L(\mathbf{q}, \dot{\mathbf{q}}, t)}{\partial \dot{q}_i} \right) - F(\mathbf{q}, t).$$

This charge can be written in the Hamiltonian framework in terms of generalised coordinates and generalised momenta as

$$Q(\mathbf{q}, \mathbf{p}, t) = \left(\sum_{i=1}^n a_i(\mathbf{q}) p_i \right) - F(\mathbf{q}, t).$$

If we now compute the Hamiltonian flow associated to this charge on the generalised coordinates we find

$$\Phi_Q^{(\epsilon)}(q_i) = q_i + \epsilon \{q_i, Q\} + O(\epsilon^2) = q_i + \epsilon a_i + O(\epsilon^2).$$

Note 5.2.13

This is a very important result: Noether's theorem told us that symmetries imply the existence of conserved quantities. We have just seen that we can go in the other direction too: conserved quantities *generate* the corresponding symmetry transformations, via the associated Hamiltonian flow.