Denote by \mathscr{F} the space of all functions from phase space \mathscr{P} to \mathbb{R} . Given any function $f \in \mathscr{F}$, we can define an operator Φ_f that generates infinitesimal transformations on \mathscr{F} using the Poisson bracket.

Definition 6.2.8. The *Hamiltonian flow* defined by $f: \mathscr{P} \to \mathbb{R}$ is the infinitesimal transformation on \mathscr{F} defined by

$$\Phi_f^{(\epsilon)} \colon \mathscr{F} \to \mathscr{F}$$

$$\Phi_f^{(\epsilon)}(g) = g + \epsilon \{g, f\} + \mathcal{O}(\epsilon^2) \,.$$

Remark 6.2.9. I am taking a small liberty with the language here to avoid having to introduce some additional formalism: what I have just introduced is the infinitesimal version of what is commonly known as "Hamiltonian flow" in the literature, which is typically defined for finite (that is, non-infinitesimal) transformations. The finite version of the transformation is obtained by exponentiation:

$$\Phi_f^{(a)}(g) = e^{a\{\cdot,f\}}g \coloneqq g + a\{g,f\} + \frac{a^2}{2!}\{\{g,f\},f\} + \frac{a^3}{3!}\{\{\{g,f\},f\},f\} + \dots$$

Remark 6.2.10. By studying the action of $\Phi_f^{(\epsilon)}$ on the coordinates \mathbf{q} , \mathbf{p} of phase space, we can also understand $\Phi_f^{(\epsilon)}$ as the generator of a map from phase space to itself. We have

$$\Phi_f^{(\epsilon)}(q_i) = q_i + \epsilon \{q_i, f\} + O(\epsilon^2) = q_i + \epsilon \frac{\partial f}{\partial p_i} + O(\epsilon^2)$$

$$\Phi_f^{(\epsilon)}(p_i) = p_i + \epsilon \{p_i, f\} + O(\epsilon^2) = p_i - \epsilon \frac{\partial f}{\partial q_i} + O(\epsilon^2)$$

The two definitions are compatible:

$$\begin{split} \Phi_f^{(\epsilon)}(g) &= g(q_1 + \epsilon\{q_1, f\}, \dots, q_n + \epsilon\{q_n, f\}, p_1 + \epsilon\{p_1, f\}, \dots, p_n + \epsilon\{p_n, f\}) \\ &= g(q_1, \dots, q_n, p_1, \dots, p_n) + \epsilon \sum_{i=1}^n \left(\frac{\partial g}{\partial q_i}\{q_i, f\} + \frac{\partial g}{\partial p_i}\{p_i, f\}\right) \\ &= g(q_1, \dots, q_n, p_1, \dots, p_n) + \epsilon \sum_{i=1}^n \left(\frac{\partial g}{\partial q_i}\frac{\partial f}{\partial p_i} - \frac{\partial g}{\partial p_i}\frac{\partial f}{\partial q_i}\right) \\ &= g + \epsilon\{g, f\} \end{split}$$

where in the second line we have done a Taylor expansion, and we have omitted higher order terms in ϵ throughout for notational simplicity.

Example 6.2.11. As a simple example, consider a particle moving in one dimension. The Hamiltonian flow Φ_p associated to the canonical momentum p acts on phase space functions as:

$$\Phi_p^{(\epsilon)}(g(q,p)) = g(q,p) + \epsilon \frac{\partial g}{\partial q} + \mathcal{O}(\epsilon^2) \,.$$

Alternatively, $\Phi_p^{(\epsilon)}$ acts on the coordinate q as $q \to q + \epsilon$, so the effect of $\Phi_p^{(\epsilon)}$ on phase space is a uniform shift in the q direction:



We can reproduce the effect on arbitrary functions of q from this viewpoint by doing a Taylor expansion:

$$g(q + \epsilon, p) = g(q, p) + \epsilon \frac{\partial g}{\partial q} + \mathcal{O}(\epsilon^2).$$

(You might also find it interesting to reproduce the full form of the Taylor expansion of f(x + a) around x using the exponentiated version in remark 6.2.9.)

Example 6.2.12. As a second example, consider a particle of unit mass moving in two dimensions, expressed in Cartesian coordinates, which we call q_1 and q_2 . We choose the Lagrangian to be of the form

$$L = \frac{1}{2}(\dot{q}_1^2 + \dot{q}_2^2) - V(q_1, q_2).$$

For the function generating the flow we will choose $J = q_1\dot{q}_2 - q_2\dot{q}_1$. (Recall from example 3.1.16 that this function is angular momentum, which Noether's theorem associated with rotations around the origin.) From the Lagrangian we have $p_1 = \dot{q}_1$ and $p_2 = \dot{q}_2$, so in terms of standard (\mathbf{q}, \mathbf{p}) coordinates of phase space we have $J(\mathbf{q}, \mathbf{p}) = q_1p_2 - q_2p_1$. The Hamiltonian flow $\Phi_J^{(\epsilon)}$ then acts on phase space as

$$\Phi_{J}^{(\epsilon)}(q_{1}) = q_{1} + \epsilon \{q_{1}, J\} = q_{1} + \epsilon \frac{\partial J}{\partial p_{1}} = q_{1} - \epsilon q_{2},$$

$$\Phi_{J}^{(\epsilon)}(q_{2}) = q_{2} + \epsilon \{q_{2}, J\} = q_{2} + \epsilon \frac{\partial J}{\partial p_{2}} = q_{2} + \epsilon q_{1},$$

$$\Phi_{J}^{(\epsilon)}(p_{1}) = p_{1} + \epsilon \{p_{1}, J\} = p_{1} - \epsilon \frac{\partial J}{\partial q_{1}} = p_{1} - \epsilon p_{2},$$

$$\Phi_{J}^{(\epsilon)}(p_{2}) = p_{2} + \epsilon \{p_{2}, J\} = p_{2} - \epsilon \frac{\partial J}{\partial q_{2}} = p_{2} + \epsilon p_{1}.$$

omitting higher orders in ϵ . So the effect of J on the coordinates can be written as an infinitesimal rotation on the \mathbf{q} and the \mathbf{p} (independently)

$$\Phi_J^{(\epsilon)}\begin{pmatrix} q_1\\q_2 \end{pmatrix} = \begin{pmatrix} 1 & -\epsilon\\\epsilon & 1 \end{pmatrix}\begin{pmatrix} q_1\\q_2 \end{pmatrix},$$

$$\Phi_J^{(\epsilon)}\begin{pmatrix} p_1\\p_2 \end{pmatrix} = \begin{pmatrix} 1 & -\epsilon\\\epsilon & 1 \end{pmatrix}\begin{pmatrix} p_1\\p_2 \end{pmatrix}.$$

For instance, the action of $\Phi_J^{(\epsilon)}$ on the (q_1, q_2) slice of phase space (which in this case has four dimensions) is as in the following picture:



§6.2.1 Flows for conserved charges

We have just seen that linear momentum p generates spatial translations, and angular momentum generates rotations. This is in fact general: assume that we have a transformation acting as $q_i \rightarrow q_i + \epsilon a_i(\mathbf{q}) + O(\epsilon^2)$ on the generalised coordinates. Noether's theorem assigns a charge to this transformation given, in the Lagrangian framework, by

$$Q(\mathbf{q}, \dot{\mathbf{q}}, t) = \left(\sum_{i=1}^{n} a_i(\mathbf{q}) \frac{\partial L(\mathbf{q}, \dot{\mathbf{q}}, t)}{\partial \dot{q}_i}\right) - F(\mathbf{q}, t) \,.$$

This charge can be written in the Hamiltonian framework in terms of generalised coordinates and generalised momenta as

$$Q(\mathbf{q}, \mathbf{p}, t) = \left(\sum_{i=1}^{n} a_i(\mathbf{q}) p_i\right) - F(\mathbf{q}, t) \,.$$

If we now compute the Hamiltonian flow associated to this charge on the generalised coordinates we find

$$\Phi_Q^{(\epsilon)}(q_i) = q_i + \epsilon \{q_i, Q\} + O(\epsilon^2) = q_i + \epsilon a_i + \mathcal{O}(\epsilon^2) \,.$$

Note 6.2.13

This is a very important result: Noether's theorem told us that symmetries imply the existence of conserved quantities. We have just seen that we can go in the other direction too: conserved quantities *generate* the corresponding symmetry transformations, via the associated Hamiltonian flow.