

§6.4 There and back again

i This section is *not examinable*. **i**

Let me finish by closing the circle of ideas that we have been developing. We have seen in §3.1 that Noether's theorem says that every symmetry implies the existence of a conserved Noether charge Q , and we have also shown in §6.2.1 that the Noether charge associated to a symmetry generates the right transformations on the generalized coordinates. It is natural to ask at this point: does any conserved charge generate a symmetry transformation? This is indeed that case, as we now show for the class of conserved charges that we have been discussing.

Theorem 6.4.1. *Assume that we have a function $Q(\mathbf{q}, \mathbf{p}, t)$ of the form*

$$Q(\mathbf{q}, \mathbf{p}, t) = \left(\sum_{i=1}^n a_i(\mathbf{q}) p_i \right) - F(\mathbf{q}, t)$$

such that

$$\frac{dQ}{dt} = \{Q, H\} + \frac{\partial Q}{\partial t} = 0,$$

Then $\Phi_Q^{(\epsilon)}(L) = L + \epsilon \frac{dF}{dt} + O(\epsilon^2)$, so Q generates a symmetry, whose Noether charge is Q .

Proof. Let me start by proving some simple auxiliary results. Note first that

$$\{q_i, Q\} = \frac{\partial Q}{\partial p_i} = a_i(\mathbf{q}) \quad (6.4.1)$$

which implies, in particular, that

$$\frac{\partial \{q_i, Q\}}{\partial t} = \frac{\partial a_i(\mathbf{q})}{\partial t} = 0.$$

Note also that since Q is conserved, and the only explicit time dependence of Q on time is via F , we have

$$\{Q, H\} = \frac{dQ}{dt} - \frac{\partial Q}{\partial t} = -\frac{\partial Q}{\partial t} = \frac{\partial F}{\partial t},$$

so

$$-\{\{Q, H\}, q_i\} = \{q_i, \{Q, H\}\} = \frac{\partial \{Q, H\}}{\partial p_i} = \frac{\partial}{\partial p_i} \left(\frac{\partial F(\mathbf{q}, t)}{\partial t} \right) = 0.$$

Using these two results we find that

$$\begin{aligned} \frac{d\{q_i, Q\}}{dt} &= \frac{\partial \{q_i, Q\}}{\partial t} + \{\{q_i, Q\}, H\} \\ &= \{\{q_i, Q\}, H\} \\ &= -\{\{H, q_i\}, Q\} - \{\{Q, H\}, q_i\} \\ &= \{\{q_i, H\}, Q\}. \end{aligned}$$

where on the third line we have used the Jacobi identity in proposition 6.2.7. Hamilton's equations then imply that

$$\frac{d\{q_i, Q\}}{dt} = \{\dot{q}_i, Q\}. \quad (6.4.2)$$

Using these results it is straightforward to compute the change in the Lagrangian due to Q . The Lagrangian in Hamiltonian coordinates is

$$L(\mathbf{q}, \mathbf{p}, t) = \left(\sum_{i=1}^n \dot{q}_i(\mathbf{q}, \mathbf{p}) p_i \right) - H(\mathbf{q}, \mathbf{p}, t).$$

Since Q is conserved, we have

$$\begin{aligned} \{L, Q\} &= \left[\sum_{i=1}^n (\{\dot{q}_i, Q\} p_i + \dot{q}_i \{p_i, Q\}) \right] + \{Q, H\} \\ &= \left[\sum_{i=1}^n (\{\dot{q}_i, Q\} p_i + \dot{q}_i \{p_i, Q\}) \right] - \frac{\partial Q}{\partial t}, \end{aligned}$$

which becomes, using the results above:

$$\begin{aligned} \{L, Q\} &= \left[\sum_{i=1}^n \left(p_i \frac{d}{dt} \{q_i, Q\} + \dot{q}_i \{p_i, Q\} \right) \right] - \frac{\partial Q}{\partial t} \\ &= \left[\sum_{i=1}^n \left(\frac{d}{dt} (\{q_i, Q\} p_i) - \dot{p}_i \{q_i, Q\} + \dot{q}_i \{p_i, Q\} \right) \right] - \frac{\partial Q}{\partial t} \\ &= \left[\sum_{i=1}^n \left(\frac{d}{dt} (\{q_i, Q\} p_i) - \dot{p}_i \frac{\partial Q}{\partial p_i} - \dot{q}_i \frac{\partial Q}{\partial q_i} \right) \right] - \frac{\partial Q}{\partial t} \\ &= \frac{d}{dt} \left(\sum_{i=1}^n \{q_i, Q\} p_i \right) - \frac{dQ}{dt}. \end{aligned}$$

where on the first line we have used (6.4.2), and in going from the third to the fourth the Chain Rule. Finally, using (6.4.1) we find that

$$\sum_{i=1}^n \{q_i, Q\} p_i = \sum_{i=1}^n a_i p_i = Q + F$$

which implies

$$\{L, Q\} = \frac{d(Q + F - Q)}{dt} = \frac{dF}{dt}.$$

Recalling the definition of the Hamiltonian flow operator, this gives

$$\Phi_Q^{(\epsilon)}(L) = L + \epsilon \{L, Q\} + \mathcal{O}(\epsilon^2) = L + \epsilon \frac{dF}{dt} + \mathcal{O}(\epsilon^2).$$

□