

## Week 9 problems

---

1. For a string obeying the wave equation  $u_{tt} = u_{xx}$  (so we set  $c^2 = 1$ ), we define a quantity

$$X(a, b) = \int_a^b ((u_t)^3 + 3u_t(u_x)^2) dx$$

where  $u_t = \frac{\partial u}{\partial t}$  and  $u_x = \frac{\partial u}{\partial x}$ . Use the wave equation to show that

$$\frac{d}{dt} X(a, b) = [g(u, u_t, u_x)]_a^b$$

for some function  $g(u, u_t, u_x)$  which you should determine. Show that if  $u_t$  and  $u_x$  vanish at  $x = \pm\infty$ , then  $X(-\infty, \infty)$  is a conserved quantity.

2. The motion of a string is governed by the wave equation  $u_{tt} = u_{xx}$  (so we are setting  $c^2 = 1$ ). The energy of the portion of the string between  $x = a$  and  $x = b$  is given by

$$E(a, b) = \frac{1}{2} \int_a^b \left( \left( \frac{\partial u}{\partial t} \right)^2 + \left( \frac{\partial u}{\partial x} \right)^2 \right) dx.$$

- (a) Use the wave equation to show that

$$\frac{d}{dt} E(a, b) = \left[ \frac{\partial u}{\partial t} \frac{\partial u}{\partial x} \right]_a^b.$$


- (b) In the case that the solution  $u$  is of the form  $u = f(x+t) - f(x-t)$ , find the form of the energy in terms of  $f$ .  
 (c) Compute what  $E(0, \infty)$  is in the case

$$u = \exp\left(-\frac{(x+t)^2}{2}\right) - \exp\left(-\frac{(x-t)^2}{2}\right)$$

You should get that  $E(0, \infty)$  a constant, whose value you should determine. (*It may be of help to recall that  $\int_{-\infty}^{\infty} e^{-y^2} dy = \sqrt{\pi}$ .*)

- (d) Use part (a) of this question to argue that  $E(0, \infty)$  is a constant whenever  $f$  is even or odd, and is such that  $\lim_{x \rightarrow \infty} f'(x) = 0$ .

3. The Dirichlet and Neumann boundary conditions studied in the lectures are those in which the energy flowing into the boundary vanishes. A simple system in which energy can flow into the boundary can be constructed by assuming that we attach a spring to the one-dimensional string at the  $x = 0$  boundary, as in the figure. The spring has constant  $\kappa$  and zero natural length, and it stores energy due to its extension, which is given by  $u(0, t)$ .



- (a) Write the equation encoding conservation of energy at the boundary.
- (b) Assuming the existence of a solution of the form

$$u(x, t) = \Re \left( (e^{ipx} + Re^{-ipx})e^{-ipct} \right)$$

for the incident and reflected monochromatic waves, solve for  $R$  (as a function of  $p$ ,  $\kappa$  and  $\tau$ ) by imposing energy conservation. [*Hint: as a consistency check, in the  $\kappa \rightarrow \infty$  limit the incoming wave will not have enough energy to excite the spring, so in this limit you should have  $R = -1$ , the result for Dirichlet boundary conditions (recall problem 6 in the Week 8 problem sheet). Similarly in the  $\kappa \rightarrow 0$  limit you should reproduce the result of Neumann boundary conditions,  $R = 1$ .]*