

Mostow's Lattices and Cone Metrics on the Sphere

Richard K. Boadi,
Department of Mathematics,
Kwame Nkrumah University of Science and Technology,
Kumasi, Ghana
email: richardkbgh@hotmail.com

John R. Parker,
Department of Mathematical Sciences,
Durham University,
Durham DH1 3LE, U.K.
email: j.r.parker@durham.ac.uk

Abstract

In [8] Mostow constructed a family of lattices in $\mathrm{PU}(2, 1)$, the holomorphic isometry group of complex hyperbolic 2-space. These groups are special cases of the lattices constructed by Deligne and Mostow [2] using monodromy of hypergeometric functions. Thurston [12] reinterpreted the work of Deligne and Mostow in terms of cone metrics on the sphere. In this paper we use Thurston's point of view to give a direct construction of fundamental domains for Mostow's lattices. Our approach is a direct generalisation of Parker's construction for Livné's lattices [10]. The details may be found in Boadi's PhD thesis [1]. We note that Deraux, Falbel and Paupert [3] have also constructed fundamental domains for Mostow's groups. Our groups are different from the ones considered in the main part of [3].

1 Introduction

A (*Euclidean*) *cone metric* on the sphere is a flat Euclidean metric on the sphere with finitely many singularities. Each singularity locally looks like the apex of a cone and may be formed by identifying the sides of a sector. The angle between the sides of the sector, or equivalently at the apex of the cone, is the *cone angle*. If the cone angle at a vertex v is $2\pi - \alpha$ then α is

the *curvature* at v . Then the sum of the curvatures at all vertices must be 4π .

For example, a cube is a cone metric on the sphere with eight singularities. At each vertex three squares meet and so the cone angle is $3\pi/2$. Clearly the curvature at each vertex is $\pi/2$ and the sum over all eight vertices gives a total curvature of $8 \cdot \pi/2 = 4\pi$.

In [12] Thurston considered the following construction. Suppose we are given a fixed number of cone singularities with prescribed cone angles (chosen so the sum of the curvatures is 4π). Keeping these angles fixed but varying the location of the singularities gives a moduli space of cone metrics. One may move the cone points in this space along a non-trivial closed path in the moduli space (for example by performing a Dehn twist). In doing so one naturally obtains a modular group. The area of the cone metric is preserved by this group. Thurston observed that this area gives an indefinite Hermitian form and one may embed the (projectivised) moduli space into complex hyperbolic space. For certain good choices of cone angle, the resulting group is a complex hyperbolic lattice. (See Weber [13] for an alternative point of view.)

Thurston's construction is an alternative point of view on the lattices constructed by Deligne and Mostow [2] via monodromy of hypergeometric functions in several variables. Thurston's good choices of angles correspond to what Deligne and Mostow call ball N -tuples satisfying the condition ΣINT . Among these groups are two families, one constructed by Mostow in [8] and the other constructed by Livné in [7]. In [10], Parker constructed Livné's lattices from first principles using Thurston's construction and he then went on to use this explicit construction to build fundamental domains for these groups. In his PhD thesis [1], Boadi extended Parker's construction to include some of Mostow's lattices. This paper is an account of this construction. We refer to [1] for many of the details.

Specifically, we consider cone metrics on the sphere with five cone points with cone angles

$$(\pi - \theta + 2\phi, \pi + \theta, \pi + \theta, \pi + \theta, 2\pi - 2\theta - 2\phi). \quad (1)$$

The angles θ and ϕ satisfy $\theta > 0$, $\phi > 0$ and $\theta + \phi < \pi$. In particular, we consider the following values of θ and ϕ which Thurston showed yield discrete groups [12]:

θ	$2\pi/3$	$2\pi/3$	$2\pi/3$	$2\pi/4$	$2\pi/4$	$2\pi/5$	$2\pi/5$	$2\pi/6$	$2\pi/6$
ϕ	$\pi/4$	$\pi/5$	$\pi/6$	$\pi/3$	$\pi/4$	$\pi/2$	$\pi/3$	$\pi/2$	$\pi/3$

(2)

Note that Mostow only considered the groups with $\theta = 2\pi/p$ for $p = 3, 4, 5$. He also considered other values of ϕ and fundamental domains for these groups were constructed by Deraux, Falbel and Paupert [3] (see also Parker [11]). The groups with $\theta = 2\pi/5$ or $2\pi/6$ and $\phi = \pi/2$ are on Livné's list. In fact the group corresponding to $(\theta, \phi) = (2\pi/6, \pi/2)$ is the Eisenstein-Picard modular group considered by Falbel and Parker in [5]. Also, the group corresponding to $(\theta, \phi) = (2\pi/3, \pi/6)$ is the so called "sister" of the Eisenstein-Picard modular group considered by Zhao [14]. The fundamental domains we construct are different from the ones constructed in [5] and [14]. Further relationships between these groups can be found in Parker's survey paper [11].

Certain automorphisms of our cone metrics yielded unitary matrices R_1 , R_2 and A_1 . (The naming of these automorphisms follows Mostow, see his survey paper [9] for example.) Our goal is to show that the group Γ generated by these automorphisms is discrete. To do so, we construct a polyhedron D . The vertices of D come from degenerate cone metrics where some of the cone points have coalesced. The boundary of D is made up of sides contained in bisectors. The maps R_1 , R_2 and A_1 lead to side pairing maps for D . Then finally we prove in Theorem 5.1, using Poincaré's polyhedron theorem, that the group Γ generated by the side pairings of D is a discrete subgroup of $\text{PU}(1, 2)$ with fundamental domain D and presentation:

$$\Gamma = \left\langle J, P, R_1, R_2 : \begin{array}{l} J^3 = R_1^p = R_2^p = (P^{-1}J)^k = I, \\ R_2 = PR_1P^{-1} = JR_1J^{-1}, P = R_1R_2 \end{array} \right\rangle.$$

The integers p and k are defined by $\theta = 2\pi/p$ and $\phi = \pi/k$ where θ and ϕ are given in (2). Equivalently, in terms of R_1 , R_2 and A_1 the presentation is

$$\Gamma = \left\langle R_1, R_2, A_1 : \begin{array}{l} R_1^p = R_2^p = A_1^k = (R_1R_2A_1)^3 = I, \\ R_1R_2R_1 = R_2R_1R_2, R_1A_1 = A_1R_1 \end{array} \right\rangle.$$

These presentations should be compared to the discussion in Mostow [9], particularly equation (5.3) and page 244. One may also write Γ as a two generator group. For example, as Mostow observes, it is easy to see that R_1 and J will generate Γ (which Mostow calls Γ_μ) but then the presentation is not so clean.

The paper is arranged as follows: Section 2 looks at the construction giving the fundamental domain for the cone structure. Section 3 looks at the construction of bisectors and vertices; Section 4 looks at the construction of the complex hyperbolic polyhedron D which sets the stage for the final Section 5 which is a summary of the proof that D is a fundamental

polyhedron for the group. Details of the work can be found in Boadi's thesis [1].

2 Construction of the Polygons and Automorphisms

2.1 The Fundamental Domain for the Cone Structures




Figure 1: *The doubled pentagon. Note that the angles indicated are the cone angles, which are twice the internal angles of the pentagon. The octagon Π is obtained by cutting along the bold lines.*

We consider Euclidean cone metrics on the sphere with five cone points with cone angles given by (1) where θ and ϕ are given in the table (2). Our goal is to try to find a unified construction for all these angles (and to verify that these angles correspond to discrete groups). If we cut the sphere open along a path through the five cone points, we obtain a Euclidean polygon Π . Conversely, if we glue the sides of Π together, we can reconstruct our cone metric on the sphere. We give an explicit parametrisation of such polygons in terms of three complex parameters (z_1, z_2, z_3) . We show that, in terms of these parameters, the area of the polygon gives a Hermitian form of signature $(1, 2)$. Thurston [12] and Weber [13] describe different ways of doing this. We follow Parker's method from [10] which is different from the

methods of Thurston and Weber.

We begin by looking at the case where the cone manifold is the double of a Euclidean pentagon. Cut the pentagon along four of its sides, as in Figure 1, with the first cut at the cone point with angle $2\pi - 2\theta - 2\phi$, then moving along the boundary of the pentagon through the three cone points v_3 , v_2 and v_1 with cone angle $\pi + \theta$, ending at the cone point v_0 with cone angle $\pi - \theta + 2\phi$. When we cut the double pentagon this way, we get an octagon, which we call Π ; see Figure 2. This octagon has a reflection symmetry. Using this symmetry to identify the boundary points reconstructs the doubled pentagon with which we began.

We now show how to construct Π geometrically. We start with a big triangle T_3 with angles θ , $\pi - \theta - \phi$ and ϕ . This will only work when $\theta > 0$, $\phi > 0$ and $\theta + \phi < \pi$. We then take off two smaller triangles T_1 with angles ϕ , $\pi/2 + \theta/2 - \phi$ and $\pi/2 - \theta/2$; and T_2 with angles θ , $\pi/2 - \theta/2$ and $\pi/2 - \theta/2$. The corners of the triangles T_1 and T_3 with angles ϕ are the same. The corners of the triangles T_2 and T_3 with angles θ are the same. The base vectors of T_1 , T_2 and T_3 are $ie^{-i\phi}z_1$, $ie^{-i\phi}z_2$ and $ie^{-i\phi}z_3$. See Figure 2 for the construction.

The vertices of the triangle T_1 are as follows:

$$\begin{aligned} v_0 &= \frac{-i \sin \theta}{\sin(\theta + \phi)} z_3 + \frac{i \sin \theta}{\sin \phi + \sin(\theta - \phi)} z_1, \\ x_0 &= \frac{-i \sin \theta}{\sin(\theta + \phi)} z_3, \\ v_1 &= ie^{-i\phi} z_1 - \frac{i \sin \theta}{\sin(\theta + \phi)} z_3. \end{aligned}$$

The vertices of triangle T_2 are as follows:

$$\begin{aligned} v_2 &= -ie^{-i\phi} z_2 + \frac{i \sin \phi e^{-i\theta - i\phi}}{\sin(\theta + \phi)} z_3, \\ x_1 &= \frac{i \sin \phi e^{-i\theta - i\phi}}{\sin(\theta + \phi)} z_3, \\ v_3 &= -ie^{-i\theta - i\phi} z_2 + \frac{i \sin \phi e^{-i\theta - i\phi}}{\sin(\theta + \phi)} z_3. \end{aligned}$$




Figure 2: The octagon Π when z_1, z_2 and z_3 are all real. Here $\psi = (\pi - \theta)/2$. Note that the arrows are $z_j i e^{-i\phi}$.

The vertices of triangle T_3 are also as follows:

$$\begin{aligned}
 &0, \\
 x_1 &= \frac{i \sin \phi e^{-i\theta - i\phi}}{\sin(\theta + \phi)} z_3, \\
 x_0 &= \frac{-i \sin \theta}{\sin(\theta + \phi)} z_3.
 \end{aligned}$$

We have constructed a pentagon whose vertices are the vertex of T_3 and the two vertices of each of T_1 and T_2 not shared by one of the other triangles. This has one edge in common with each of T_1 and T_2 . Consider the edge of this pentagon joining the vertices of T_1 with angle $\pi + \theta/2 - \phi$ and the vertex of T_3 with angle $\pi - \theta - \phi$. Reflect the pentagon across this side to form an octagon, see Figure 2. The image of the triangle T_1 under this reflection will be a new triangle T_{-1} . Similarly, the images of T_2 and T_3 will be triangles T_{-2} and T_{-3} . The images of vertices v_1, v_2 and v_3 will be v_{-1}, v_{-2} and v_{-3} respectively. Our resulting octagon is preserved by reflection in the imaginary axis and we label its vertices so that this reflection interchanges

v_j and v_{-j} . Moreover, gluing points of the boundary Π to their image under this reflection reconstructs the doubled pentagon we begun with. Below are the vertices of the octagon.

$$\begin{aligned}
v_0 &= \frac{-i \sin \theta}{\sin(\theta + \phi)} z_3 + \frac{i \sin \theta}{\sin \phi + \sin(\theta - \phi)} z_1, \\
v_1 &= ie^{-i\phi} z_1 - \frac{i \sin \theta}{\sin(\theta + \phi)} z_3, \\
v_2 &= -ie^{-i\phi} z_2 + \frac{i \sin \phi e^{-i\theta - i\phi}}{\sin(\theta + \phi)} z_3, \\
v_3 &= -ie^{-i\theta - i\phi} z_2 + \frac{i \sin \phi e^{-i\theta - i\phi}}{\sin(\theta + \phi)} z_3, \\
v_{-1} &= ie^{i\phi} z_1 - \frac{i \sin \theta}{\sin(\theta + \phi)} z_3, \\
v_{-2} &= -ie^{i\phi} z_2 + \frac{i \sin \phi e^{i\theta + i\phi}}{\sin(\theta + \phi)} z_3, \\
v_{-3} &= -ie^{i\theta + i\phi} z_2 + \frac{i \sin \phi e^{i\theta + i\phi}}{\sin(\theta + \phi)} z_3.
\end{aligned}$$

The vertices of T_{-1} are v_0 , x_0 and v_{-1} ; the vertices of T_{-2} are v_{-2} , x_{-1} and v_{-3} and the vertices of T_3 are 0 , x_{-1} and x_{-2} . We remark that substituting $\phi = \pi/2$ we recover the octagon from [10].

In the above construction, the octagon was formed by cutting a doubled pentagon along four of its edges. We now consider how to build an octagon associated with a more general cone metric on the sphere with five cone points whose angles are given by (1). We do this by following the above construction but with complex parameters z_1 , z_2 , z_3 in place of the real parameters x_1 , x_2 , x_3 . A typical example of an octagon with complex parameters is shown in Figure 3. The triangles T_1 , T_{-1} , T_3 and T_{-3} share the vertex x_0 , the triangles T_2 and T_3 share the vertex x_1 and the triangles T_{-2} and T_{-3} share the vertex x_{-1} . The base of the triangle T_1 is still $v_1 - x_0 = ie^{-i\phi} z_1$, the base of T_2 is $x_1 - v_2 = ie^{-i\phi} z_2$ and the base of T_3 is $x_1 - x_0 = ie^{-i\phi} z_3$. In general these are no longer real multiples of each other. Similarly, the bases of T_{-1} , T_{-2} and T_{-3} are $v_{-1} - x_0 = ie^{i\phi} z_1$, $x_{-1} - v_{-2} = ie^{i\phi} z_2$ and $x_{-1} - x_0 = ie^{i\phi} z_3$ respectively. Note that v_{-j} is no longer the image of v_j under reflection in the imaginary axis.




Figure 3: The octagon Π with complex z_1, z_2 and z_3 . Again $\psi = (\pi - \theta)/2$ and the arrows are $z_j i e^{-i\phi}$.

Simple geometry shows that the areas of the triangles are as follows:

$$\begin{aligned} \text{Area}(T_1) &= \frac{\sin \theta \sin \phi}{2(\sin \phi + \sin(\theta - \phi))} |z_1|^2, \\ \text{Area}(T_2) &= \frac{1}{2} \sin \theta |z_2|^2, \\ \text{Area}(T_3) &= \frac{\sin \theta \sin \phi}{2 \sin(\theta + \phi)} |z_3|^2. \end{aligned}$$

When the parameters z_j are real it is easy to see that the area of octagon Π is

$$\text{Area}(\Pi) = 2\text{Area}(T_3) - 2\text{Area}(T_1) - 2\text{Area}(T_2).$$

When the z_j are complex, one can show by a simple cut and paste argument

that this formula still holds. Therefore:

$$\begin{aligned}
\text{Area}(\Pi) &= 2\text{Area}(T_3) - 2\text{Area}(T_1) - 2\text{Area}(T_2) \\
&= \sin \theta \left(-\frac{\sin \phi}{(\sin \phi + \sin(\theta - \phi))} |z_1|^2 - |z_2|^2 + \frac{\sin \phi}{\sin(\theta + \phi)} |z_3|^2 \right) \\
&= \sin \theta \begin{bmatrix} \bar{z}_1 & \bar{z}_2 & \bar{z}_3 \end{bmatrix} \begin{bmatrix} \frac{-\sin \phi}{(\sin \phi + \sin(\theta - \phi))} & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & \frac{\sin \phi}{\sin(\theta + \phi)} \end{bmatrix} \begin{bmatrix} z_1 \\ z_2 \\ z_3 \end{bmatrix} \\
&= \mathbf{z}^* H \mathbf{z},
\end{aligned}$$

where H is the Hermitian matrix:

$$H = \sin \theta \begin{bmatrix} -\sin \phi / (\sin \phi + \sin(\theta - \phi)) & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & \sin \phi / \sin(\theta + \phi) \end{bmatrix}. \quad (3)$$

We observe that the area gives a Hermitian form of signature (1,2) on \mathbb{C}^3 . This leads to a complex hyperbolic structure on the moduli space of such polygons. This is a special case of Proposition 3.3 of Thurston [12].

There is a natural way to construct a particular Euclidean cone manifold from Π . The following, σ_j , are edge pairing maps of Π , which are Euclidean isometries which preserve orientation and so are completely determined on each edge by their value on the vertices v_j, v_{j+1} . The maps are:

$$\begin{aligned}
\sigma_1(0) = 0, \quad \sigma_1(v_3) = v_{-3}; & \quad \sigma_2(v_3) = v_{-3}, \quad \sigma_1(v_2) = v_{-2}; \\
\sigma_3(v_2) = v_{-2}, \quad \sigma_3(v_1) = v_{-1}; & \quad \sigma_4(v_1) = v_{-1}, \quad \sigma_4(v_0) = v_0.
\end{aligned}$$

Let M be the Euclidean cone manifold given by identifying the edges of Π using the maps σ_j . It is clear that M is homeomorphic to a sphere and has five cone points corresponding to $0, v_0, v_{\pm 1}, v_{\pm 2}, v_{\pm 3}$ with cone angles $\pi - \theta + 2\phi, \pi + \theta, \pi + \theta, \pi + \theta, 2\pi - 2\theta - 2\phi$ respectively. These are the cone angles given in (1).

2.2 Moves on the cone structure

We define automorphisms which we call *moves* on such polygons in the spirit of Thurston [12]. These generalise the moves constructed by Parker in [10] in an obvious way.

We define them as follows. Our cone manifold has five cone points. The two corresponding to 0 and v_0 have cone angles $2\pi - 2\theta - 2\phi$ and $\pi + \phi - \theta$,




Figure 4: *The cut for move R_1 .*

respectively. The other three vertices have the same cone angle, which is $\pi + \theta$. In cutting our cone manifold to get back our octagon, there is no canonical ordering of these three vertices, hence we can change the order of the cut. This results in the moves we will consider namely R_1 and R_2 . We introduce a third move A_1 which generalises Thurston's butterfly moves.

- **The move R_1** The move R_1 fixes the vertex 0, v_0 and $v_{\pm 1}$ and then interchanges $v_{\pm 2}$ and $v_{\pm 3}$. This corresponds to a Dehn Twist along a simple closed curve through $v_{\pm 2}$ and $v_{\pm 3}$ that does not separate the other cone points. This is described on page 242 of Mostow's survey article [9]. When cutting open the cone manifold, one must begin cutting from 0 and then to $v_{\pm 2}$, then to $v_{\pm 3}$, and then to $v_{\pm 1}$ and v_0 ; see Figure 4. When we cut open the double pentagon, we obtain an octagon, shown in Figure 5.

Using cut and paste, one can obtain the new octagon from the old. The cut goes from 0 directly to v_2 . Then the triangle 0, v_2 , v_3 must be glued back on along the edge 0, v_{-3} according to the side identification σ_1 . In the same way, the triangle v_{-1} , v_{-2} , v_{-3} must be glued by σ_3^{-1} to the side v_1 , v_2 ; see Figure 5.




Figure 5: *The octagon obtained after performing move R_1 .*

We now find the new parameters w_1, w_2, w_3 for the new polygon by analysing the vertices. We write the new vertices as v'_j . Then: $v'_0 = v_0$, $v'_1 = v_1$, $v'_3 = v_2$. Thus:

$$\begin{aligned} \frac{-i \sin \theta w_3}{\sin(\theta + \phi)} + \frac{i \sin \theta w_1}{\sin \phi + \sin(\theta - \phi)} &= \frac{-i \sin \theta z_3}{\sin(\theta + \phi)} + \frac{i \sin \theta z_1}{\sin \phi + \sin(\theta - \phi)}, \\ ie^{-i\phi} w_1 - \frac{i \sin \theta w_3}{\sin(\theta + \phi)} &= ie^{-i\phi} z_1 - \frac{i \sin \theta z_3}{\sin(\theta + \phi)}, \\ -ie^{-i\theta - i\phi} w_2 + \frac{i \sin \phi e^{-i\theta - i\phi} w_3}{\sin(\theta + \phi)} &= -ie^{-i\phi} z_2 + \frac{i \sin \phi e^{-i\theta - i\phi} z_3}{\sin \theta + \phi}. \end{aligned}$$

Solving these simultaneous equations give you the following:

$$w_1 = z_1, \quad w_2 = e^{i\theta} z_2, \quad w_3 = z_3.$$

Hence the triangle v_1, v_2, v_3 should be glued by σ_2 to v_{-2}, v_{-3} . We also analyse the vertices to find the new coordinates; $v'_0 = v_0, v'_2 = v_1, v'_3 = v_3$ as before:




Figure 7: *The octagon obtained after performing move R_2 .*

$$\begin{aligned} \frac{-i \sin \theta w_3}{\sin(\theta + \phi)} + \frac{i \sin \theta w_1}{\sin \phi + \sin(\theta - \phi)} &= \frac{-i \sin \theta z_3}{\sin(\theta + \phi)} + \frac{i \sin \theta z_1}{\sin \phi + \sin(\theta - \phi)}, \\ -ie^{-i\phi} w_2 + \frac{i \sin \phi e^{-i\theta - i\phi} w_3}{\sin \theta + \phi} &= ie^{-i\phi} z_1 - \frac{i \sin \theta z_3}{\sin(\theta + \phi)}, \\ -ie^{-i\theta - i\phi} w_2 + \frac{i \sin \phi e^{-i\theta - i\phi} w_3}{\sin(\theta + \phi)} &= -ie^{-i\theta - i\phi} z_2 + \frac{i \sin \phi e^{-i\theta - i\phi} z_3}{\sin(\theta + \phi)} \end{aligned}$$

Solving these simultaneously results in the following matrix R_2 as the solution:

$$\begin{aligned} &(1 - e^{-i\theta}) \sin(\phi) R_2 \\ &= \begin{bmatrix} -\sin(\theta)e^{-i\phi} & -\sin(\phi) - \sin(\theta - \phi) & \sin(\phi) + \sin(\theta - \phi) \\ -\sin(\phi) & -\sin(\phi)e^{-i\theta} & \sin(\phi) \\ -\sin(\theta + \phi) & -\sin(\theta + \phi) & \sin(\phi) + \sin(\theta)e^{i\phi} \end{bmatrix} \quad (5) \end{aligned}$$

Again, R_2 preserves area and so is unitary with respect to H .

- **The move A_1** The third move is a generalisation of the ‘butterfly’ move discussed by Thurston [12] and generalises the move I_1 in Parker




Figure 8: *The octagon obtained after performing move A_1 .*

[10]. In terms of monodromy, it is defined in equations (5.2) and (5.3) and illustrated in Figure 1.5 of Mostow [9]. Thurston's butterfly operation moves one edge of the pentagon across a butterfly-shaped quadrilateral of zero signed area, yielding a new polygon of the same area. In our case, fix $v_{\pm 2}, v_{\pm 3}$ and we rotate the triangle T_1 so that $v'_1 = v_{-1}$. The resulting octagon has a point of self intersection, but by using signed area we still preserve H . The move A_1 preserves the triangles T_2 and T_3 and so it fixes z_2 and z_3 . The triangle T_1 is rotated by 2ϕ and so z_1 is sent to $e^{2i\phi}z_1$. That is, as a matrix, A_1 is given by:

$$A_1 = \begin{bmatrix} e^{2i\phi} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}. \quad (6)$$

Our goal will be to consider the group $\Gamma = \langle R_1, R_2, A_1 \rangle$ generated by the moves R_1, R_2 and A_1 . We view these moves as the matrices given in

(4), (5) and (6). All these moves preserve the (signed) area of Π and so the matrices are all unitary with respect to the Hermitian form given by the matrix H given in (3). We will show that Γ is discrete for the various values of θ and ϕ given in table (2).

3 Construction of Bisectors and Vertices

3.1 Introduction

In this section, we show how the collection of polygons Π (cone metrics on the sphere) can be parametrised by a subset of complex hyperbolic space and we discuss how the moves (automorphisms) R_1 , R_2 and A_1 act as complex hyperbolic isometries. We will actually be looking at the geometry of the action of the isometries. We construct a polyhedron D whose sides are contained in bisectors and whose vertices correspond to certain cone metrics which have degenerated. This degeneration is obtained either from the collision of three cone points or from the collision of two pairs of cone points.

3.2 New Coordinates

Complex hyperbolic space can be defined to be the projectivisation of those points in the space for which the Hermitian form form is positive. We achieve the projectivisation by considering the section for which $z_3 = 1$. Using the definition of H in (3), we see that our model of complex hyperbolic space is defined as follows:

$$\mathbf{H}_{\mathbb{C}}^2 = \left\{ \mathbf{z} : \langle \mathbf{z}, \mathbf{z} \rangle = \mathbf{z}^* H \mathbf{z} > 0 \right\}.$$

Equivalently:

$$\mathbf{H}_{\mathbb{C}}^2 = \left\{ \begin{bmatrix} z_1 \\ z_2 \\ 1 \end{bmatrix} : \frac{-|z_1|^2 \sin \theta \sin \phi}{(\sin \phi + \sin(\theta - \phi))} - |z_2|^2 \sin \theta + \frac{\sin \theta \sin \phi}{\sin(\theta + \phi)} > 0 \right\}. \quad (7)$$

There will be two elements of the group $\Gamma = \langle R_1, R_2, A_1 \rangle$ that are of particular interest to us, namely $P = R_1 R_2$ and $J = P A_1 = R_1 R_2 A_1$. Using the matrices for R_1 , R_2 and A_1 in equations (4), (5) and (6), we find the

following matrices for P and J :

$$(1 - e^{-i\theta}) \sin(\phi) P = \begin{bmatrix} -\sin(\theta)e^{-i\phi} & -\sin(\phi) - \sin(\theta - \phi) & \sin(\phi) + \sin(\theta - \phi) \\ -\sin(\phi)e^{i\theta} & -\sin(\phi) & \sin(\phi)e^{i\theta} \\ -\sin(\theta + \phi) & -\sin(\theta + \phi) & \sin(\phi) + \sin(\theta)e^{i\phi} \end{bmatrix}, \quad (8)$$

$$(1 - e^{-i\theta}) \sin(\phi) J = \begin{bmatrix} -\sin(\theta)e^{i\phi} & -\sin(\phi) - \sin(\theta - \phi) & \sin(\phi) + \sin(\theta - \phi) \\ -\sin(\phi)e^{i(2\phi+\theta)} & -\sin(\phi) & \sin(\phi)e^{i\theta} \\ -\sin(\theta + \phi)e^{2i\phi} & -\sin(\theta + \phi) & \sin(\phi) + \sin(\theta)e^{i\phi} \end{bmatrix}. \quad (9)$$

Note that $\text{tr}(J) = 0$ and so J has order 3. Also, we have put the scalar factor of $(1 - e^{-i\theta}) \sin(\phi)$ (which comes from R_2) on the left hand side of the equation. In our application we are only interested in projective classes of matrices. Therefore this factor may be dropped. We denote projective equality by \sim .

We remark that, using the action of R_1 and R_2 on the cone points, as discussed above, we can summarise the action of P on the cone points as follows:

Lemma 3.1 *The map $P = R_1 R_2$ fixes the cone points 0 and v_0 and maps the other cone points as follows:*

$$P : v_{\pm 1} \mapsto v_{\pm 3}, \quad P : v_{\pm 2} \mapsto v_{\pm 1}, \quad P : v_{\pm 3} \mapsto v_{\pm 2}.$$

We now define our second set of coordinates denoted by \mathbf{w} , which is the preimage under P of the first set of coordinates. Geometrically, the w_j -parameters have the following meaning. Before cutting the sphere to form an octagon, cyclically permute the cone points $v_{\pm j}$ as described in Lemma 3.1. The resulting octagon may be built up from three triangles in just the same way that Π was built up from the triangles T_1 , T_2 and T_3 . The coordinates w_1 , w_2 and w_3 are then the bases of the new triangles. We consider the section with $w_3 = 1$.

In terms of parameters, the new coordinates are given by:

$$\begin{aligned} \mathbf{w} &= \begin{bmatrix} w_1 \\ w_2 \\ 1 \end{bmatrix} = \begin{bmatrix} P^{-1}(\mathbf{z}) \end{bmatrix} \\ &\sim \begin{bmatrix} -\sin(\theta)e^{i\phi} & -(\sin(\phi) + \sin(\theta - \phi))e^{-i\theta} & \sin(\phi) + \sin(\theta - \phi) \\ -\sin(\phi) & -\sin(\phi) & \sin(\phi) \\ -\sin(\theta + \phi) & -\sin(\theta + \phi)e^{-i\theta} & \sin(\phi) + \sin(\theta)e^{-i\phi} \end{bmatrix} \begin{bmatrix} z_1 \\ z_2 \\ 1 \end{bmatrix}. \end{aligned}$$

Hence finding w_1 and w_2 as rational functions of z_1 and z_2 , we obtain

$$\begin{aligned}
w_1 &= \frac{-\sin(\theta)e^{i\phi}z_1 - (\sin(\phi) + \sin(\theta - \phi))e^{-i\theta}z_2 + \sin(\phi) + \sin(\theta - \phi)}{-\sin(\theta + \phi)z_1 - \sin(\theta + \phi)e^{-i\theta}z_2 + \sin(\phi) + \sin(\theta)e^{-i\phi}} \quad (10) \\
w_2 &= \frac{-\sin(\phi)z_1 - \sin(\phi)z_2 + \sin(\phi)}{-\sin(\theta + \phi)z_1 - \sin(\theta + \phi)e^{-i\theta}z_2 + \sin(\phi) + \sin(\theta)e^{-i\phi}}. \quad (11)
\end{aligned}$$

By a similar procedure, we obtain z_1 and z_2 in terms of w_1 and w_2 :

$$\begin{aligned}
\mathbf{z} &= \begin{bmatrix} z_1 \\ z_2 \\ 1 \end{bmatrix} = [P(\mathbf{w})] \\
&\sim \begin{bmatrix} -\sin(\theta)e^{-i\phi} & -\sin(\phi) - \sin(\theta - \phi) & \sin(\phi) + \sin(\theta - \phi) \\ -\sin(\phi)e^{i\theta} & -\sin(\phi) & \sin(\phi)e^{i\theta} \\ -\sin(\theta + \phi) & -\sin(\theta + \phi) & \sin(\phi) + \sin(\theta)e^{i\phi} \end{bmatrix} \begin{bmatrix} w_1 \\ w_2 \\ 1 \end{bmatrix}.
\end{aligned}$$

and hence

$$\begin{aligned}
z_1 &= \frac{-\sin(\theta)e^{-i\phi}w_1 - (\sin(\phi) + \sin(\theta - \phi))w_2 + \sin(\phi) + \sin(\theta - \phi)}{-\sin(\theta + \phi)w_1 - \sin(\theta + \phi)w_2 + \sin(\phi) + \sin(\theta)e^{i\phi}} \quad (12) \\
z_2 &= \frac{-\sin(\phi)e^{i\theta}w_1 - \sin(\phi)w_2 + \sin(\phi)e^{i\theta}}{-\sin(\theta + \phi)w_1 - \sin(\theta + \phi)w_2 + \sin(\phi) + \sin(\theta)e^{i\phi}}. \quad (13)
\end{aligned}$$

Our reason of keeping track of two coordinates is that it gives a simple description of the polyhedron D in terms of the arguments of z_1 , z_2 , w_1 and w_2 .

3.3 Vertices

In this section, we obtain some distinguished points of $\mathbf{H}_{\mathbb{C}}^2$ which will be the vertices of our polyhedron. It will be useful to give these points in the two sets of coordinates \mathbf{w} and \mathbf{z} constructed in the previous section. The distinguished points (cone structures) are obtained by letting some of the cone points approach each other until in the limit they coalesce, and then result in a new point. The complementary angle (curvature) of this new cone point (that is 2π minus the cone angle) is the sum of the complementary angles of the cone points that have coalesced. Considering this from the view point of the octagon Π considered in Section 2, obtaining the new cone points is the same as either expanding or contracting the triangles T_1 and T_2 till some of the vertices become the same point. If such vertices are adjacent




Figure 9: Degenerate octagons corresponding to vertices of D .

to each other then the edge between them has degenerated to a point. We define the following vertices by where various cone points coalesce, and we give the cone angle at the resulting new vertex.

Point	Cone Points	Angle	Cone Points	Angle
p_1	$v_0, v_{\pm 1}$	2ϕ	$v_{\pm 2}, v_{\pm 3}$	2θ
p_2	$v_0, v_{\pm 3}$	2ϕ	$v_{\pm 1}, v_{\pm 2}$	2θ
p_{231}	$v_{\pm 1}, v_{\pm 2}, v_{\pm 3}$	$3\theta - \pi$		
p_{23}	$v_0, v_{\pm 2}, v_{\pm 3}$	$\theta + 2\phi - \pi$		
p_{31}	$v_0, v_{\pm 1}, v_{\pm 2}$	$\theta + 2\phi - \pi$		
p_{12}	$v_0, v_{\pm 1}, v_{\pm 3}$	$\theta + 2\phi - \pi$		

One can notice from the above table that $3\theta \geq \pi$ and $\theta + 2\phi \geq \pi$. This will be the case for all the angles we are interested in; see Table (2). When $3\theta = \pi$ the vertex p_{231} will be on the boundary of complex hyperbolic space. Likewise, when $\theta + 2\phi = \pi$ the vertices p_{12} , p_{23} and p_{31} will be on the boundary of complex hyperbolic space. We now describe the corresponding degenerate octagons in detail; see Figure 9.

p_1 : When v_0 and $v_{\pm 1}$ coalesce, the triangle T_1 shrinks to a point and so $z_1 = 0$. Likewise, when $v_{\pm 2}$ and $v_{\pm 3}$ coalesce then T_2 also shrinks to a point and so $z_2 = 0$. Thus p_1 is given by $z_1 = z_2 = 0$. This is the

origin in the z -coordinates. Putting $z_1 = z_2 = 0$ into (10) and (11) gives:

$$w_1 = \frac{\sin \phi + \sin(\theta - \phi)}{\sin \phi + \sin \theta e^{-i\phi}}, \quad w_2 = \frac{\sin \phi}{\sin \phi + \sin \theta e^{-i\phi}}.$$

p_2 : When v_0 and v_3 coalesce and v_1 and v_2 coalesce, the triangles T_1 and T_2 share an edge. Also T_{-1} and T_{-2} share the vertex $v_{-1} = v_{-2}$. Alternatively, applying the map P permutes the cone points as in Lemma 3.1. This has the effect of making us use the w -coordinates. Therefore, we can repeat our argument for p_1 to see that p_2 corresponds to $w_1 = w_2 = 0$. Putting this into equations (12) and (13) gives:

$$z_1 = \frac{\sin \phi + \sin(\theta - \phi)}{\sin \phi + \sin \theta e^{i\phi}}, \quad z_2 = \frac{\sin \phi e^{i\phi}}{\sin \phi + \sin \theta e^{i\phi}}.$$

Note that (using $z_3 = 1$) when $v_{\pm 1}$ and $v_{\pm 2}$ coalesce we have $z_1 + z_2 = 1$. Likewise, when v_0 and v_3 coalesce we have

$$ie^{-i\phi} = \frac{i \sin \theta}{\sin \phi + \sin(\theta - \phi)} z_1 + ie^{-i\theta - i\phi} z_2. \quad (14)$$

These two equations yield the same solution for z_1 and z_2 .

p_{231} : In this case, the cone points $v_{\pm 1}$, $v_{\pm 2}$ and $v_{\pm 3}$ coalesce. Once again, as $v_{\pm 1}$ and $v_{\pm 2}$ coalesce we have $z_1 + z_2 = z_3 = 1$. Also, since $v_{\pm 2}$ and $v_{\pm 3}$ coalesce, the triangle T_2 shrinks to a point and we have $z_2 = 0$. Hence $z_1 = 1$. Either using Lemma 3.1 or substituting directly, we see that $w_1 = 1$ and $w_2 = 0$.

p_{23} : In this case v_0 , $v_{\pm 2}$ and $v_{\pm 3}$ coalesce. Once again, when $v_{\pm 2}$ and $v_{\pm 3}$ coalesce we obtain $z_2 = 0$ and when v_0 , v_3 coalesce we have equation (14). Putting this together gives

$$z_1 = \frac{\sin \phi + \sin(\theta - \phi)}{\sin \theta e^{i\phi}}, \quad z_2 = 0.$$

In terms of the w -coordinates, when v_0 and v_3 coalesce then $w_1 = 0$ and when v_2 and v_3 coalesce we get $w_1 + e^{-i\theta} w_2 = 1$. Hence $w_1 = 0$ and $w_2 = e^{i\theta}$.

p_{31} : Here v_0 , $v_{\pm 1}$ and $v_{\pm 2}$ coalesce. Arguing as before, we have $z_1 = 0$ and $z_1 + z_2 = 1$. Hence $z_2 = 1$. In terms of w -coordinates, we have $w_2 = 0$. A slightly more involved argument shows that

$$w_1 = \frac{\sin \phi + \sin(\theta - \phi)}{\sin \theta e^{-i\phi}}.$$

p_{12} : Here $v_0, v_{\pm 1}$ and $v_{\pm 3}$ coalesce. The above arguments yield $z_1 = 0$ and $z_1 + z_2 = e^{i\theta}$. Thus $z_2 = e^{i\theta}$. Also, $w_1 = 0$ and $w_1 + w_2 = 1$. Hence $w_2 = 1$.

In coordinates (normalising so $z_3 = w_3 = 1$) we have

Point	z_1	z_2	w_1	w_2
p_1	0	0	$\frac{\sin \phi + \sin(\theta - \phi)}{\sin \phi + \sin \theta e^{-i\phi}}$	$\frac{\sin \phi}{\sin \phi + \sin \theta e^{-i\phi}}$
p_2	$\frac{\sin \phi + \sin(\theta - \phi)}{\sin \phi + \sin \theta e^{i\phi}}$	$\frac{\sin \phi e^{i\theta}}{\sin \phi + \sin \theta e^{i\phi}}$	0	0
p_{231}	1	0	1	0
p_{23}	$\frac{\sin \phi + \sin(\theta - \phi)}{\sin \theta} e^{-i\phi}$	0	0	$e^{i\theta}$
p_{31}	0	1	$\frac{\sin \phi + \sin(\theta - \phi)}{\sin \theta} e^{i\phi}$	0
p_{12}	0	$e^{i\theta}$	0	1

In concluding this section, we show that the collection of vertices described above is symmetrical with respect to an involution. The polyhedron D will also demonstrate this symmetry when we get to Section 4. Let us consider the antiholomorphic isometry ι given by $\iota(\mathbf{z}) = R_1 R_2 R_1(\bar{\mathbf{z}})$, which is the same as $\iota(\mathbf{z}) = P R_1(\bar{\mathbf{z}})$. In coordinates:

$$\iota \begin{bmatrix} z_1 \\ z_2 \\ 1 \end{bmatrix} \sim \begin{bmatrix} \bar{w}_1 \\ \bar{w}_2 e^{i\theta} \\ 1 \end{bmatrix}. \quad (15)$$

Notice that \sim refers to projective equality. The following lemma deduced from the above equation can be verified using the vertices obtained in the above table of vertices.

Lemma 3.2 *The isometry ι has order 2 and acts on the p_j by*

$$\iota(p_1) = p_2, \quad \iota(p_{231}) = p_{231}, \quad \iota(p_{23}) = p_{31}, \quad \iota(p_{12}) = p_{12}.$$

PROOF: This follows by direct calculation. For example to see that ι fixes p_{231} observe

$$\iota \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} = \frac{1}{(1 - e^{-i\theta}) \sin(\phi)} \begin{bmatrix} -\sin(\theta) e^{-i\phi} + \sin(\phi) + \sin(\theta - \phi) \\ -\sin(\phi) e^{i\theta} + \sin(\phi) e^{i\theta} \\ -\sin(\theta + \phi) + \sin(\phi) + \sin(\theta) e^{i\phi} \end{bmatrix} \sim \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}.$$

where we have used

$$\sin(\theta - \phi) - \sin \theta e^{-i\phi} = -\cos \theta \sin \phi + i \sin \theta \sin \phi = -\sin(\theta + \phi) + \sin \theta e^{i\phi}.$$

Similarly to see that $\iota(p_{31}) = p_{23}$ observe

$$\iota \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} = \frac{1}{(1 - e^{-i\theta}) \sin(\phi)} \begin{bmatrix} (1 - e^{i\theta})(\sin \phi + \sin(\theta - \phi)) \\ 0 \\ \sin \phi + \sin \theta e^{i\phi} - \sin(\theta + \phi)e^{i\theta} \end{bmatrix} \sim \begin{bmatrix} \frac{\sin \phi + \sin(\theta - \phi)}{\sin \theta e^{i\phi}} \\ 0 \\ 1 \end{bmatrix}$$

where we have used

$$\sin \phi + \sin \theta e^{i\phi} - \sin(\theta + \phi)e^{i\theta} = \sin \theta e^{i\phi}(1 - e^{i\theta}).$$

The other identities follow similarly by substituting and then simplifying using trigonometric formulae. \square

4 Construction of the Complex Hyperbolic Polyhedron D

4.1 Introduction

In this section we construct a polyhedron D in complex hyperbolic space. In the next section we will use Poincaré's polyhedron theorem to demonstrate that this is a fundamental polyhedron for $\Gamma = \langle R_1, R_2, A_1 \rangle$.

The vertices of D will be the five special cone manifolds $p_1, p_2, p_{231}, p_{23}, p_{31}$ and p_{12} constructed above in Section 3.3. Combinatorially, D is almost as simple as it can be: it will be the union of two four-simplices. The vertices of the first simplex are all the vertices except p_2 and the vertices of the second are all except p_1 .

The co-dimension 1 sides of D will be contained in bisectors. A bisector B is the locus of points equidistant from a given pair of points. They have been studied extensively and we will briefly summarise their properties. For more detail see Mostow [8] or Goldman [6]. If the bisector B is equidistant from points q_1 and q_2 then the complex line $\Sigma = \Sigma(B)$ spanned by q_1 and q_2 is called the *complex spine* of B . The geodesic $\sigma = \sigma(B)$ in Σ equidistant from q_1 and q_2 is called the *spine* of B . Bisectors are not totally geodesic but are foliated by totally geodesic subspaces in two different ways. First, if Π_Σ is orthogonal projection onto Σ then $B = \Pi_\Sigma^{-1}(\sigma)$. For each point s on σ , the preimage $\Pi_\Sigma^{-1}(s)$ is a complex line, called a *slice* of Σ ; see Mostow [8]. The slices foliate B . Secondly, B is the union of all totally real Lagrangian planes containing σ . Such a plane is called a *meridian*; see Goldman [6]. We will use this structure very little. One property we will use is that each

codimension 1 side of D will be a 3-simplex in a bisector with one edge in σ , two faces in meridians and one face in a slice. We will see, in Proposition 4.8, this has the consequence that the 1-skeleton of D is made up of geodesic arcs.

4.2 The polyhedron D

We define the polyhedron D to be those points of $\mathbf{H}_{\mathbb{C}}^2$ for which the arguments of z_1 , z_2 , w_1 and w_2 lie in the following intervals (compare Equation (17) of Parker [10]):

$$D = \left\{ \mathbf{z} = P(\mathbf{w}) : \begin{array}{l} \arg(z_1) \in (-\phi, 0), \quad \arg(z_2) \in (0, \theta), \\ \arg(w_1) \in (0, \phi), \quad \arg(w_2) \in (0, \theta) \end{array} \right\}. \quad (16)$$

We want to characterise D in terms of inequalities involving the Hermitian form. The following lemma generalises Lemmas 4.4 and 4.6 of [10]

Lemma 4.1 *Let $\langle \cdot, \cdot \rangle$ be the Hermitian form given by H . Using the coordinates \mathbf{z} and \mathbf{w} we have:*

- $\text{Im}(z_1 e^{i\phi}) > 0$ if and only if $|\langle \mathbf{z}, p_{231} \rangle| < |\langle \mathbf{z}, J^{-1}(p_{231}) \rangle|$;
- $\text{Im}(z_1) < 0$ if and only if $|\langle \mathbf{z}, p_{23} \rangle| < |\langle \mathbf{z}, P^{-1}(p_{31}) \rangle|$;
- $\text{Im}(z_2) > 0$ if and only if $|\langle \mathbf{z}, p_{12} \rangle| < |\langle \mathbf{z}, R_1^{-1}(p_{31}) \rangle|$;
- $\text{Im}(z_2 e^{-i\theta}) < 0$ if and only if $|\langle \mathbf{z}, p_{31} \rangle| < |\langle \mathbf{z}, R_1(p_{12}) \rangle|$;
- $\text{Im}(w_1) > 0$ if and only if $|\langle \mathbf{w}, p_{31} \rangle| < |\langle \mathbf{w}, P(p_{23}) \rangle|$;
- $\text{Im}(w_1 e^{-i\phi}) < 0$ if and only if $|\langle \mathbf{w}, p_{231} \rangle| < |\langle \mathbf{w}, J(p_{231}) \rangle|$;
- $\text{Im}(w_2) > 0$ if and only if $|\langle \mathbf{w}, p_{23} \rangle| < |\langle \mathbf{w}, R_2^{-1}(p_{12}) \rangle|$;
- $\text{Im}(w_2 e^{-i\theta}) < 0$ if and only if $|\langle \mathbf{w}, p_{12} \rangle| < |\langle \mathbf{w}, R_2(p_{23}) \rangle|$.

PROOF: We illustrate one case of this theorem. In \mathbf{z} -coordinates

$$p_{231} = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \quad J^{-1}(p_{231}) = \begin{bmatrix} e^{-2i\phi} \\ 0 \\ 1 \end{bmatrix}.$$

Therefore

$$\begin{aligned} \langle \mathbf{z}, p_{231} \rangle &= \frac{-\sin \phi}{\sin \phi + \sin(\theta - \phi)} z_1 + \frac{\sin \phi}{\sin(\theta + \phi)}, \\ \langle \mathbf{z}, J^{-1}(p_{231}) \rangle &= \frac{-\sin \phi}{\sin \phi + \sin(\theta - \phi)} e^{2i\phi} z_1 + \frac{\sin \phi}{\sin(\theta + \phi)}. \end{aligned}$$

Hence $|\langle \mathbf{z}, p_{231} \rangle| < |\langle \mathbf{z}, J^{-1}(p_{231}) \rangle|$ if and only if $-2\operatorname{Re}(z_1) < -2\operatorname{Re}(z_1 e^{2i\phi})$. This is true if and only if $\operatorname{Im}(z_1 e^{i\phi}) > 0$. This is the first part. The other parts are all similar. \square

Corollary 4.2 *The polyhedron D is defined by points $p \in \mathbf{H}_{\mathbb{C}}^2$ (which may be written in \mathbf{z} or \mathbf{w} coordinates) satisfying the inequalities:*

$$\begin{array}{ll} |\langle p, p_{231} \rangle| < |\langle p, J^{-1}(p_{231}) \rangle|, & |\langle p, p_{23} \rangle| < |\langle p, P^{-1}(p_{31}) \rangle|, \\ |\langle p, p_{12} \rangle| < |\langle p, R_1^{-1}(p_{31}) \rangle|, & |\langle p, p_{31} \rangle| < |\langle p, R_1(p_{12}) \rangle|, \\ |\langle p, p_{31} \rangle| < |\langle p, P(p_{23}) \rangle|, & |\langle p, p_{231} \rangle| < |\langle p, J(p_{231}) \rangle|, \\ |\langle p, p_{23} \rangle| < |\langle p, R_2^{-1}(p_{12}) \rangle|, & |\langle p, p_{12} \rangle| < |\langle p, R_2(p_{23}) \rangle|. \end{array}$$

We refer to the codimension 1 facets of D as *sides*. Each side corresponds to one of the eight inequalities in Lemma 4.1 being replaced with equality. Therefore each of the eight sides of D is contained in a bisector. Moreover, for each of these bisectors B , we have:

- (i) either the point p_1 or p_2 lies on B ;
- (ii) three of the four points p_{231} , p_{23} , p_{31} and p_{12} lie on B ;
- (iii) the fourth of these points lies on the complex spine Σ of B but not on B ;
- (iv) the spine σ of B passes through one of p_1 and p_2 and one of p_{231} , p_{23} , p_{31} or p_{12} .

We name the bisectors $B(X)$ where X is one of J, P, R_1, R_2 or their inverses so that the isometry X will send $B(X)$ to $B(X^{-1})$. For example, $B(R_1)$ is given by the equality

$$|\langle p, p_{12} \rangle| = |\langle p, R_1^{-1}(p_{31}) \rangle|.$$

Applying R_1 sends p_{12} to $R_1(p_{12})$ and sends $R_1^{-1}(p_{31})$ to p_{31} . Therefore R_1 sends $B(R_1)$ to the bisector defined by

$$|\langle p, R_1(p_{12}) \rangle| = |\langle p, p_{31} \rangle|.$$

This is $B(R_1^{-1})$.

The summary is:

Bisector	Definition	Equidistant from	Points on spine	Other points
$B(J)$	$\text{Im}(z_1 e^{i\phi}) = 0$	$p_{231}, J^{-1}(p_{231})$	p_1, p_{23}	p_{31}, p_{12}
$B(J^{-1})$	$\text{Im}(w_1 e^{-i\phi}) = 0$	$p_{231}, J(p_{231})$	p_2, p_{31}	p_{12}, p_{23}
$B(P)$	$\text{Im}(z_1) = 0$	$p_{23}, P^{-1}(p_{31})$	p_1, p_{231}	p_{31}, p_{12}
$B(P^{-1})$	$\text{Im}(w_1) = 0$	$p_{31}, P(p_{23})$	p_2, p_{231}	p_{12}, p_{23}
$B(R_1)$	$\text{Im}(z_2) = 0$	$p_{12}, R_1^{-1}(p_{31})$	p_1, p_{31}	p_{23}, p_{231}
$B(R_1^{-1})$	$\text{Im}(z_2 e^{-i\theta}) = 0$	$p_{31}, R_1(p_{12})$	p_1, p_{12}	p_{23}, p_{231}
$B(R_2)$	$\text{Im}(w_2) = 0$	$p_{23}, R_2^{-1}(p_{12})$	p_2, p_{12}	p_{31}, p_{231}
$B(R_2^{-1})$	$\text{Im}(w_2 e^{-i\theta}) = 0$	$p_{12}, R_2(p_{23})$	p_2, p_{23}	p_{31}, p_{231}

We go through these properties in two cases. The others are similar; see Boadi [1] for details.

- (1) Example, we consider $B(J)$. In \mathbf{z} -coordinates this is given by

$$B(J) = \left\{ (x_1 e^{-i\phi}, z_2) \in \mathbf{H}_{\mathbb{C}}^2 : x_1 \in \mathbb{R}, z_2 \in \mathbb{C} \right\}.$$

The spine $\sigma(J)$ of $B(J)$ has \mathbf{z} -coordinates

$$\sigma(J) = \left\{ (x_1 e^{-i\phi}, 0) \in \mathbf{H}_{\mathbb{C}}^2 : x_1 \in \mathbb{R} \right\}.$$

The complex spine $\Sigma(J)$ of $B(J)$ has \mathbf{z} -coordinates

$$\Sigma(J) = \left\{ (z_1, 0) \in \mathbf{H}_{\mathbb{C}}^2 : z_1 \in \mathbb{C} \right\}.$$

- p_1 is given by $(z_1, z_2) = (0, 0)$. This clearly lies in $B(J)$ and $\sigma(J)$.
- p_{231} is given by $(z_1, z_2) = (1, 0)$ and $J^{-1}(p_{231})$ is given by $(z_1, z_2) = (e^{-2i\phi}, 0)$. These points clearly do not lie on $B(J)$ but do lie on $\Sigma(J)$, and $B(J)$ is equidistant from them.
- p_{23} is given by

$$(z_1, z_2) = \left(\frac{\sin \phi + \sin(\theta - \phi)}{\sin \theta} e^{-i\phi}, 0 \right).$$

Since $(\sin \phi + \sin(\theta - \phi)) / \sin \theta$ is real, this lies on $B(J)$ and $\sigma(J)$.

- p_{31} is given by $(z_1, z_2) = (0, 1)$. This clearly lies on $B(J)$ but does not lie on $\sigma(J)$.
- p_{12} is given by $(z_1, z_2) = (0, e^{i\theta})$. This clearly lies on $B(J)$ but does not lie on $\sigma(J)$.

(2) Next we consider $B(R_2)$. In \mathbf{w} -coordinates this is given by

$$B(R_2) = \left\{ (w_1, w_2) \in \mathbf{H}_{\mathbb{C}}^2 : w_1 \in \mathbb{C}, w_2 \in \mathbb{R} \right\}.$$

The spine $\sigma(R_2)$ of $B(R_2)$ has \mathbf{w} coordinates

$$\sigma(R_2) = \left\{ (0, w_2) \in \mathbf{H}_{\mathbb{C}}^2 : w_2 \in \mathbb{R} \right\}.$$

The complex spine $\Sigma(R_2)$ of $B(R_2)$ has \mathbf{w} coordinates

$$\Sigma(R_2) = \left\{ (0, w_2) \in \mathbf{H}_{\mathbb{C}}^2 : w_2 \in \mathbb{C} \right\}.$$

- p_2 is given by $(w_1, w_2) = (0, 0)$. This clearly lies in $B(R_2)$ and $\sigma(R_2)$.
- p_{231} is given by $(w_1, w_2) = (1, 0)$. This clearly lies in $B(R_2)$ but not in $\sigma(R_2)$.
- p_{31} is given by

$$(w_1, w_2) = \left(\frac{\sin \phi + \sin(\theta - \phi)}{\sin \theta} e^{i\phi}, 0 \right).$$

This clearly lies on $B(R_2)$ but not on $\sigma(R_2)$.

- p_{12} is given by $(w_1, w_2) = (0, 1)$. This lies on both $B(R_2)$ and $\sigma(R_2)$.
- p_{23} is given by $(w_1, w_2) = (0, e^{i\theta})$ and $R_2^{-1}(p_{12})$ is given by $(w_1, w_2) = (0, e^{-i\theta})$. These points clearly do not lie on $B(R_2)$. However, they do lie on $\Sigma(R_2)$, and $B(R_2)$ is equidistant from them.

4.3 Some useful inequalities

When we give details of the faces of D we will need to use the following lemmas which give inequalities satisfied by all points of $\mathbf{H}_{\mathbb{C}}^2$.

Lemma 4.3 *If $\mathbf{z} \in \mathbf{H}_{\mathbb{C}}^2$ then*

$$|z_1| < \frac{\sin \phi + \sin(\theta - \phi)}{\sin(\theta + \phi)}, \quad |w_1| < \frac{\sin \phi + \sin(\theta - \phi)}{\sin(\theta + \phi)}.$$

PROOF: We prove this by contradiction. If $|z_1| \geq (\sin \phi + \sin(\theta - \phi)) / \sin(\theta + \phi)$ then

$$\begin{aligned} \mathbf{z}^* H \mathbf{z} &= \frac{-\sin \phi}{\sin \phi + \sin(\theta - \phi)} |z_1|^2 - |z_2|^2 + \frac{\sin \phi}{\sin(\theta + \phi)} \\ &\leq \frac{-\sin \phi (\sin \phi + \sin(\theta - \phi))}{\sin^2(\theta - \phi)} - |z_2|^2 + \frac{\sin \phi}{\sin(\theta + \phi)} \\ &\leq \frac{\sin^2 \phi (2 \cos \theta - 1)}{\sin^2(\theta + \phi)} - |z_2|^2 \leq 0, \end{aligned}$$

where we have used $\cos \theta \leq 1/2$ on the last line. Thus \mathbf{z} is not in $\mathbf{H}_{\mathbb{C}}^2$. A similar argument holds for \mathbf{w} . \square

Lemma 4.4 *If $\mathbf{z} \in \mathbf{H}_{\mathbb{C}}^2$ then*

$$|z_1|, |w_1| < \frac{\sin \theta}{\sin(\theta + \phi)}, \quad |z_2|, |w_2| < \frac{\sin \phi}{\sin(\theta + \phi)},$$

PROOF: In order to prove the lemma, first observe from the combination of ϕ and θ in Table (2) that we have $\theta + 2\phi \geq \pi$ and hence

$$\pi - (\theta + \phi) \leq \phi \leq \theta + \phi.$$

Therefore $\sin(\phi) \geq \sin(\theta + \phi)$.

Now we prove the inequality for z_1 . If $|z_1| \geq \sin(\theta) / \sin(\theta + \phi)$ then from the area

$$\begin{aligned} \mathbf{z}^* H \mathbf{z} &= \frac{-\sin \phi}{\sin \phi + \sin(\theta - \phi)} |z_1|^2 - |z_2|^2 + \frac{\sin \phi}{\sin(\theta + \phi)} \\ &\leq \frac{\sin \phi}{\sin(\theta + \phi)} \frac{\sin(\theta + \phi) (\sin \phi + \sin(\theta - \phi)) - \sin^2 \theta}{\sin(\theta + \phi) (\sin \phi + \sin(\theta - \phi))} - |z_2|^2 \\ &= \frac{\sin \phi}{\sin(\theta + \phi)} \frac{\sin(\theta + \phi) \sin \phi + \sin^2 \theta \cos^2 \phi - \cos^2 \theta \sin^2 \phi - \sin^2 \theta}{\sin(\theta + \phi) (\sin \phi + \sin(\theta - \phi))} - |z_2|^2 \\ &= \frac{\sin \phi}{\sin(\theta + \phi)} \frac{\sin(\theta + \phi) \sin \phi - \sin^2 \phi}{\sin(\theta + \phi) (\sin \phi + \sin(\theta - \phi))} - |z_2|^2 \\ &= -\frac{\sin^2 \phi}{\sin^2(\theta + \phi)} \frac{\sin \phi - \sin(\theta + \phi)}{\sin \phi + \sin(\theta - \phi)} - |z_2|^2 \leq 0. \end{aligned}$$

This a contradiction. Hence $|z_1| < \sin(\theta) / \sin(\theta + \phi)$. Proving the inequality for w_1 is identical.

Similarly, to prove the inequality for z_2 assume that $|z_2| \geq \sin \phi / \sin(\theta + \phi)$. Then

$$\begin{aligned}
\mathbf{z}^* H \mathbf{z} &= \frac{-\sin \phi}{\sin \phi + \sin(\theta - \phi)} |z_1|^2 - |z_2|^2 + \frac{\sin \phi}{\sin(\theta + \phi)} \\
&\leq \frac{-\sin \phi}{\sin \phi + \sin(\theta - \phi)} |z_1|^2 - \frac{\sin^2 \phi}{\sin^2(\theta + \phi)} + \frac{\sin \phi}{\sin(\theta + \phi)} \\
&\leq \frac{-\sin \phi}{\sin \phi + \sin(\theta - \phi)} |z_1|^2 - \frac{\sin \phi}{\sin(\theta + \phi)} \left(\frac{\sin \phi}{\sin(\theta + \phi)} - 1 \right) \\
&\leq 0,
\end{aligned}$$

as $\sin \phi / \sin(\theta + \phi) \geq 1$. The inequality for w_2 is similar. \square

4.4 Faces of the polyhedron

We refer to codimension 2 facets of the polyhedron D as *faces*. Each face is contained in the intersection of two of the bisectors defining the sides of D . The other bisectors determine further inequalities defining the edges bounding each face so that the face becomes a triangle in this bisector intersection. The faces come in three types: (a) faces contained in a common slice of two of the bisectors; (b) faces contained in a common meridian of two bisectors or (c) faces that are not contained in either a complex line or a Lagrangian plane. In fact the intersections of type (c) are contained in a *Giraud disc*, which is a particularly nice type of bisector intersection; see Theorem 8.3.3 of Goldman [6]. We give representative examples of each type. In Proposition 4.5 we give details of a face in a common slice, in Proposition 4.6 consider a face in a common meridian and in Proposition 4.7 we give a face in a Giraud disc. For the complete list, see Boadi's thesis [1].

First we give a face contained in a complex line that is a common slice of the two bisectors.

Proposition 4.5 *A point in the face of D contained in $B(J) \cap B(P)$ has coordinates $\mathbf{z} = (0, x + ye^{i\theta})$ and $\mathbf{w} = (w_1, w_2)$ with*

$$\begin{aligned}
w_1 &= \frac{(\sin(\phi) + \sin(\theta - \phi))(1 - e^{-i\theta}x - y)}{\sin \phi + \sin \theta e^{-i\phi} - \sin(\theta + \phi)(e^{-i\theta}x + y)}, \\
w_2 &= \frac{\sin \phi(1 - x - e^{i\theta}y)}{\sin \phi + \sin \theta e^{-i\phi} - \sin(\theta + \phi)(e^{-i\theta}x + y)},
\end{aligned}$$

where x, y are non-negative real numbers satisfying

$$0 \leq \sin \phi(1 - x - y) - \sin(\theta + \phi)(x + y - x^2 - y^2 - 2xy \cos \theta).$$

PROOF: Since the point is on $B(J)$ we have $\text{Im}(z_1 e^{i\phi}) = 0$ and as it is on $B(P)$ we have $\text{Im}(z_1) = 0$. Hence the intersection of these bisectors is the complex line $z_1 = 0$, which is a common slice. The conditions $\text{Im}(z_2) \geq 0$ and $\text{Im}(z_2 e^{-i\theta}) \leq 0$ mean that if we write $z_2 = x + ye^{i\theta}$ then $x \geq 0$ and $y \geq 0$. The expressions for w_1 and w_2 follow by substituting these values of z_1 and z_2 into (10) and (11).

It is not hard to check that $\text{Im}(w_1 e^{-i\phi}) \leq 0$ if and only if $\text{Im}(w_2) \geq 0$ if and only if

$$0 \leq \sin \phi(1 - x - y) - \sin(\theta + \phi)(x + y - x^2 - y^2 - 2xy \cos \theta).$$

Note that the curve given by equality in this expression intersects the line $y = 0$ (respectively $x = 0$) at $x = 1$ and $x = \sin \phi / \sin(\theta + \phi)$ (respectively $y = 1$ and $y = \sin \phi / \sin(\theta + \phi)$). The latter points are outside complex hyperbolic space, by Lemma 4.4. Therefore this curve and the lines $x = 0$, $y = 0$ bound a triangle with vertices $(x, y) = (0, 0)$, $(1, 0)$ and $(0, 1)$.

Finally, we must check that $\text{Im}(w_1) \geq 0$ and $\text{Im}(w_2 e^{-i\theta}) \leq 0$. It is easy to check that $\text{Im}(w_1) \geq 0$ if and only if

$$0 \leq 1 - y + x - 2 \cos \theta x.$$

If $y = 1 + x - 2 \cos \theta$ then

$$\sin \phi(1 - x - y) - \sin(\theta + \phi)(x + y - x^2 - y^2 - 2xy \cos \theta) = -2(1 - \cos \theta)x(\sin \phi - \sin(\theta + \phi)x).$$

Using Lemma 4.4 we see that $\sin \phi > \sin(\theta + \phi)x$ and so this expression is negative. Hence $\text{Im}(w_1 e^{-i\phi}) \leq 0$ implies $\text{Im}(w_1) \geq 0$.

Also, $\text{Im}(w_2 e^{-i\theta}) \leq 0$ if and only if

$$0 \leq (\sin \phi + \sin(\theta - \phi))(1 - x + y) - 2 \cos \theta \sin \phi y.$$

A similar argument shows that $\text{Im}(w_2) \geq 0$ implies $\text{Im}(w_2 e^{-i\theta}) \leq 0$. We leave this to the reader. \square

Now we give a face contained in a Lagrangian plane that is a common meridian of the two bisectors.

Proposition 4.6 *A point in the face of D contained in $B(J) \cap B(R_1)$ has coordinates $\mathbf{z} = (xe^{-i\phi}, y)$ and $\mathbf{w} = (w_1, w_2)$ with*

$$w_1 = \frac{-\sin(\theta)x - (\sin(\phi) + \sin(\theta - \phi))e^{-i\theta}y + \sin(\phi) + \sin(\theta - \phi)}{-\sin(\theta + \phi)e^{-i\phi}x - \sin(\theta + \phi)e^{-i\theta}y + \sin(\phi) + \sin(\theta)e^{-i\phi}},$$

$$w_2 = \frac{-\sin(\phi)e^{-i\phi}x - \sin(\phi)y + \sin(\phi)}{-\sin(\theta + \phi)e^{-i\phi}x - \sin(\theta + \phi)e^{-i\theta}y + \sin(\phi) + \sin(\theta)e^{-i\phi}},$$

where x and y are non-negative real numbers satisfying

$$0 \leq (\sin \phi + \sin(\theta - \phi))(1 - y) - \sin \theta x.$$

PROOF: Since the point is on $B(J)$ we have $\text{Im}(z_1 e^{i\phi}) = 0$ and as it is on $B(R_1)$ we have $\text{Im}(z_2) = 0$. Hence the intersection of these bisectors is on the Lagrangian plane where $(z_1, z_2) = (xe^{-i\phi}, y)$ for real x and y . The conditions $\text{Im}(z_1) \leq 0$ and $\text{Im}(z_2 e^{-i\theta}) \leq 0$ imply $x \geq 0$ and $y \geq 0$. The expressions for w_1 and w_2 follow from (10) and (11).

It is not hard to check that $\text{Im}(w_1 e^{-i\phi}) \leq 0$ if and only if

$$0 \leq (\sin \phi - \sin(\theta + \phi)y)((\sin \phi + \sin(\theta - \phi))(1 - y) - \sin \theta x).$$

Similarly $\text{Im}(w_2 e^{-i\theta}) \leq 0$ if and only if

$$0 \leq (\sin \theta - \sin(\theta + \phi)x)((\sin \phi + \sin(\theta - \phi))(1 - y) - \sin \theta x).$$

Using Lemma 4.4 we see that $\sin \theta - \sin(\theta + \phi)x > 0$ and $\sin \phi - \sin(\theta + \phi)y > 0$. Finally, we must check $\text{Im}(w_1) \geq 0$ and $\text{Im}(w_2) \geq 0$. We leave this to the reader. \square

Finally we give a face contained in a Giraud disc.

Proposition 4.7 *A point on the face of D contained in $\mathbf{z} \in B(J) \cap B(J^{-1})$ has coordinates $\mathbf{z} = (xe^{-i\phi}, z_2)$ and $\mathbf{w} = (ue^{i\phi}, w_2)$ where*

$$z_2 = e^{i\theta} \frac{xu \sin(\theta + \phi) - u(e^{i\phi} \sin \phi + \sin \theta) - x \sin \theta + \sin \phi + \sin(\theta - \phi)}{\sin \phi + \sin(\theta - \phi) - ue^{i\phi} \sin(\theta + \phi)},$$

$$w_2 = \frac{xu \sin(\theta + \phi) - x(e^{-i\phi} \sin \phi + \sin \theta) - u \sin \theta + \sin \phi + \sin(\theta - \phi)}{\sin \phi + \sin(\theta - \phi) - x \sin(\theta + \phi)e^{-i\phi}},$$

where x and u are non-negative real numbers satisfying

$$0 \leq xu \sin(\theta - \phi) \sin(\theta + \phi) - (x + u) \sin \theta (\sin \phi + \sin(\theta - \phi)) + (\sin \phi + \sin(\theta - \phi))^2.$$

PROOF: Arguing as before, since the point is on $B(J)$ we have $z_1 = xe^{-i\phi}$ and as it is on $B(J)$ we have $w_1 = ue^{i\phi}$ for real x and u . Since $\text{Im}(z_1) \leq 0$ and $\text{Im}(w_1) \geq 0$ we have $x \geq 0$ and $u \geq 0$. Substituting these in (12) and (10) gives:

$$\begin{aligned} xe^{-i\phi} &= \frac{-u \sin \theta - (\sin \phi + \sin(\theta - \phi))w_2 + \sin \phi + \sin(\theta - \phi)}{-u \sin(\theta + \phi)e^{i\phi} - \sin(\theta + \phi)w_2 + \sin \phi + \sin \theta e^{i\phi}}, \\ ue^{i\phi} &= \frac{-x \sin \theta - e^{-i\theta}(\sin \phi + \sin(\theta - \phi))z_2 + \sin \phi + \sin(\theta - \phi)}{-x \sin(\theta + \phi)e^{-i\phi} - \sin(\theta + \phi)e^{-i\theta}z_2 + \sin \phi + e^{-i\phi} \sin \theta}. \end{aligned}$$

Solving for w_2 and z_2 gives the expressions in the statement of the proposition. The condition $\text{Im}(z_2) \geq 0$ is equivalent to

$$0 \leq (\sin \theta - u \sin(\theta + \phi))p(u, x)$$

where

$$\begin{aligned} p(u, x) &= xu \sin(\theta - \phi) \sin(\theta + \phi) - (x + u) \sin \theta (\sin \phi + \sin(\theta - \phi)) \\ &\quad + (\sin \phi + \sin(\theta - \phi))^2. \end{aligned}$$

The condition that $\text{Im}(w_2 e^{-i\theta}) \leq 0$ is equivalent to

$$0 \leq (\sin \theta - x \sin(\theta + \phi))p(u, x).$$

From Lemma 4.4 we see that $\sin \theta - u \sin(\theta + \phi) > 0$ and $\sin \theta - x \sin(\theta + \phi) > 0$, thus we must have $p(u, x) \geq 0$ as claimed. Finally we must check $\text{Im}(z_2 e^{-i\theta}) \leq 0$ and $\text{Im}(w_2) \geq 0$. We leave this to the reader. \square

We conclude this section by listing all the faces. We use the notation that $F(X, Y)$ is the face of D contained in the intersections of the bisectors $B(X)$ and $B(Y)$. In the following table, the letters S, M, G in the last

column indicate whether the face is in a slice, a meridian or a Giraud disc.

Face	Vertices	Coordinates	
$F(J, J^{-1})$	p_{12}, p_{23}, p_{31}	$\text{Im}(z_1 e^{i\phi}) = \text{Im}(w_1 e^{-i\phi}) = 0$	G
$F(J, P)$	p_1, p_{31}, p_{12}	$\text{Im}(z_1 e^{i\phi}) = \text{Im}(z_1) = 0$	S
$F(J, R_1)$	p_1, p_{31}, p_{23}	$\text{Im}(z_1 e^{i\phi}) = \text{Im}(z_2) = 0$	M
$F(J, R_1^{-1})$	p_1, p_{12}, p_{23}	$\text{Im}(z_1 e^{i\phi}) = \text{Im}(z_2 e^{-i\theta}) = 0$	M
$F(J^{-1}, P^{-1})$	p_2, p_{12}, p_{23}	$\text{Im}(w_1 e^{-i\phi}) = \text{Im}(w_1) = 0$	S
$F(J^{-1}, R_2)$	p_2, p_{31}, p_{12}	$\text{Im}(w_1 e^{-i\phi}) = \text{Im}(w_2) = 0$	M
$F(J^{-1}, R_2^{-1})$	p_2, p_{31}, p_{23}	$\text{Im}(w_1 e^{-i\phi}) = \text{Im}(w_2 e^{-i\theta}) = 0$	M
$F(P, R_1)$	p_1, p_{31}, p_{231}	$\text{Im}(z_1) = \text{Im}(z_2) = 0$	M
$F(P, R_1^{-1})$	p_1, p_{12}, p_{231}	$\text{Im}(z_1) = \text{Im}(z_2 e^{-i\theta}) = 0$	M
$F(P, R_2)$	p_{12}, p_{31}, p_{231}	$\text{Im}(z_1) = \text{Im}(w_2) = 0$	G
$F(P^{-1}, R_1^{-1})$	p_{12}, p_{23}, p_{231}	$\text{Im}(w_1) = \text{Im}(z_2 e^{-i\theta}) = 0$	G
$F(P^{-1}, R_2)$	p_2, p_{12}, p_{231}	$\text{Im}(w_1) = \text{Im}(w_2) = 0$	M
$F(P^{-1}, R_2^{-1})$	p_2, p_{23}, p_{231}	$\text{Im}(w_1) = \text{Im}(w_2 e^{-i\theta}) = 0$	M
$F(R_1, R_1^{-1})$	p_1, p_{23}, p_{231}	$\text{Im}(z_2) = \text{Im}(z_2 e^{-i\theta}) = 0$	S
$F(R_1, R_2^{-1})$	p_{23}, p_{31}, p_{231}	$\text{Im}(z_2) = \text{Im}(w_2 e^{-i\theta}) = 0$	G
$F(R_2, R_2^{-1})$	p_2, p_{31}, p_{231}	$\text{Im}(w_2) = \text{Im}(w_2 e^{-i\theta}) = 0$	S

4.5 Other facets of D

We have discussed the vertices and faces of D , that is the facets of dimension 0 and 2. In this section we discuss the rest of the facets of D . Figure 10 shows that sides of D . We begin with the edges, that is facets of dimension 1. We refer to the edge joining vertices p_a and p_b as $\gamma(a, b)$.

Proposition 4.8 *Each edge $\gamma(a, b)$ of D is a geodesic segment joining a pair of the vertices p_a and p_b .*

PROOF: Each edge will be contained in either three or four of the faces listed in the previous section. We refer to faces contained in a common slice, a common meridian or a Giraud disc as S-faces, M-faces or G-faces

respectively. We now list the edges together with the faces containing them.

Edge	S-face	M-face	M-face	G-face	G-face
$\gamma(1, 12)$	$F(J, P)$	$F(J, R_1^{-1})$	$F(P, R_1^{-1})$		
$\gamma(1, 23)$	$F(R_1, R_1^{-1})$	$F(J, R_1)$	$F(J, R_1^{-1})$		
$\gamma(1, 31)$	$F(J, P)$	$F(J, R_1)$	$F(P, R_1)$		
$\gamma(1, 231)$	$F(R_1, R_1^{-1})$	$F(P, R_1)$	$F(P, R_1^{-1})$		
$\gamma(2, 12)$	$F(J^{-1}, P^{-1})$	$F(J^{-1}, R_2)$	$F(P^{-1}, R_2)$		
$\gamma(2, 23)$	$F(J^{-1}, P^{-1})$	$F(J^{-1}, R_2^{-1})$	$F(P^{-1}, R_2^{-1})$		
$\gamma(2, 31)$	$F(R_2, R_2^{-1})$	$F(J^{-1}, R_2)$	$F(J^{-1}, R_2^{-1})$		
$\gamma(2, 231)$	$F(R_2, R_2^{-1})$	$F(P^{-1}, R_2)$	$F(P^{-1}, R_2^{-1})$		
$\gamma(231, 12)$		$F(P, R_1^{-1})$	$F(P^{-1}, R_2)$	$F(P, R_2)$	$F(P^{-1}R_1^{-1})$
$\gamma(231, 23)$	$F(R_1, R_1^{-1})$	$F(P^{-1}, R_2^{-1})$		$F(P^{-1}R_1^{-1})$	$F(R_1, R_2^{-1})$
$\gamma(231, 31)$	$F(R_2, R_2^{-1})$	$F(P, R_1)$		$F(P, R_2)$	$F(R_1, R_2^{-1})$
$\gamma(12, 23)$	$F(J^{-1}, P^{-1})$	$F(J, R_1^{-1})$		$F(J, J^{-1})$	$F(P^{-1}, R_1^{-1})$
$\gamma(23, 31)$		$F(J, R_1)$	$F(J^{-1}, R_2^{-1})$	$F(J, J^{-1})$	$F(R_1, R_2^{-1})$
$\gamma(31, 12)$	$F(J, P)$	$F(J^{-1}, R_2)$		$F(J, J^{-1})$	$F(P, R_2)$

For each edge, at least two of these faces are totally geodesic, that is they are contained in a complex line or Lagrangian plane. This means that the edge is a geodesic segment. In particular:

- (1) Each edge ending at either p_1 or p_2 is on one S-face and two M-faces. Moreover, the two M-faces are contained in meridians of the same bisector. Therefore this edge is a geodesic segment contained in the spine of this bisector.
- (2) Each edge of the other edges is in two G-faces and either in two M-faces or an S-face and an M-face.

This completes the proof. \square

Let $S(X)$ denote the side (that is codimension 1 facet) of D contained in the bisector $B(X)$. A consequence of the analysis we have done is:

Proposition 4.9 *Each side $S(X)$ of D is a 3-simplex (solid tetrahedron) contained in the bisector $B(X)$. It has one face contained in a slice of $B(X)$ and two faces contained in meridians of $B(X)$, intersecting in an arc of the spine $\sigma(X)$. The fourth face is contained in a Giraud disc.*

We also have a proposition describing the faces more precisely:

Proposition 4.10 *Each face $F(X, Y)$ of D is a 2-simplex (solid triangle) homeomorphic to an open disc in \mathbb{R}^2 and the boundary of $F(X, Y)$ is made up of three geodesic segments joining the vertices.*

5 Proof that D is a fundamental Polyhedron for the Group

5.1 Introduction

In this section we show that the group Γ generated by R_1 , R_2 and A_1 is discrete group with fundamental polyhedron D . To do so, we use Poincaré's polyhedron theorem. For accounts of Poincaré's polyhedron theorem see Parker [10] or Mostow [8]. An account of Poincaré's theorem in the constant curvature setting is given by Epstein and Petronio [4]. Our proof is based on Parker's proof of a similar result for Livné's lattices. Full details are given in Boadi's thesis [1]. We will indicate how the proof goes but we do not give full details.

Our main theorem is

Theorem 5.1 *Let R_1 , R_2 , P and J be given by (4), (5), (8) and (9) respectively. The subgroup Γ of $\text{PU}(H)$ generated by these maps is discrete and the polyhedron D constructed in Section 4 is a fundamental polyhedron for Γ . Moreover, Γ has the following presentation:*

$$\Gamma = \left\langle J, P, R_1, R_2 : \begin{array}{l} J^3 = R_1^p = R_2^p = (P^{-1}J)^k = I, \\ R_2 = PR_1P^{-1} = JR_1J^{-1}, P = R_1R_2 \end{array} \right\rangle \quad (17)$$

where the values of p and k are given by:

p	3	3	3	4	4	5	5	6	6
k	4	5	6	3	4	2	3	2	3

We remark that, since $P = R_1R_2$ and $J = R_1R_2A_1$ the group Γ is the same as $\langle R_1, R_2, A_1 \rangle$. Substituting this into the presentation (17) gives:

$$\Gamma = \left\langle R_1, R_2, A_1 : \begin{array}{l} R_1^p = R_2^p = A_1^k = (R_1R_2A_1)^3 = I, \\ R_1R_2R_1 = R_2R_1R_2, R_1A_1 = A_1R_1 \end{array} \right\rangle.$$

5.2 The side pairing maps

There are eight sides of D , namely $S(J^{\pm 1})$, $S(P^{\pm 1})$, $S(R_1^{\pm 1})$ and $S(R_2^{\pm 1})$ as constructed in Section 4. These sides are paired by the maps J , P , R_1 and

R_2 as described in the following proposition. Recall that D , as defined in (16), is an open polyhedron. We use the convention that that sides, faces, edges and vertices are closed sets.

Proposition 5.2 *Let X be one of $J^{\pm 1}$, $P^{\pm 1}$, $R_1^{\pm 1}$ or $R_2^{\pm 1}$. Then:*

- (i) X sends $S(X)$ bijectively to $S(X^{-1})$ sending vertices, edges and faces to vertices, edges and faces respectively.
- (ii) $X^{-1}(D) \cap D = \emptyset$ and $X^{-1}(\overline{D}) \cap \overline{D} = S(X)$.

PROOF: For each X the side $S(X)$ is contained in a bisector $B(X)$. By the construction of $B(X)$ in Section 4, for each X there are vertices p_a and p_b so that p_a is not on $B(X)$ and $B(X)$ is equidistant from p_a and $X^{-1}(p_b)$. Moreover, D is contained in the half-space closer to p_a than to $X^{-1}(p_b)$.

Applying X we see that $X(B(X))$ is equidistant from $X(p_a)$ and p_b . In other words, $X(p(X)) = B(X^{-1})$. Moreover $X(D)$ is contained in the halfspace closer to $X(p_a)$ than to p_b . This is the opposite halfspace to D . Thus $X(D) \cap D = \emptyset$ and $X(\overline{D}) \cap \overline{D} \subset B(X^{-1})$.

The rest of the proposition follows from the fact that $\overline{D} \cap B(X) = S(X)$, which comes out of our construction. \square

This proposition verifies the side pairing conditions (S.1), (S.2), (S.3) and (S.4) of Poincaré's theorem as given in Section 4.1 of Parker [10]. The condition (S.5) is obvious and (S.6) is vacuous in this case.

5.3 Face cycles

For each face $F(X, Y^{-1}) = S(X) \cap S(Y^{-1})$ we construct an element of Γ called a cycle transformation as follows. Let $X = X_1$. We know from Proposition 5.2 that $X = X_1$ sends faces of $S(X_1)$ to faces of $S(X_1^{-1})$. Suppose that $X_1(F(X_1, Y^{-1})) = F(X_2, X_1^{-1}) = S(X_2) \cap S(X_1^{-1})$, a face of $S(X_1^{-1})$. We can repeat this process, and consider $X_2(F(X_2, X_1^{-1}))$, a face $F(X_3, X_2^{-1})$ of $S(X_2^{-1})$. Eventually, we will find an n so that $X_n = Y$ and $Y(F(Y, X_n^{-1})) = F(X, Y^{-1})$, the first face we considered. In other words we have

$$F(X_1, X_n^{-1}) \xrightarrow{X_1} F(X_2, X_1^{-1}) \xrightarrow{X_2} \dots \xrightarrow{X_{n-1}} F(X_n, X_{n-1}^{-1}) \xrightarrow{X_n} F(X_1, X_n^{-1}).$$

The collection of faces $F(X_i, X_{i-1}^{-1})$, for $i = 1$ to n with $X_0 = X_n$, is called the *face cycle* associated to $F(X_1, X_n^{-1}) = F(X, Y^{-1})$ and the composition $T = X_n \circ \dots \circ X_2 \circ X_1$ is called the *cycle transformation* associated to $F(X, Y^{-1})$.

Proposition 5.3 For each face $F(X_1, X_n^{-1})$ with cycle transformation $T = X_n \circ \dots \circ X_2 \circ X_1$:

- (i) There is an integer ℓ so that the restriction of T^ℓ to $F(X_1, X_n^{-1})$ is the identity.
- (ii) There is an integer m so that $T^{\ell m} = (T^\ell)^m$ is the identity on the whole space.
- (iii) For $i = 0, \dots, n-1$ and $j = 0, \dots, \ell m - 1$ the images $T^{-j} \circ X_1^{-1} \circ \dots \circ X_i^{-1}(D)$ are disjoint. (Here $i = 0$ means the identity and so $T^{-j} \circ X_1^{-1} \circ \dots \circ X_i^{-1}(D) = D$ when $i = j = 0$.)
- (iv) The union over $i = 0, \dots, n-1$ and $j = 0, \dots, \ell m - 1$

$$\bigcup_{i,j} T^{-j} \circ X_1^{-1} \circ \dots \circ X_i^{-1}(\bar{D})$$

covers a neighbourhood of the interior of $F(X_1, X_n^{-1})$.

PROOF: We list the face cycles and the cycle transformation associated to the first face in the cycle and the integers l and m .

Face cycle	Transformation	ℓ	m
$F(J, J^{-1})$	J	3	1
$F(J, P)$	$F(P^{-1}, J^{-1})$	1	k
$F(J, R_1)$	$F(R_2, J^{-1})$	1	1
$F(P, R_1)$	$F(R_2, P^{-1})$	1	1
$F(R_1, R_1^{-1})$	R_1	1	p
$F(R_2, R_2^{-1})$	R_2	1	p
$F(R_1, R_2^{-1})$	$F(P^{-1}, R_1^{-1})$	1	1

This proves (i) and (ii). We now prove (iii) and (iv) in the cases where $F(X_1, X_n^{-1})$ is contained in a Giraud disc, a slice and a meridian. We do a single example in each case.

First consider faces contained in Giraud discs. We will give the details for the face $F(J, J^{-1})$. It is defined by

$$|\langle p, p_{231} \rangle| = |\langle p, J^{-1}(p_{231}) \rangle| = |\langle p, J(p_{231}) \rangle|.$$

There are three sectors around this face, each where one of these quantities is smallest. First, using Corollary 4.2, D is contained in the sector with

$$|\langle p, p_{231} \rangle| < |\langle p, J^{-1}(p_{231}) \rangle|, \quad |\langle p, p_{231} \rangle| < |\langle p, J(p_{231}) \rangle|.$$

Thus $J(D)$ is contained in the sector where

$$|\langle p, J(p_{231}) \rangle| < |\langle p, p_{231} \rangle|, \quad |\langle p, J(p_{231}) \rangle| < |\langle p, J^2(p_{231}) \rangle|.$$

As J has order 3 we see that $J^2 = J^{-1}$. Likewise $J^{-1}(D)$ is contained in the sector where

$$|\langle p, J^{-1}(p_{231}) \rangle| < |\langle p, J(p_{231}) \rangle|, \quad |\langle p, J^{-1}(p_{231}) \rangle| < |\langle p, p_{231} \rangle|.$$

It is easy to see that these three sectors are disjoint and that their closures cover a neighbourhood of $B(J) \cap B(J^{-1})$. Adding the extra inequalities defining the boundary of $F(J, J^{-1})$, we see that D , $J(D)$ and $J^{-1}(D)$ cover a neighbourhood of the interior of $F(J, J^{-1})$. This works because p_{231} is the intersection of the complex spines $\Sigma(J)$ and $\Sigma(J^{-1})$ of $B(J)$ and $B(J^{-1})$. For the other faces in Giraud discs we must modify the inequalities in Corollary 4.2 accordingly.

Now consider the face $F(J, P)$ contained in the complex line $z_1 = 0$. The polyhedron D is contained in the sector where $\arg(z_1) \in (-\phi, 0)$. Also note that D is contained in the sector $0 < \arg(w_1) < \phi$. since $\mathbf{w} = P^{-1}(\mathbf{z})$, we see $P^{-1}(D)$ is contained in the sector $\arg(z_1) < (0, \phi)$. Since $P^{-1}J = A_1$, which multiplies z_1 by $e^{2i\phi}$ we see that $J^{-1}(D) = (P^{-1}J)^{-1}P^{-1}(D)$ is contained in the sector with $\arg(z_1) < (-2\phi, -\phi)$. Continuing in this way we see see:

Image of D	Sector containing $\arg(z_1)$
$(P^{-1}J)^{-j}(D)$	$(-2j - 1)\phi < \arg(z_1) < -2j\phi$
$(P^{-1}J)^{-j}J^{-1}(D)$	$(-2j - 2)\phi < \arg(z_1) < (-2j - 1)\phi$

Since $\phi = 2\pi/k$, clearly these sectors are disjoint and the union as j varies from 0 to $k - 1$ of their closures covers a neighbourhood of $B(J) \cap B(P)$. Adding in the other inequalities shows that a neighbourhood of the interior of $F(J, P)$ is covered by

$$\bigcup_{j=0}^{k-1} \left((P^{-1}J)^{-j}(\overline{D}) \cup (P^{-1}J)^{-j}J^{-1}(\overline{D}) \right).$$

Now consider the face $F(J, R_1)$ contained in the Lagrangian plane where $\text{Im}(z_1 e^{i\phi}) = 0$ and $\text{Im}(z_2) = 0$. Similar arguments to those given above show that D is contained in the quadrant where $\text{Im}(z_1 e^{i\phi}) > 0$ and $\text{Im}(z_2) > 0$; $J^{-1}(D)$ is contained in the quadrant where $\text{Im}(z_1 e^{i\phi}) < 0$ and $\text{Im}(z_2) > 0$; $J^{-1}R_2^{-1}(D)$ is contained in the quadrant where $\text{Im}(z_1 e^{i\phi}) < 0$ and $\text{Im}(z_2) < 0$ and $J^{-1}R_2^{-1}J(D)$ is contained in the quadrant where $\text{Im}(z_1 e^{i\phi}) > 0$ and

$\text{Im}(z_2) < 0$. Arguing as above, we see that these images of D are disjoint and their closures cover a neighbourhood of the interior of $F(J, R_1)$. \square

Therefore we have proved the face conditions (F.1), which was done in Proposition 4.10, (F.2) and (F.3) given in Section 4.1 of Parker [10].

We also have to be careful in the case where one of the vertices is on the ideal boundary. In this case, we must show that there is a *consistent horosphere* about this point; see Epstein and Petronio [4]. (This was not necessary for the groups considered in [10] as the lattices were cocompact.) In other words, there is a horosphere based at this point that is preserved under its stabiliser and mapped off itself by all other group elements. Since we have finitely many faces, it will suffice to show that when a point is on the boundary, its stabiliser is generated by elliptic maps.

- (1) It is clear that p_1 and p_2 are never on the ideal boundary.
- (2) The vertex p_{231} is on the ideal boundary exactly when $\theta = 2\pi/p = \pi/3$ and so $p = 6$. The stabiliser of p_{231} is generated by R_1 and R_2 . Since these maps are elliptic they preserve all horospheres centred at p_{231} .
- (3) The vertices p_{12} , p_{23} and p_{31} lie on the ideal boundary exactly when $\theta + 2\phi = 2\pi/p + 2\pi/k = \pi$ and so (p, k) is one of $(3, 6)$, $(4, 4)$ or $(6, 3)$. The stabiliser of p_{31} is generated by R_2 and A_1 . Since these maps are elliptic they preserve all horospheres centred at p_{231} .

We have verified the hypotheses of Poincaré's polyhedron theorem. Therefore the group generated by the side pairing transformations J, P, R_1 and R_2 is discrete, D is a fundamental polyhedron and the relations are generated by the cycle relations $T^{\ell m} = I$ for each face $F(X, Y^{-1})$ (there are no reflection relations in this case). Therefore the relations are generated by

$$J^3 = (P^{-1}J)^k = R_1^{-1}J^{-1}R_2J = R_1^{-1}P^{-1}R_2P = R_1^p = R_2^p = R_2P^{-1}R_1 = I.$$

It is clear that these may be rewritten to give the relations in (17). Therefore we have proved Theorem 5.1.

References

- [1] Boadi, R.K.; *New Cone Metrics on the Sphere*. PhD thesis, KNUST, 2011.

- [2] Deligne, P. & Mostow, G. D.; *Monodromy of hypergeometric functions and non-lattice integral monodromy*, Publ. Math. IHES **63** (1986) 5–89.
- [3] Deraux, M., Falbel, E & Paupert, J.; *New constructions of fundamental polyhedra in complex hyperbolic space*, Acta Math. **194** (2005). 155–201.
- [4] Epstein, D.B.A., Petronio, C., *An exposition of Poincaré’s polyhedron theorem*, L’Enseignement Mathématique, **40** (1994), 113–170.
- [5] Falbel, E. & Parker, J. R.; *The geometry of the Eisenstein-Picard modular group*, Duke Math. J **131** (2006), 249–289.
- [6] Goldman, W.M.; *Complex Hyperbolic Geometry*. Oxford Mathematical Monographs. Oxford University Press (1999).
- [7] Livné, R.A.; *On Certain Covers of the Universal Elliptic Curve*, PhD thesis, Harvard University, 1981.
- [8] Mostow, G.D.; *On a remarkable class of polyhedra in complex hyperbolic space*, Pacific J. Maths. **86** (1980), 171–276.
- [9] Mostow, G.D.; *Braids, hypergeometric functions and lattices*, Bull. A.M.S. **16** (1987), 225–246.
- [10] Parker, J.R.; *Cone metrics on the sphere and Livné’s lattices*, Acta Math. **196** (2006), 1–64.
- [11] Parker, J.R.; *Complex Hyperbolic Lattices*, in “Discrete Groups and Geometric Structures”, ed: K. Dekimpe, P. Igodt & A. Valette, Contemporary Mathematics **501** AMS, (2009) 1–42.
- [12] Thurston, W.P.; *Shapes of polyhedra and triangulations of the sphere*, In “The Epstein Birthday Schrift”, ed: I. Rivin, C. Rourke & C. Series, Geometry and Topology Monographs **1** (1998), 511–549.
- [13] Weber, M.; *Fundamentaltbereiche komplex hyperbolische Flächen*, Bonner Mathematische Schriften **254** 1993.
- [14] Zhao, T.; *A minimal volume arithmetic cusped complex hyperbolic orbifold*, Math. Proc. Cambridge Philos. Soc. **150** (2011) 313–342.




Figure 10: *The side pairing maps. The solid lines denote edges in the spine of the relevant bisector.*