Dr. Norbert Peyerimhoff, Durham University 28/11/2008

Riemannian Geometry IV

Solutions, set 6.

Exercise 12. $c(t) = (\cos^3(t), \sin^3(t))$ implies that

$$c'(t) = 3\sin t \cos t(-\cos(t), \sin(t)).$$

So we obtain

$$\|c'(t)\| = 3|\sin t \cos t| = \frac{3}{2}|\sin(2t)|,$$

and the length is given by

$$\begin{aligned} L(c) &= \int_0^{2\pi} \|c'(t)\| dt \\ &= \frac{3}{2}\left(\int_0^{\pi/2} \sin(2t)dt - \int_{\pi/2}^{\pi} \sin(2t)dt + \int_{\pi}^{\frac{3\pi}{2}} \sin(2t)dt - \int_{\frac{3\pi}{2}}^{2\pi} \sin(2t)dt\right) \\ &= \frac{3}{2} \cdot 4 \cdot \int_0^{\pi/2} \sin(2t)dt = 6. \end{aligned}$$

Exercise 13. (a) We have $c'(t) = i$ for all $t \in [a, b]$. The function $l : [a, b] \to [0, L(c)]$ is given by

$$l(t) = \int_a^t \|c'(s)\|_{c(s)}\, ds = \ln \frac{t}{a}.$$

So $l : [a, b] \to [0, \ln(b/a)]$ is bijective, strictly monotone increasing and differentiable. We calculate its inverse:

$$s = l(t) \Leftrightarrow s = \ln \frac{t}{a} \Leftrightarrow e^s = \frac{t}{a} \Leftrightarrow t = ae^s.$$

So $l^{-1}(s) = ae^s$ and an arc length reparametrization of c is given by $\gamma = c \circ l^{-1} : [0, \ln(b/a)] \to \mathbb{H}^2$,

$$\gamma(s) = c(l^{-1}(s)) = c(ae^s) = ae^s i.$$

(b) We have

$$c(t) = \frac{(ai\cos t + \sin t)(ai\sin t + \cos t)}{\cos^2 t + a^2\sin^2 t} = \frac{\sin t\cos t(1-a^2) + ia}{\cos^2 t + a^2\sin^2 t},$$

so

$$\mathrm{Im}(c(t)) = \frac{a}{\cos^2 t + a^2\sin^2 t}.$$

On the other hand, we have

$$c'(t) = \frac{(-ai\sin t + \cos t)^2 + (ai\cos t + \sin t)^2}{(-ai\sin t + \cos t)^2} = \frac{1-a^2}{(-ai\sin t + \cos t)^2}.$$

This implies that

$$|c'(t)| = \frac{a^2-1}{\cos^2 t + a^2\sin^2 t},$$

and

$$\|c'(t)\|_{c(t)} = \frac{a^2-1}{\cos^2 t + a^2\sin^2 t}\frac{\cos^2 t + a^2\sin^2 t}{a} = \frac{a^2-1}{a} = a - \frac{1}{a}.$$

So we obtain

$$L(c) = \int_0^\pi \|c'(t)\|_{c(t)}\,dt = \pi\left(a - \frac{1}{a}\right).$$

Exercise 14.

(a) If $z_1 = z_2$ there is nothing to check. Let $z_1 = ai$ and $z_2 = bi$. In this case we have

$$d_{\mathbb{H}}(z_1, z_2) = |\ln b/a|,$$

which implies that

$$\sinh(\frac{1}{2}d(z_1,z_2)) = \frac{1}{2}\Big|\exp(\frac{\ln(b/a)}{2}) - \exp(-\frac{\ln(b/a)}{2})\Big| = \frac{1}{2}\Big|\sqrt{\frac{a}{b}} - \sqrt{\frac{b}{a}}\Big|.$$

On the other hand we obtain

$$\frac{|z_1 - z_2|}{2\sqrt{\mathrm{Im}(z_1)\mathrm{Im}(z_2)}} = \frac{|a-b|}{2\sqrt{ab}} = \frac{1}{2}\Big|\frac{a}{\sqrt{ab}} - \frac{b}{\sqrt{ab}}\Big|,$$

which shows the validity of the formula in this particular case.

(b) Since f_A is an isometry (see Exercise 9), we have $d(f_A(z_1), f_A(z_2)) = d(z_1, z_2)$. This immediately implies invariance of the left hand side under f_A. As for the right-hand side note first that

$$|f_A(z_1) - f_A(z_2)| = \frac{|(az_1+b)(cz_2+d) - (az_2+b)(cz_1+d)|}{|cz_1+d|\,|cz_2+d|}$$
$$= \frac{|z_1 - z_2|}{|cz_1+d|\,|cz_2+d|}.$$

Using the identity

$$\mathrm{Im}(f_A(z)) = \frac{\mathrm{Im}(z)}{|cz+d|^2},$$

we obtain

$$\frac{|f_A(z_1) - f_A(z_2)|}{2\sqrt{\mathrm{Im}(f_A(z_1))\mathrm{Im}(f_A(z_2))}} = \frac{|z_1 - z_2|}{|cz_1+d|\,|cz_2+d|} \cdot \frac{|cz_1+d|\,|cz_2+d|}{2\sqrt{\mathrm{Im}(z_1)\mathrm{Im}(z_2)}}$$
$$= \frac{|z_1 - z_2|}{2\sqrt{\mathrm{Im}(z_1)\mathrm{Im}(z_2)}}.$$

(c) The map $f(z) = z - x$ coincides with the map f_A for $A = \begin{pmatrix} 1 & -x \\ 0 & 1 \end{pmatrix} \in \mathrm{SL}(2,\mathbb{R})$. So we can apply (b). Note that the points $w_1 = f(z_1), w_2 = f(z_2)$ satisfy the requirements in (a). We conclude that

$$\begin{aligned}
\sinh(\tfrac{1}{2}d(z_1,z_2)) &= \sinh(\tfrac{1}{2}d(w_1,w_2)) && \text{by (b)} \\
&= \frac{|w_1 - w_2|}{2\sqrt{\mathrm{Im}(w_1)\mathrm{Im}(w_2)}} && \text{by (a)} \\
&= \frac{|z_1 - z_2|}{2\sqrt{\mathrm{Im}(z_1)\mathrm{Im}(z_2)}} && \text{by (b).}
\end{aligned}$$

(d) Using $c(t) = x + Re^{it}$ we calculate

$$f(c(t)) = \frac{R(e^{it} - 1)}{R(e^{it} + 1)} = i\frac{\sin(t/2)}{\cos(t/2)} = i\tan(t/2).$$

As t runs from 0 to π, $i\tan(t/2)$ runs from the origin along the positive imaginary axis to infinity.

Note that $f(z)$ coincides with the map f_A for

$$A = \begin{pmatrix} \frac{1}{\sqrt{2R}} & -\frac{x+r}{\sqrt{2R}} \\ \frac{1}{\sqrt{2R}} & -\frac{x-R}{\sqrt{2R}} \end{pmatrix} \in \mathrm{SL}(2, \mathbb{R}).$$

So we can apply (b). Note that the points $w_1 = f(z_1), w_2 = f(z_2)$ satisfy the requirements in (a). We conclude that

$$\begin{aligned} \sinh(\frac{1}{2}d(z_1, z_2)) &= \sinh(\frac{1}{2}d(w_1, w_2)) && \text{by (b)} \\ &= \frac{|w_1 - w_2|}{2\sqrt{\mathrm{Im}(w_1)\mathrm{Im}(w_2)}} && \text{by (a)} \\ &= \frac{|z_1 - z_2|}{2\sqrt{\mathrm{Im}(z_1)\mathrm{Im}(z_2)}} && \text{by (b).} \end{aligned}$$