


Topics in Combinatorics IV, Solutions 17 (Week 17)

17.1. (★) Draw the Hasse diagram of the root poset of root system A_4 .

Solution: Every root $e_i - e_j$ for $i < j$ can be written as

$$e_i - e_j = \sum_{k=i}^{j-1} e_k - e_{k+1} = \sum_{k=i}^{j-1} \alpha_k$$

which implies that $e_i - e_j \geq e_k - e_l$ if and only if $i \leq k$ and $j \geq l$. We get the following Hasse diagram:



17.2. (★) Let (W, S) be a Coxeter system. A subgroup H of W is a *parabolic subgroup* if it is conjugated to a standard parabolic subgroup W_T for some $T \subset S$ (see HW 15.2), i.e. $H = w^{-1}W_T w$ for some $w \in W$. Show that for any $p \in \mathbb{R}^n$ the stabilizer $\text{Stab}_W(p)$ is a parabolic subgroup.

Solution: This follows from HW 16.4. Let C be the initial chamber of W . There exists $w \in W$ such that $wp \in \overline{C}$. By HW 16.4, the stabilizer of wp is a standard parabolic subgroup W_T , where $s \in T$ if and only if $s(wp) = wp$. Therefore, the stabilizer of p is precisely $w^{-1}W_T w$.

17.3. Let (W, S) be an irreducible Coxeter system. Denote $c_n = \#\{w \in W \mid l(w) = n\}$, and define the generating function

$$W(q) = \sum_{n \geq 0} c_n q^n = \sum_{w \in W} q^{l(w)},$$

which is called the *Poincaré series* of W . In the case when W is finite, $W(q)$ is called the *Poincaré polynomial* of W .

Recall that if $T \subset S$ then W_T denotes a standard parabolic subgroup, and $W^T = \{w \in W \mid l(wt) > l(w) \forall t \in T\}$ (see HW 16.2).

For every $X \subset W$ denote also $X(q) = \sum_{w \in X} q^{l(w)}$.

- (a) Show that if $T \subset S$ then $W(q) = W_T(q)W^T(q)$.
- (b) Let $w \in W$, define $F = F(w) = \{s \in S \mid l(ws) > l(w)\}$. Show that $\sum_{T \subset F} (-1)^{|T|} = 0$ unless W is finite and $w = w_0$ is the longest element of W .
- (c) Show that

$$\sum_{T \subset S} (-1)^{|T|} \frac{W(q)}{W_T(q)} = \sum_{T \subset S} (-1)^{|T|} W^T(q) = \begin{cases} 0 & \text{if } W \text{ is infinite,} \\ q^N & \text{if } W \text{ is finite,} \end{cases}$$

where N is the length of the longest element of W .

- (d) Assume W is finite. Show that

$$\sum_{T \subset S} (-1)^{|T|} \frac{|W|}{|W_T|} = 1$$

- (e) Apply the formula from (d) to compute the order of the group H_3 . Can you compute the order of H_4 in this way?

Solution:

- (a) This follows from HW 16.2: take any $w \in W$, then there are unique $u \in W_T$ and $v \in W^T$ such that $w = uv$. Since $l(w) = l(u) + l(v)$, we have $q^{l(w)} = q^{l(u)}q^{l(v)}$, so

$$W(q) = \sum_{w \in W} q^{l(w)} = \sum_{u \in W_T, v \in W^T} q^{l(u)}q^{l(v)} = \sum_{u \in W_T} q^{l(u)} \sum_{v \in W^T} q^{l(v)} = W_T(q)W^T(q)$$

- (b) Suppose F is empty, then $l(ws) < l(w)$ for any $s \in S$, and thus W is finite and $w = w_0$. Otherwise, F is non-empty, and the statement claims that the alternating sum of binomial coefficients is zero, which follows from the identity $(1 - 1)^{|F|} = 0$.
- (c) The first equality follows from (a). To prove the second equality, write

$$\sum_{T \subset S} (-1)^{|T|} W^T(q) = \sum_{T \subset S} (-1)^{|T|} \left(\sum_{w \in W^T} q^{l(w)} \right)$$

and observe that $w \in W^T$ if and only if $T \subset F(w)$. Therefore, for a given w the coefficient of $q^{l(w)}$ is precisely $\sum_{T \subset F(w)} (-1)^{|T|}$, which vanishes for every $w \in W$ unless W is finite due to

(b). If W is finite, then the only $w \in W$ such that $F(w)$ is empty is w_0 , and thus the only non-zero term is $q^{l(w_0)}$.

- (d) Plug in $q = 1$ in (c).
- (e) Let $H_3 = \langle s_1, s_2, s_3 \mid s_i^2, (s_1 s_2)^3, (s_2 s_3)^5, (s_1 s_3)^2 \rangle$. Then we have the following subsets of S and the orders of the standard parabolic subgroups:

T	\emptyset	$\{s_1\}$	$\{s_2\}$	$\{s_3\}$	$\{s_1, s_2\}$	$\{s_1, s_3\}$	$\{s_2, s_3\}$	$\{s_1, s_2, s_3\}$
W_T	$\{e\}$	A_1	A_1	A_1	A_2	$A_1 \times A_1$	$I_2(5)$	H_3
$ W_T $	1	2	2	2	6	4	10	$ W(H_3) $

Therefore, we have

$$|W| \left(\frac{1}{1} - \frac{1}{2} - \frac{1}{2} - \frac{1}{2} + \frac{1}{6} + \frac{1}{4} + \frac{1}{10} \right) - 1 = 1,$$

which is equivalent to $|W(H_3)| \cdot \frac{2}{120} = 2$, so we get $|W(H_3)| = 120$.

The group $W(H_4)$ has rank 4, which implies that the summand $\frac{|W|}{|W|}$ in the sum will have coefficient +1. Therefore, the formula in (c) will become $|W| \cdot (\dots) = 0$, which cannot be used for computation of the order.