

## Differential Geometry III, Homework 5 (Week 5)

Due date for starred problems: **Thursday, November 21.**

### Space curves - 2

**5.1.** (★) A curve  $\alpha : I \rightarrow \mathbb{R}^3$  is called a (*generalized*) *helix* if its tangent lines make a constant angle with a fixed direction in  $\mathbb{R}^3$ . Suppose that  $\tau(s) \neq 0$  for all  $s \in I$ .

(a) Prove that the curve

$$\alpha(s) = \left( \frac{a}{c} \int_{s_0}^s \sin \vartheta(v) \, dv, \frac{a}{c} \int_{s_0}^s \cos \vartheta(v) \, dv, \frac{b}{c} s \right),$$

with  $s_0 \in I$ ,  $c^2 = a^2 + b^2$ ,  $a \neq 0$ ,  $b \neq 0$  and  $\vartheta'(s) > 0$  is a (*generalized*) helix.

(b) Prove that  $\alpha$  is a (*generalized*) helix if and only if  $\kappa/\tau$  is constant.

**5.2.** Let  $\alpha, \beta$  be regular curves in  $\mathbb{R}^3$  such that, for each  $u$ , the principal normals  $\mathbf{n}_\alpha(u)$  and  $\mathbf{n}_\beta(u)$  are parallel. Prove that the angle between  $\mathbf{t}_\alpha(u)$  and  $\mathbf{t}_\beta(u)$  is independent of  $u$ . Prove also that if the line through  $\alpha(u)$  in direction  $\mathbf{n}_{\alpha(u)}$  coincides with the line through  $\beta(u)$  in direction  $\mathbf{n}_{\beta(u)}$  then

$$\beta(u) = \alpha(u) + r\mathbf{n}_\alpha(u)$$

for some real number  $r$ .

**5.3.** (★) Let  $\alpha$  be the curve in  $\mathbb{R}^3$  given by

$$\alpha(u) = e^u(\cos u, \sin u, 1), \quad u \in \mathbb{R}.$$

If  $0 < \lambda_0 < \lambda_1$ , find the length of the segment of  $\alpha$  which lies between the planes  $z = \lambda_0$  and  $z = \lambda_1$ . Show also that the curvature and torsion of  $\alpha$  are both inversely proportional to  $e^u$ .

**5.4.** Let  $\alpha$  be a curve parametrized by arc length with nowhere vanishing curvature  $\kappa$  and torsion  $\tau$ . Show that if the trace of  $\alpha$  lies on a sphere then

$$\frac{\tau}{\kappa} = \left( \frac{\kappa'}{\tau\kappa^2} \right)'$$

Is the converse true?

**5.5.** Let  $\alpha$  be a regular curve parametrized by arc length with  $\kappa > 0$  and  $\tau \neq 0$ . Denote by  $\mathbf{n}$  and  $\mathbf{b}$  the principal normal and the binormal of  $\alpha$ .

(a) If  $\alpha$  lies on a sphere of centre  $\mathbf{c} \in \mathbb{R}^3$  and radius  $r > 0$ , show that

$$\alpha - \mathbf{c} = -\rho\mathbf{n} - \rho'\sigma\mathbf{b},$$

where  $\rho = 1/\kappa$  and  $\sigma = -1/\tau$ . Deduce that  $r^2 = \rho^2 + (\rho'\sigma)^2$ .

(b) Conversely, if  $\rho^2 + (\rho'\sigma)^2$  has constant value  $r^2$  and  $\rho' \neq 0$ , show that  $\alpha$  lies on a sphere of radius  $r$ .

*Hint:* Show that the curve  $\alpha + \rho\mathbf{n} + \rho'\sigma\mathbf{b}$  is constant.