

## Topics in Combinatorics IV, Term 1<sup>1</sup>

### 1 Catalan numbers

#### 1.1 Definitions

**Definition 1.1** (one of many). The  $n$ -th Catalan number  $C_n$  is the number of sequences  $(\varepsilon_1, \dots, \varepsilon_{2n})$  with  $\varepsilon_i = \pm 1$  such that

$$\begin{aligned} & \cdot \sum_{i=1}^{2n} \varepsilon_i = 0; \\ & \cdot \sum_{i=1}^k \varepsilon_i \geq 0 \text{ for every } k \leq 2n. \end{aligned}$$

**Remark.** Sequences in the definition above are called *ballot sequences*.

**Example 1.2.**  $n = 2$ : the only sequences of length  $2n = 4$  are  $(1, 1, -1, -1)$  and  $(1, -1, 1, -1)$ , so  $C_2 = 2$ .  
 $n = 3$ : there are five sequences (list them!), so  $C_3 = 5$ .

Two equivalent definitions of  $C_n$ :

- the number of “bracketings” of a non-associative product of  $n + 1$  variables;
- the number of triangulations of a convex  $(n + 2)$ -gon on a plane (here by a triangulation we mean a maximal collection of non-crossing diagonals, it automatically subdivides the polygon into triangles).

**Example 1.3.** For  $n = 3$ , there are precisely five bracketings:

$$((a_1 a_2) a_3) a_4 \quad (a_1 (a_2 a_3)) a_4 \quad (a_1 a_2) (a_3 a_4) \quad a_1 ((a_2 a_3) a_4) \quad a_1 (a_2 (a_3 a_4)).$$

There are also precisely five triangulations of a pentagon: two non-crossing diagonals share a vertex, and there are five vertices to choose from.

**Exercise 1.4.** Show that the two definitions above are equivalent to Definition 1.1.

Ballot sequences can be represented by *Dyck paths*: these are paths in a  $n \times n$  square going along the grid from one corner (say,  $(0, n)$ ) to the opposite (i.e.,  $(n, 0)$ ) and staying above the main diagonal. The bijection with ballot sequences is obvious:  $\varepsilon_i = 1$  becomes a move to the right, and  $\varepsilon_i = -1$  becomes a move down.

One can draw the top right half of the square only, in this case a Dyck path can be understood as a path from  $(0, 0)$  to  $(2n, 0)$  with steps  $(1, 1)$  and  $(1, -1)$  never going below  $x$ -axis. This is the model we will usually use in the sequel.

**Remark.** Paths whose all steps are vectors of  $\mathbb{Z}^d$  are called *lattice paths*. The two models above for Dyck paths sit in  $\mathbb{Z}^2$  generated, respectively, by  $(1, 0)$ ,  $(0, -1)$ , and  $(1, 1)$ ,  $(1, -1)$ .

---

<sup>1</sup>Based on Alex Postnikov’s notes

**Example 1.5.**




Figure 1.1: Dyck paths with steps (a)  $(1, 0)$  and  $(0, -1)$  and (b)  $(1, 1)$  and  $(1, -1)$ .

**1.2 Explicit formula for  $C_n$**

**Theorem 1.6.**  $C_n = \frac{1}{n+1} \binom{2n}{n}$ .

**Example 1.7.**  $C_3 = \frac{1}{4} \binom{6}{3} = \frac{1}{4} \frac{6!}{3!3!} = \frac{6 \cdot 5 \cdot 4}{4 \cdot 6} = 5$ ;  $C_4 = \frac{1}{5} \binom{8}{4} = \frac{1}{5} \frac{8!}{4!4!} = \frac{8 \cdot 7 \cdot 6 \cdot 5}{5 \cdot 8 \cdot 3} = 14$ ;  $C_5 = \frac{1}{6} \binom{10}{5} = 42$ .

We will look at three different proofs of the theorem.

*Proof 1: by reflection.* The number of all lattice paths from  $(0, 0)$  to  $(2n, 0)$  with steps  $(1, 1)$  and  $(1, -1)$  is equal to  $\binom{2n}{n}$  – choose  $n$  steps “up” out of  $2n$  steps. We will compute the number of ones which go below the  $x$ -axis (call them *bad*) and subtract.

Observe that a path is bad if and only if it intersects the line  $y = -1$ . Find the first point where it touches the line  $y = -1$ , and reflect the part of the path to the right of this point with respect to the line  $y = -1$ . We get a new lattice path, it goes from  $(0, 0)$  to  $(2n, -2)$ . We now claim that this map establishes a bijection between *bad* paths from  $(0, 0)$  to  $(2n, 0)$  and *all* lattice paths from  $(0, 0)$  to  $(2n, -2)$ . Indeed, this map is injective, and there is an inverse: take any lattice path from  $(0, 0)$  to  $(2n, -2)$ , take the first point where it touches the line  $y = -1$ , and reflect the right part – we get a bad path.

Now, the paths from  $(0, 0)$  to  $(2n, -2)$  contain  $n - 1$  steps up and  $n + 1$  steps down, so the number of all paths from  $(0, 0)$  to  $(2n, -2)$  is  $\binom{2n}{n+1}$ . Thus,

$$C_n = \binom{2n}{n} - \binom{2n}{n+1} = \frac{(2n)!}{n!n!} - \frac{(2n)!}{(n-1)!(n+1)!} = \frac{(2n)!}{n!n!} - \frac{n}{n+1} \frac{(2n)!}{n!n!} = \frac{1}{n+1} \binom{2n}{n}$$

□

*Proof 2: by cyclic shifts.* First, observe that

$$\frac{1}{n+1} \binom{2n}{n} = \frac{(2n)!}{n!(n+1)!} = \frac{1}{2n+1} \frac{(2n+1)!}{n!(n+1)!} = \frac{1}{2n+1} \binom{2n+1}{n}$$

Interpret  $\binom{2n+1}{n}$  as all lattice paths from  $(0, 0)$  to  $(2n+1, -1)$ , i.e. paths with  $n$  steps up and  $n+1$  steps down. Then  $C_n$  is the number of those paths (call them *good*) that go below the  $x$ -axis at the last step only. We want to show that good paths constitute precisely  $1/(2n+1)$  part of all paths.

Let us switch to sequences of  $\pm 1$ . Take any sequence  $(\varepsilon_1, \dots, \varepsilon_{2n+1})$  of length  $2n+1$  adding up to  $-1$ , and consider all its  $2n+1$  *cyclic shifts*:

$$(\varepsilon_1, \dots, \varepsilon_{2n+1}), (\varepsilon_2, \varepsilon_3, \dots, \varepsilon_{2n+1}, \varepsilon_1), (\varepsilon_3, \dots, \varepsilon_{2n+1}, \varepsilon_1, \varepsilon_2), \dots, (\varepsilon_{2n+1}, \varepsilon_1, \dots, \varepsilon_{2n}).$$

Now the proof follows from the following statement which we leave as an exercise

**Exercise.** (1) Show that all cyclic shifts are distinct.

(2) Show that out of  $2n + 1$  shifts of one sequence precisely one is good. (Hint: consider the leftmost lowest point of the path). □

**Definition 1.8.** Let  $(a_n)_{n \in \mathbb{Z}_{\geq 0}}$  be a sequence of non-negative integers. A formal power series  $\sum_{k=0}^{\infty} a_k x^k$  is called a *generating function* of  $(a_n)$ .

**Example 1.9.** Let  $a_n = 1$  for every  $n$ . The the generating function is  $A(x) = 1 + x + \dots = 1/(1 - x)$ .

In general, how to find a closed formula for the generating function  $A(x)$  of a sequence  $(a_n)$ ? One can follow the following plan:

- write a recurrence relation on  $a_n$ ;
- interpret the recurrence relation as an equation on  $A(x)$ ;
- solve the equation and get an explicit expression for  $a_n$ .

We will proceed along this plan to get a *third proof* of the theorem.

**Lemma 1.10** (Recurrence on Catalan numbers).

$$C_n = C_0 C_{n-1} + C_1 C_{n-2} + \dots + C_{n-1} C_0 = \sum_{k=1}^n C_{k-1} C_{n-k}$$

*Proof.* Take any Dyck path of length  $2n$ , and consider the first point  $2k > 0$  where the path touches the  $x$ -axis. Then on the right there is a Dyck path of length  $2n - 2k = 2(n - k)$ . On the left there is a Dyck path of length  $2k$  which stays strictly above  $x$ -axis, which means that we can think of it as a Dyck path of length  $2k - 2$  between points  $(1, 1)$  and  $(2k - 1, 1)$ . Both Dyck paths on the left and on the right are arbitrary, so there are precisely  $C_{k-1} \cdot C_{n-k}$  Dyck paths which touch  $x$ -axis for the first time at  $2k$ , and the result follows. □

Defining  $C_0 = 1$ , we can now recursively compute any Catalan number.

**Example 1.11.**

$$C_1 = C_0 \cdot C_0 = 1; \quad C_2 = C_0 \cdot C_1 + C_1 \cdot C_0 = 1 + 1 = 2; \quad C_3 = C_0 \cdot C_2 + C_1 \cdot C_1 + C_2 \cdot C_0 = 2 + 1 + 2 = 5.$$

**Lemma 1.12.** *The generating function  $C(x)$  satisfies the equation  $x C(x)^2 - C(x) + 1 = 0$ .*

*Proof.* We know that  $C_n = \sum_{k=1}^n C_{k-1} C_{n-k}$ . Multiplying by  $x^n$ , we get

$$C_n x^n = \sum_{k=1}^n C_{k-1} C_{n-k} x^n = x \sum_{k=1}^n (C_{k-1} x^{k-1}) (C_{n-k} x^{n-k})$$

Now, summing on  $n > 0$ , we get on the left  $\sum_{n=1}^{\infty} C_n x^n = C(x) - C_0 = C(x) - 1$ . On the right, we get

$$\begin{aligned} \sum_{n=1}^{\infty} \sum_{k=1}^n x(C_{k-1}x^{k-1})(C_{n-k}x^{n-k}) &= x \sum_{n=1}^{\infty} \sum_{i=0}^{n-1} (C_i x^i)(C_{n-i-1}x^{n-i-1}) = \\ &= x \sum_{m=0}^{\infty} \sum_{i=0}^m (C_i x^i)(C_{m-i}x^{m-i}) = x \cdot C(x) \cdot C(x), \end{aligned}$$

so we obtain the equation  $C(x) - 1 = xC(x)^2$ , which is precisely what we wanted.  $\square$

**Lemma 1.13.**

$$C(x) = \frac{1 - \sqrt{1 - 4x}}{2x}$$

*Proof.* Solving quadratic equation with respect to  $C(x)$  (while considering  $x$  as a parameter), we see that  $C(x) = \frac{1 \pm \sqrt{1 - 4x}}{2x}$ . The reason to choose the sign is the following: we want  $\lim_{x \rightarrow 0} C(x) = C(0)$ , where we already know  $C(0) = C_0 = 1$ . Now, if we consider the positive sign, then  $\lim_{x \rightarrow 0} C(x) = \infty$ , while the negative sign gives the required limit.  $\square$

We are left to extract the explicit expression for  $C_n$  from the generating function. For this, we recall the definition of a generalization of binomial coefficients.

**Definition 1.14.** Given  $\alpha \in \mathbb{R}$  and  $k \in \mathbb{N}$ , the *binomial coefficient*  $\binom{\alpha}{k}$  is given by

$$\binom{\alpha}{k} = \frac{\alpha(\alpha - 1)(\alpha - 2) \dots (\alpha - k + 1)}{k!}$$

**Exercise.** Show that

$$(1 + y)^\alpha = \sum_{k \geq 0} \binom{\alpha}{k} y^k$$

We now apply the exercise above to  $\alpha = 1/2$ ,  $y = -4x$ . We get

$$\sqrt{1 - 4x} = \sum_{k \geq 0} \binom{\frac{1}{2}}{k} (-4x)^k = 1 + \binom{\frac{1}{2}}{1} (-4x) + \binom{\frac{1}{2}}{2} (-4x)^2 + \dots,$$

so

$$C(x) = \frac{1 - \sqrt{1 - 4x}}{2x} = -\frac{1}{2} \left( \binom{\frac{1}{2}}{1} (-4) + \binom{\frac{1}{2}}{2} (-4)^2 x + \dots \right) = -\frac{1}{2} \left( \sum_{n \geq 0} \binom{\frac{1}{2}}{n+1} (-4)^{n+1} x^n \right).$$

Therefore,

$$\begin{aligned} C_n &= -\frac{1}{2} \binom{\frac{1}{2}}{n+1} (-4)^{n+1} = -\frac{1}{2} \frac{\frac{1}{2}(\frac{1}{2} - 1)(\frac{1}{2} - 2) \dots (\frac{1}{2} - n)}{(n+1)!} (-4)^{n+1} = \\ &= -\frac{1}{2} \frac{\frac{1}{2}(-\frac{1}{2})(-\frac{3}{2}) \dots (-\frac{2n-1}{2})}{(n+1)!} (-4)^{n+1} = \frac{1 \cdot 3 \cdot \dots \cdot (2n-1)}{(n+1)!} \frac{2^{2n+2}}{2^{n+2}} = \frac{(2n)!}{(2^n n!)(n+1)!} \cdot 2^n = \frac{1}{n+1} \binom{2n}{n} \end{aligned}$$

as required.

### 1.3 Further examples and Applications

**Example 1.15** (Drunkard's walk). A drunkard walks randomly along a line in steps of length 1 to the left or right with probability  $1/2$ . The starting point is  $x = 1$ . The walk is terminated when the drunkard reaches the point  $x = 0$ . Question: what is the probability the walk terminates?

Reaching  $x = 0$  after  $2k + 1$  steps is equivalent to returning to  $x = 1$  after  $2k$  steps and then going left, so this can be understood as a Dyck path. The probability of any individual path is  $(1/2)^{2k+1}$ , the number of such paths of length  $2k + 1$  is clearly  $C_k$ . Thus,

$$P = \sum_{k=0}^{\infty} C_k \frac{1}{2^{2k+1}} = \frac{1}{2} \sum_{k=0}^{\infty} C_k \frac{1}{2^{2k}} = \frac{1}{2} \sum_{k=0}^{\infty} C_k \left(\frac{1}{4}\right)^k = \frac{1}{2} C\left(\frac{1}{4}\right) = \frac{1}{2} \frac{1 - \sqrt{1 - 4 \cdot \frac{1}{4}}}{2 \cdot \frac{1}{4}} = 1$$

We list below some further interpretations of Catalan numbers  $C_n$ .

- The number of *plane binary trees* with  $n$  vertices.

A plane binary tree can be defined recursively: if not empty, it has a root vertex, a left subtree, and a right subtree, both of which are binary trees (either of them can be empty). When drawing such trees, the root is drawn on the top, with an edge drawn from it to the root of each of its subtrees. All binary trees with 5 vertices are shown in Fig. 1.2.



Figure 1.2: Plane binary trees with 3 vertices

- The number of *complete plane binary trees* with  $n + 1$  leaves.

A plane binary tree is complete if it has no vertices of valence 2, i.e., for every vertex both left and right subtrees are either simultaneously nonempty or empty (in the latter case the vertices are called *leaves*). All complete binary trees with 4 leaves are shown in Fig. 1.3.




Figure 1.3: Complete plane binary trees with 4 leaves

- The number of *plane trees* with  $n + 1$  vertices.

A plane tree can also be defined recursively: it has a root vertex, and, in the case the whole tree is not a single vertex, it has a sequence  $(T_1, \dots, T_k)$  of subtrees  $T_i$ ,  $1 \leq i \leq k$ , each of which is also a plane tree. In particular, the subtrees of each vertex are ordered; when drawing such trees, the subtrees are drawn from left to right. The root is on the top, with an edge drawn from it to the root of each of its subtrees. All plane trees with 4 vertices are shown in Fig. 1.4.




Figure 1.4: Plane trees with 4 vertices

- The number of *non-crossing matchings* on  $2n$  vertices.

A matching on  $2n$  vertices can be drawn as a way to connect  $2n$  nodes on the  $x$ -axis by arcs, with every node connected to precisely one other node by an arc drawn in the upper halfplane. A matching is *non-crossing* if all arcs can be drawn in a way such that no pair of arcs intersects.

All non-crossing matchings on 6 vertices are shown in Fig. 1.5.




Figure 1.5: Non-crossing matchings on 6 vertices

- The number of *non-nesting matchings* on  $2n$  vertices.

A matching is *non-nesting* if all the arcs can be drawn in a way such that no arc is above the other, or, equivalently, there is no pair of arcs with ends at points  $(a_1, b_1)$  and  $(a_2, b_2)$  such that  $a_1 < a_2 < b_2 < b_1$ .

All non-nesting matchings on 6 vertices are shown in Fig. 1.6.



Figure 1.6: Non-nesting matchings on 6 vertices

## 2 Partitions and Young Diagrams

### 2.1 Definitions

**Definition 2.1.** A *partition* of  $n \in \mathbb{N}$  is a sequence of integers  $\lambda = (\lambda_1, \dots, \lambda_k)$  such that  $\sum_{i=1}^k \lambda_i = n$  and  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k > 0$ . If  $(\lambda_1, \dots, \lambda_k)$  is a partition of  $n$ , we denote  $(\lambda_1, \dots, \lambda_k) \vdash n$ .

**Example.**  $4 = 3 + 1 = 2 + 2 = 2 + 1 + 1 = 1 + 1 + 1 + 1$ .

Graphically, a partition can be represented by a *Young diagram* which is a left-justified array of boxes arranged in  $k$  rows, such that  $i$ -th row contains  $\lambda_i$  boxes, see Fig. 2.1, left, for an example.




Figure 2.1: Young diagram of the partition  $\lambda = (5, 3, 2, 2) \vdash 12$  and a standart Young tableau of type  $\lambda$

**Definition 2.2.** A *standard Young tableau* (SYT for short) is a filling of a Young diagram of type  $\lambda = (\lambda_1, \dots, \lambda_k) \vdash n$  by numbers  $1, \dots, n$  such that each number appears precisely once, and entries increase in rows (to the right) and in columns (downwards).

See Fig. 2.1, right, for an example.

**Example 2.3.** Let us compute the number of SYT of shape  $(n, n)$ . One can assign to any such SYT a Dyck path: put a step up at every place appearing in the first row, and a step down at every place appearing in the second row. It is easy to see that this is a bijection, so the number of SYT of shape  $(n, n)$  is just the Catalan number  $C_n$ .

Denote by  $f_\lambda$  the number of SYT of type  $\lambda$  (for example,  $f_{(n,n)} = C_n$  as we have seen above). Our next goal is to obtain a formula for  $f_\lambda$ .

## 2.2 The Hook Length Formula

**Definition 2.4.** Let  $(i, j)$  be a box in a Young diagram  $\lambda$ . The *hook* of  $(i, j)$  is the union of all boxes in  $i$ -th row to the right of  $(i, j)$  and all boxes in  $j$ -th column to the bottom of  $(i, j)$ , i.e.,  $\{(i, b) \mid b \geq j\} \cup \{(a, j) \mid a \geq i\}$ . The *hook length*  $h(i, j)$  is the number of boxes in the hook of  $(i, j)$ .

An example is shown in Fig. 2.2.




Figure 2.2: The hook of box  $(2, 2)$  is shaded,  $h(2, 2) = 8$

Denote  $H(\lambda) = \prod_{(i,j) \in \lambda} h(i, j)$ .

**Theorem 2.5** (The Hook Length Formula). *For every partition  $\lambda \vdash n$ ,  $f_\lambda = \frac{n!}{H(\lambda)}$ .*

**Example 2.6.** · Let  $\lambda = (4, 2, 2) \vdash 8$ . Then  $H(\lambda) = (6 \cdot 5 \cdot 2 \cdot 1) \cdot (3 \cdot 2) \cdot (2 \cdot 1) = 6!$ , so  $f_\lambda = \frac{8!}{6!} = 56$ .

· Let  $\lambda = (n, n)$ . Then  $f(1, j) = n - j + 2$  and  $f(2, j) = n - j + 1$ . Therefore,

$$H(\lambda) = ((n + 1) \cdot \dots \cdot 2) \cdot (n \cdot \dots \cdot 1) = (n + 1)!n!,$$

and thus

$$f_{(n,n)} = \frac{(2n)!}{(n + 1)!n!} = C_n$$

Let us first prove a recurrence relation on  $f_\lambda$ . Let  $\lambda$  be a Young diagram, and let  $c$  be a *corner* of  $\lambda$ , i.e. a box which does not have neighbors from right and bottom. Denote by  $\lambda - c$  a Young diagram obtained from  $\lambda$  by removing the box  $c$ .

**Lemma 2.7.** Define  $f_\emptyset = 1$ . Then

$$f_\lambda = \sum_{c \text{ is a corner of } \lambda} f(\lambda - c)$$

*Proof.* Observe that  $n$  is always in the corner of any SYT of shape  $\lambda \vdash n$ . If  $c$  is a corner of  $\lambda \vdash n$ , there is a clear bijection between SYT of shape  $\lambda$  with  $n$  located at box  $c$  and SYT of shape  $\lambda - c \vdash (n - 1)$ . Therefore,

$$f_\lambda = \sum_{c \text{ is a corner of } \lambda} (\text{number of SYT of shape } \lambda \text{ with } n \text{ at } c) = \sum_{c \text{ is a corner of } \lambda} f(\lambda - c)$$

as required. □

To prove the theorem, we will use induction on the size of the Young diagram: the induction step will be to show that the numbers  $n!/H(\lambda)$  satisfy the same recursion as in Lemma 2.7. The base is easy: for a one-box Young diagram, all numbers in the equation are equal to 1. Thus, we need to prove the following.

**Lemma 2.8.** For every Young diagram  $\lambda \vdash n$ , we have

$$\sum_{c \text{ is a corner of } \lambda} \frac{(n-1)!}{H(\lambda - c)} = \frac{n!}{H(\lambda)}$$

The Theorem 2.5 then follows: once we prove Lemma 2.8, we have

$$f_\lambda = \sum_{c \text{ is a corner of } \lambda} f(\lambda - c) = \sum_{c \text{ is a corner of } \lambda} \frac{(n-1)!}{H(\lambda - c)} = \frac{n!}{H(\lambda)},$$

where the first equality follows from Lemma 2.7, the second one is the induction assumption (applied to every corner of  $\lambda$ ), and the last one is Lemma 2.8.

Observe that the equality in Lemma 2.8 can be reformulated as

$$\sum_{c \text{ is a corner of } \lambda} \frac{1}{n} \frac{H(\lambda)}{H(\lambda - c)} = 1$$

Therefore, we can try to interpret the summands in the left hand side as probabilities of some events. In other words, we will proceed according to the following plan: given  $\lambda \vdash n$ , find a random process, such that the space of outcomes is the set of corners of  $\lambda$ , and the probability of the outcome  $c$  is precisely  $\frac{1}{n} \frac{H(\lambda)}{H(\lambda - c)}$ . Then the sum of probabilities is automatically equal to 1, and thus we get the statement of the lemma.

The random process we will use is called a *hook walk* and is defined as follows.

- (1) Choose randomly a box  $b_1$  in  $\lambda$ , the probability of this is  $p(b_1) = 1/n$ .
- (2) Choose randomly a box  $b_2 \neq b_1$  in the hook of  $b_1$ , the probability of this is  $1/(h(b_1) - 1)$ .
- (3) Repeat Step (2) until reach a corner.



Denote by  $P(b, c)$  the probability of the walk starting at a box  $b$  ends in a corner  $c$ . The probability of an individual hook walk  $b = b_1 \rightarrow b_2 \rightarrow \dots \rightarrow b_k = c$  is equal to

$$\frac{1}{h(b_1) - 1} \cdots \frac{1}{h(b_{k-1}) - 1},$$

which implies that

$$P(b, c) = \sum_{\text{all walks } b=b_1 \rightarrow b_2 \rightarrow \dots \rightarrow b_k=c} \frac{1}{h(b_1) - 1} \cdots \frac{1}{h(b_{k-1}) - 1}$$

Define

$$P(c) = \frac{1}{n} \sum_{b \in \lambda} P(b, c),$$

then  $\sum_{c \text{ is a corner of } \lambda} P(c) = 1$ . Therefore, we need to prove that

$$P(c) = \frac{1}{n} \frac{H(\lambda)}{H(\lambda - c)},$$

or, equivalently,

$$\sum_{b \in \lambda} P(b, c) = \frac{H(\lambda)}{H(\lambda - c)}$$

**Example 2.9.** Let  $\lambda = (3, 2, 1) \vdash 6$ . There are four hook paths ending in the corner located in the first row (denote it by  $c_1$ ), same for the corner  $c_3$  in the third row, and five hook paths ending in the remaining corner  $c_2$ . Then  $P(c_1) = P(c_3) = 5/16$ , and  $P(c_2) = 3/8$ .

Let us make some observations.

- Let  $c = (u, v)$  be a corner. Then any hook walk ending in  $c$  is contained in the rectangle with vertices at  $(1, 1), (1, v), (u, 1)$  and  $(u, v)$ .
- If  $c = (u, v)$  is a corner, and  $p \leq u, q \leq v$ , then  $h(p, q) + h(u, v) = h(p, v) + h(u, q)$ . Since  $h(u, v) = 1$ , this implies the equality

$$h(p, q) - 1 = (h(p, v) - 1) + (h(u, q) - 1)$$

Define a *weight*  $\text{wt}(p, q)$  of a box  $(p, q)$  by

$$\text{wt}(p, q) = \frac{1}{h(p, q) - 1}$$

Then, if we denote  $1/x = \text{wt}(u, q)$  and  $1/y = \text{wt}(p, v)$ , the equality above says that  $\text{wt}(p, q) = 1/(x + y)$ .

- For any path, define its weight as the product of weights of all its boxes. Then we can write  $P(b, c)$  as the sum of weights of all hook walks from  $b$  to  $c$ .

To complete the proof, we need further two technical lemmas.

**Lemma 2.10.** Consider all lattice paths in a rectangle  $(l + 1) \times (k + 1)$  from the top left box to the bottom right box. Denote

$$\frac{1}{x_j} = \text{wt}(l + 1, j), \quad \frac{1}{y_i} = \text{wt}(i, k + 1), \quad x_{k+1} = y_{l+1} = 1.$$

Then

$$\sum_{\gamma \text{ is a lattice path}} \text{wt}(\gamma) = \frac{1}{x_1 \cdots x_k y_1 \cdots y_l}$$

$\frac{1}{x_1 + y_1}$	$\frac{1}{x_2 + y_1}$	$\frac{1}{y_1}$	$l = 1$
$\frac{1}{x_1}$	$\frac{1}{x_2}$	$1$	
$k = 2$			

Figure 2.3: Weights of boxes in a  $2 \times 3$  rectangle

**Example.** Let  $l = 1, k = 2$ , so we consider paths in a  $2 \times 3$  rectangle, see Fig. 2.3. There are  $\binom{k+l}{k} = \binom{3}{2} = 3$  lattice paths from top left box to the bottom right one. Then we can compute the sum of weights of the three paths:

$$\sum_{\gamma \text{ is a lattice path}} \text{wt}(\gamma) = \frac{1}{x_1 + y_1} \cdot \frac{1}{x_2 + y_2} \cdot \frac{1}{y_1} \cdot 1 + \frac{1}{x_1 + y_1} \cdot \frac{1}{x_2 + y_2} \cdot \frac{1}{x_2} \cdot 1 + \frac{1}{x_1 + y_1} \cdot \frac{1}{x_1} \cdot \frac{1}{x_2} \cdot 1 = \frac{1}{x_1 x_2 y_1}$$

*Proof of Lemma 2.10.* We use induction on  $k + l$ . If  $k + l = 0$ , then we have a  $1 \times 1$  square and both sides are equal to one. To prove the induction step, observe that every path starts either from a horizontal step or from a vertical one, i.e. the second box  $b_2$  in the path has coordinates  $(1, 2)$  or  $(2, 1)$  respectively. In both cases the remaining part of the path lies in a smaller rectangle, so we may use the induction assumption. More precisely, we have

$$\begin{aligned} \sum_{\gamma \text{ is a lattice path}} \text{wt}(\gamma) &= \sum_{\gamma | b_2=(1,2)} \text{wt}(\gamma) + \sum_{\gamma | b_2=(2,1)} \text{wt}(\gamma) = \\ &= \frac{1}{x_1 + y_1} \frac{1}{x_1 \dots x_k y_2 \dots y_l} + \frac{1}{x_1 + y_1} \frac{1}{x_2 \dots x_k y_1 \dots y_l} = \frac{1}{x_1 + y_1} \frac{1}{x_2 \dots x_k y_2 \dots y_l} \frac{x_1 + y_1}{x_1 y_1} = \\ &= \frac{1}{x_1 \dots x_k y_1 \dots y_l} \end{aligned}$$

□

In the next lemma we will consider hook walks, i.e. we are allowed to miss steps in a lattice path.

**Lemma 2.11.** Consider all hook walks in a rectangle  $(l + 1) \times (k + 1)$  ending in the bottom right box. Define  $x_j$  and  $y_i$  as in Lemma 2.10. Then

$$\sum_{\gamma \text{ is a hook walk}} \text{wt}(\gamma) = \left(1 + \frac{1}{x_1}\right) \dots \left(1 + \frac{1}{x_k}\right) \left(1 + \frac{1}{y_1}\right) \dots \left(1 + \frac{1}{y_l}\right)$$

*Proof.* The proof follows from an easy observation: all hook walks are precisely lattice paths in some subrectangles (i.e., a rectangle constructed from boxes lying in the intersection of some of rows and some of columns of the initial rectangle). Now, choosing 1 or  $1/x_j$  in the product is equivalent to the column  $j$  to be either absent or present in the subrectangle, respectively. The same holds for choosing 1 or  $1/y_i$  and presence of column  $i$ . Then every such term has the form as in Lemma 2.10.

□

We can now complete the proof of Lemma 2.8. Recall that the corner  $c$  has coordinates  $(u, v)$ . By Lemma 2.11, we have

$$\begin{aligned}
 P(c) &= \frac{1}{n} \sum_{b \in \lambda} P(b, c) = \frac{1}{n} \sum_{\gamma \text{ is a hook walk}} \text{wt}(\gamma) = \\
 &= \frac{1}{n} \left(1 + \frac{1}{x_1}\right) \dots \left(1 + \frac{1}{x_k}\right) \left(1 + \frac{1}{y_1}\right) \dots \left(1 + \frac{1}{y_l}\right) = \frac{1}{n} \prod_{\substack{b = (u, j) \mid j < v \\ b = (i, v) \mid i < u}} \left(1 + \frac{1}{h(b) - 1}\right) = \\
 &= \frac{1}{n} \prod_{\substack{b = (u, j) \mid j < v \\ b = (i, v) \mid i < u}} \left(\frac{h(b)}{h(b) - 1}\right) = \frac{1}{n} \frac{H(\lambda)}{H(\lambda - c)},
 \end{aligned}$$

where the last equality holds since the only boxes for which the hook lengths differ in  $\lambda$  and  $\lambda - c$  are precisely those located at row  $u$  and column  $v$ , and their hook lengths in  $\lambda - c$  are one less than in  $\lambda$ .

### 2.3 Set partitions

In the previous sections we considered “unlabeled” partitions. We will now consider the “labeled” version.

Denote by  $[n]$  the set of integers  $1, \dots, n$ .

**Definition 2.12.** A *set partition* of  $[n]$  is a subdivision of  $[n]$  into a disjoint union of non-empty subsets (*blocks*).

**Example 2.13.** Let  $n = 7$ , we can write  $[7] = \{1, 7\} \cup \{2, 3, 5\} \cup \{4, 6\}$  – this is one of partitions of shape  $(3, 2, 2) \vdash 7$ . Notation: if we call this partition  $\pi$ , we write  $\pi = (17 \mid 235 \mid 46)$  (note that the order of blocks does not matter).

Graphically, we can draw set partitions as arc diagrams: vertices correspond to set elements  $1, \dots, n$ , consecutive elements in one block are joined by an arc, see Fig. 2.4, left.




Figure 2.4: Partition  $(17 \mid 235 \mid 46)$  of  $[7]$ : arc diagram and rook placement

The arc diagrams representing set partitions are characterized as follows: every vertex is incident to at most one arc from the left and at most one arc from the right.

**Example 2.14.** Another graphical interpretation of set partitions is a *non-attacking rook placement*. If we have a partition of  $[n]$ , consider a chess “half-board” obtained from the square  $n \times n$  by taking all boxes  $(i, j)$  with  $i < j$ . Then place a rook in the box  $(i, j)$  if the corresponding arc diagram contains an arc connecting  $i$  and  $j$ , see Fig. 2.4, right.

**Exercise.** Show that rooks placed as described above do not attack each other. Show that this map has an inverse: any non-attacking rook placement gives rise to a set partition.

Note that the number of rooks (which is also the number of arcs) is equal to  $n - k$ , where  $k$  is the number of blocks.

**Definition 2.15.** The number of set partitions of  $[n]$  is called *Bell number*  $B(n)$ . The number of set partitions of  $[n]$  into  $k$  blocks is called *Stirling number of second kind*  $S(n, k)$ , i.e.  $B(n) = \sum_{k=1}^n S(n, k)$ .

**Example 2.16.** There are five set partitions of  $[3]$ , namely

$$(1 \mid 2 \mid 3), \quad (12 \mid 3), \quad (1 \mid 23), \quad (13 \mid 2), \quad (123),$$

so  $B(3) = 5$ ,  $S(3, 1) = S(3, 3) = 1$ ,  $S(3, 2) = 3$ .

**Definition 2.17.** A set partition is *non-crossing* if the arc diagram is non-crossing. A set partition is *non-nesting* if the arc diagram is non-nesting.

**Example 2.18.** For  $n = 1, 2, 3$  all set partitions are non-crossing and non-nesting. For  $n = 4$ , there is precisely one set partition which is not non-crossing and one which is not non-nesting.

**Exercise.** Compute  $B(4)$  by listing all arc diagrams.

Note that arc diagrams of non-crossing (non-nesting) partitions are not the same as of non-crossing (non-nesting) matchings. However, the following result holds.

**Theorem 2.19.** *The number of non-crossing set partitions of  $[n]$  is equal to the number of non-nesting set partitions of  $[n]$  and is equal to the Catalan number  $C_n$ .*

**Definition 2.20.** Given a Dyck path, a *peak* is a local maximum (i.e., a step up followed by a step down), and a *valley* is a local minimum (i.e., a step down followed by a step up). Clearly, the number of peaks exceeds the number of valleys by one.

*Proof of Theorem 2.19.* First we construct a bijection between non-nesting set partitions of  $[n]$  and Dyck paths of length  $2n$ . A partition is not non-nesting if and only if there are two arcs  $(a_i, b_i)$  such that  $a_i < a_j < b_j < b_i$ . This is equivalent to the rook  $(a_j, b_j)$  in the corresponding rook placement being located in the positive octant with respect to the origin at  $(a_i, b_i)$  (here by “positive” we mean standard coordinates in  $\mathbb{R}^2$ ). Therefore, the equivalent criterion for a set partition being non-nesting is the following: the positive octant with respect to every rook does not contain any other rook.

Now we can construct the map: take a set partition, consider the corresponding rook placement, draw the union of all positive octants centered at rooks, and then the boundary of this domain will be a Dyck path. The inverse map is constructed as follows: put rooks in all valleys of a Dyck path.

Given a Dyck path, a non-crossing partition can be constructed via a “shelling algorithm”, see Fig. 2.5 for an example.

□

**Exercise 2.21.** Fill in the details of the proof.

**Definition 2.22.** *Narayana number*  $N(n, k)$  is the number of Dyck paths of length  $2n$  with  $k$  peaks.

**Corollary 2.23** (of the proof of Thm. 2.19). *The number of non-nesting partitions of  $[n]$  with  $k$  blocks is equal to  $N(n, n - (k - 1))$ .*

**Example 2.24.** Let us compute  $N(4, 2)$ , i.e. the number of Dyck paths of length 8 with 2 peaks. By Cor. 2.23,  $N(4, 2)$  is equal to the number of non-nesting partitions of  $[4]$  with 3 blocks, which, in its turn, is equal to the number of rook placements, where the number of rooks is equal to  $4 - 3 = 1$ . The number of boxes in the upper half of the  $4 \times 4$  board is equal to 6, so there are 6 ways to place one rook, and thus  $N(4, 2) = 6$ .



Figure 2.5: (a) non-nesting partition  $(1\ 3\ 5\ 6 \mid 2\ 4 \mid 7)$  of  $[7]$ : rook placement and Dyck path; (b) non-crossing partition  $(1\ 7 \mid 2\ 3 \mid 4\ 6 \mid 5)$  of  $[7]$  and its Dyck path

## 2.4 Generating functions

Recall: if we have a sequence  $(a_n)$ , we can define a generating function of  $(a_n)$ , which is a formal power series  $a_0 + a_1x + a_2x^2 + \dots$ . This is an *ordinary* generating function. We can also define an *exponential* generating function.

**Definition 2.25.** *Exponential generating function* of a sequence  $(a_n)_{n \geq 0}$  is a formal power series  $\sum_{n=0}^{\infty} a_n \frac{x^n}{n!}$ .

Usually, ordinary generating functions are used when the objects are unlabeled, and exponential generating functions are used when the objects are labeled.

**Example 2.26.** The number  $p_n$  counts *integer partitions* of  $n$ , i.e. the number of Young diagrams  $\lambda \vdash n$ . Bell number  $B(n)$  counts partitions of  $[n]$ . Bell numbers are “labeled versions” of  $p(n)$ , so it is reasonable to look for an ordinary generating function for  $(p_n)$  and an exponential generating function for  $(B_n)$ .

Our next goal is to write the generating functions for  $p(n)$  and  $B(n)$ .

### 2.4.1 Integer partitions

Let  $\lambda = (\lambda_1, \dots, \lambda_k) \vdash n$  be a partition. We can write it as  $\lambda = 1^{m_1} 2^{m_2} 3^{m_3} \dots$ , where  $m_i$  is the number of  $\lambda_j = i$ .

**Example.**  $\lambda = (4, 2, 1, 1, 1) \vdash 9$  can be written as  $\lambda = 1^3 2^1 3^0 4^1$  (formally, there are infinitely many zero powers at the end, but we omit all factors  $r^0$  for  $r > \lambda_1$ ).

The notation above provides a bijection between all partitions of natural numbers and all sequences  $(m_i)_{i > 0}$  of non-negative integers with finite number of positive entries. Given a sequence  $(m_i)$ , it corresponds to a partition of  $n = 1 \cdot m_1 + 2 \cdot m_2 + 3 \cdot m_3 + \dots$ . If  $\lambda \vdash n$ , denote  $|\lambda| = n$ .

Then the (ordinary) generating function of  $p(n)$  is

$$\begin{aligned} \sum_{n \geq 0} p(n)x^n &= \sum_{n \geq 0} \underbrace{x^n + x^n + \dots + x^n}_{p(n) \text{ summands}} = \sum_{n \geq 0, \lambda \vdash n} x^{|\lambda|} = \sum_{\lambda \text{ any partition}} x^{|\lambda|} = \\ &= \sum_{m_1, m_2, \dots \geq 0} x^{1 \cdot m_1 + 2 \cdot m_2 + 3 \cdot m_3 + \dots} = \left( \sum_{m_1 \geq 0} x^{m_1} \right) \left( \sum_{m_2 \geq 0} x^{2m_2} \right) \left( \sum_{m_3 \geq 0} x^{3m_3} \right) \dots = \\ &= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1-x^3} \dots = \prod_{k \geq 1} \frac{1}{1-x^k} \end{aligned}$$

Note that although the product is infinite, the number of terms contributing to any coefficient is finite.

### 2.4.2 The exponential formula

Let numbers  $c_n$  count some objects on  $n$  labeled nodes (call them  $c$ -objects), where  $n > 0$ . Let  $c(x) = \sum_{n \geq 1} c_n \frac{x^n}{n!}$  be the exponential generating function of  $(c_n)$ .

Define  $d$ -objects on  $n$  nodes as collections of  $c$ -objects on  $n_1, \dots, n_k$  nodes such that  $n_1 + \dots + n_k = n$ . Let  $d(x) = \sum_{n \geq 0} d_n \frac{x^n}{n!}$  be the exponential generating function of  $(d_n)$ .

**Example 2.27.** Let  $c$ -objects be linear graphs on  $n$  nodes (or, equivalently, sequences of  $n$  distinct numbers  $1, \dots, n$ ). Then  $c_n = n!$ , and thus  $c(x) = x + x^2 + \dots = \frac{x}{1-x}$ .

Then  $d$ -objects are graphs on  $n$  nodes, such that any connected component is a linear graph (or, equivalently, unions of sequences of total length  $n$ ). How to compute  $d(x)$ ?

**Lemma 2.28** (Exponential formula).

$$d(x) = e^{c(x)}$$

*Proof.* A  $d$ -object on  $n$  nodes consists of a set partition of  $[n]$ ,  $n = n_1 + \dots + n_k$ , and then a  $c$ -object on every block. Let us fix  $k$  and  $n_1, \dots, n_k$ , and compute  $d$ -objects.

The number of ways to subdivide  $n$  into  $k$  blocks of size  $n_1, \dots, n_k$  is equal to the multinomial coefficient

$$\binom{n}{n_1 \ n_2 \ \dots \ n_k} = \frac{n!}{n_1! n_2! \ \dots \ n_k!}$$

We have a  $c$ -object on every block, so the number of  $d$ -object with fixed  $k$  and  $n_1, \dots, n_k$  is equal to

$$\frac{1}{k!} \binom{n}{n_1 \ n_2 \ \dots \ n_k} c_{n_1} \dots c_{n_k},$$

where we divide by  $k$  as we are not interested in the order of the blocks.

Now take a sum over  $k \geq 0$  and  $n_1 + \dots + n_k = n$ :

$$d_n = \sum_{k \geq 0} \sum_{\substack{n_1 + \dots + n_k = n \\ n_1, \dots, n_k \geq 1}} \frac{1}{k!} \frac{n!}{n_1! n_2! \ \dots \ n_k!} c_{n_1} \dots c_{n_k},$$

which implies that

$$\frac{d_n}{n!} = \sum_{k \geq 0} \sum_{\substack{n_1 + \dots + n_k = n \\ n_1, \dots, n_k \geq 1}} \frac{1}{k!} \frac{c_{n_1}}{n_1!} \frac{c_{n_2}}{n_2!} \dots \frac{c_{n_k}}{n_k!},$$

and thus

$$\begin{aligned} d(x) &= \sum_{n \geq 0} d_n \frac{x^n}{n!} = \sum_{n \geq 0} \sum_{k \geq 0} \sum_{\substack{n_1 + \dots + n_k = n \\ n_1, \dots, n_k \geq 1}} \frac{1}{k!} \frac{c_{n_1}}{n_1!} \frac{c_{n_2}}{n_2!} \dots \frac{c_{n_k}}{n_k!} x^n \\ &= \sum_{k \geq 0} \frac{1}{k!} \sum_{n_1, \dots, n_k \geq 1} \frac{c_{n_1} x^{n_1}}{n_1!} \frac{c_{n_2} x^{n_2}}{n_2!} \dots \frac{c_{n_k} x^{n_k}}{n_k!} = \sum_{k \geq 0} \frac{1}{k!} \left( \sum_{m \geq 1} \frac{c_m x^m}{m!} \right)^k = \sum_{k \geq 0} \frac{1}{k!} c(x)^k = e^{c(x)}. \end{aligned}$$

□

**Corollary 2.29.** *The exponential generating function for Bell numbers is*

$$B(x) = e^{e^x - 1}$$

*Proof.* Let  $c$ -objects be just sets (so,  $c_n = 1$ ), and  $d$ -objects be collections of sets of total cardinality  $n$ . Then  $d$ -objects are precisely partitions of  $[n]$ . Since  $c_n = 1$ , we have  $c(x) = e^x - 1$ , and thus  $B(x) = d(x) = e^{e^x - 1}$  by the exponential formula. □

## 3 Permutations

### 3.1 Definitions and notation

Recall that a *permutation* is a bijection  $w : [n] \rightarrow [n]$ . Permutations form a group (called *symmetric group*)  $S_n$  with respect to composition, the order of the group is  $n!$ .

There are several notations for  $w \in S_n$ .

- *2-line notation:* if  $w(i) = w_i$ , we write

$$w = \begin{pmatrix} 1 & 2 & \dots & n \\ w_1 & w_2 & \dots & w_n \end{pmatrix}, \quad \text{e.g.} \quad w = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 3 & 2 & 6 & 1 & 5 & 7 & 4 \end{pmatrix} \in S_7;$$

- *1-line notation:* we simply write  $w = w_1, w_2, \dots, w_n$ , or even  $w = w_1 w_2 \dots w_n$ , e.g.  $w = 3261574$ ;
- *cycle notation:* recall that every permutation can be decomposed a a product of disjoint cycles. For the permutation above, we have  $w = (1\ 3\ 6\ 7\ 4)(2)(5)$ . Note that the order of cycles is irrelevant, as well as the starting point of each cycle: we could also write  $w = (2)(6\ 7\ 4\ 1\ 3)(5)$ .

### 3.2 Statistics on permutations

**Definition 3.1.** A *statistics* on permutations is a function  $\sigma : S_n \rightarrow \mathbb{Z}_{\geq 0}$ . Its *generating function* is a polynomial  $f_\sigma(x) = \sum_{w \in S_n} x^{\sigma(w)}$ .

Two statistics  $\sigma$  and  $\mu$  are *equidistributed* if  $f_\sigma(x) = f_\mu(x)$ , i.e. the number of permutations with the same value of statistics is the same.

**Definition 3.2.** Let  $w \in S_n$ .

- An *inversion* is a pair  $(i, j)$  such that  $i < j$ ,  $w_i > w_j$ ;
- A *descent* is  $i \in [n - 1]$  such that  $w_{i+1} < w_i$ .

We can now look at some statistics on  $S_n$ .

- $\text{inv}(w)$  is the number of inversions in  $w$ ;
- $\text{des}(w)$  is the number of descents in  $w$ ;
- $\text{cyc}(w)$  is the number of cycles in  $w$ .

**Example 3.3.** Let  $w = 3261574 \in S_7$ . Then  $\text{inv}(w) = 8$ ,  $\text{des}(w) = 3$ ,  $\text{cyc}(w) = 3$ .

**Definition 3.4.** A statistics on  $S_n$  is called *Mahonian* if it is equidistributed with  $\text{inv}$ , and *Eulerian* if it is equidistributed with  $\text{des}$ .

**Example 3.5.** Let  $n = 3$ , order permutations by the number of inversions. Then we have the following.

$w$	1 2 3	2 1 3	1 3 2	2 3 1	3 1 2	3 2 1
$\text{inv}(w)$	0	1	1	2	2	3
$\text{des}(w)$	0	1	1	1	1	2
$\text{cyc}(w)$	3	2	2	1	1	2

In particular, we can see that all three statistics are different.

We can also compute the generating functions:

$$\begin{aligned} f_{\text{inv}}(x) &= 1 \cdot x^0 + 2 \cdot x^1 + 2 \cdot x^2 + 1 \cdot x^3 = (1+x)(1+x+x^2) \\ f_{\text{des}}(x) &= 1 \cdot x^0 + 4 \cdot x^1 + 1 \cdot x^2 = 1 + 4x + x^2 \\ f_{\text{cyc}}(x) &= 0 \cdot x^0 + 2 \cdot x^1 + 3 \cdot x^2 + 1 \cdot x^3 = x(x+1)(x+2) \end{aligned}$$

**Theorem 3.6.** For every  $n \in \mathbb{N}$ ,

$$f_{\text{inv}}(x) = (1+x)(1+x+x^2) \dots (1+x+\dots+x^{n-1}) = \frac{\prod_{k=1}^n (1-x^k)}{(1-x)^n}$$

*Proof.* We use induction on  $n$ . For  $n = 1$  the result is clear (as for  $n = 2, 3$ ). Assume that the theorem holds for  $n - 1$ . Take any permutation  $w' \in S_{n-1}$  and “insert”  $n$  in all possible  $n$  places. Depending on the place where  $n$  is inserted, we add from 0 to  $n - 1$  inversions, where all numbers show up. Therefore, the generating function is

$$\begin{aligned} f_{\text{inv}}(x) &= \sum_{w \in S_n} x^{\text{inv}(w)} = \sum_{w' \in S_{n-1}} x^{\text{inv}(w') + x} \sum_{w' \in S_{n-1}} x^{\text{inv}(w') + x^2} \sum_{w' \in S_{n-1}} x^{\text{inv}(w') + \dots + x^{n-1}} \sum_{w' \in S_{n-1}} x^{\text{inv}(w')} = \\ &= \left( \sum_{w' \in S_{n-1}} x^{\text{inv}(w')} \right) (1+x+\dots+x^{n-1}) = (1+x)(1+x+x^2) \dots (1+x+\dots+x^{n-1}), \end{aligned}$$

where the last equality follows from the induction assumption. □

**Definition 3.7.** The *Major index* of  $w \in S_n$  is  $\text{maj}(w) = \sum_{k \text{ is a descent of } w} k$ .

**Example 3.8.** For  $w = 3261574 \in S_7$ , we have  $\text{maj}(w) = 1 + 3 + 6 = 10$ .

For  $n = 3$ , we have

$w$	1 2 3	2 1 3	1 3 2	2 3 1	3 1 2	3 2 1
$\text{maj}(w)$	0	1	2	2	1	3

and thus  $f_{\text{maj}}(x) = 1 + 2x + 2x^2 + x^3 = (1+x)(1+x+x^2) = f_{\text{inv}}(x)$ .

**Exercise.** Show that  $f_{\text{maj}}(x) = f_{\text{inv}}(x)$  for every  $n \in \mathbb{N}$ , i.e.  $\text{maj}$  is Mahonian.

**Definition 3.9.** Let  $w = w_1 w_2 \dots w_n \in S_n$ ,  $w_i$  is a *record* of  $w$  if  $w_i > w_j$  for all  $j < i$ . Denote by  $\text{rec}(w)$  the number of records of  $w$ .



**Example.** for  $w = 3261574 \in S_7$ , we have  $\text{rec}(w) = 3$ .

For  $n = 3$ , we have

$w$	1 2 3	2 1 3	1 3 2	2 3 1	3 1 2	3 2 1
$\text{rec}(w)$	3	2	2	2	1	1

and thus  $f_{\text{rec}}(x) = 2x + 3x^2 + x^3 = f_{\text{cyc}}(x)$ .

**Theorem 3.10.**  $f_{\text{rec}}(x) = f_{\text{cyc}}(x)$  for every  $n \in \mathbb{N}$ .

*Proof.* We construct a bijection  $f : S_n \rightarrow S_n$ , such that if  $w' = f(w)$  then  $\text{rec}(w') = \text{cyc}(w)$ .

First, write the cycle decomposition in a standard way: if  $w = (a_1 \dots)(a_2 \dots) \dots (a_k \dots)$  then every  $a_i$  is maximal in its cycle, and  $a_1 < a_2 < \dots < a_k$ .

Now, the map  $f$  takes  $w$  written in the standard way above to  $w'$  by erasing all brackets (and considering the result as a 1-line notation for  $w'$ ). Clearly, records are first elements of cycles, so  $\text{rec}(w') = \text{cyc}(w)$ .

The map  $f$  has an inverse: take any permutation  $w'$  in 1-line notation, put brackets in the beginning and at the end, and then put closing and opening brackets before every record. We get a cycle decomposition of some permutation  $w$ . Then  $f(w) = w'$ . □

**Example 3.11.** The preimage of  $w = 3261574 \in S_7$  under  $f$  is  $u = (326)(15)(74)$ .

Now,  $w = 3261574 = (13674)(2)(5) = (2)(5)(74136)$ , so  $f(w) = 2574136$ .

**Definition 3.12.** Let  $w = w_1 w_2 \dots w_n \in S_n$ ,  $i \in [n]$  is an *excedance* of  $w$  if  $i < w_i$ . Denote by  $\text{exc}(w)$  the number of excedances of  $w$ .

**Example.** For  $w = 3261574 \in S_7$ , we have  $\text{exc}(w) = 3$ .

For  $n = 3$ , we have

$w$	1 2 3	2 1 3	1 3 2	2 3 1	3 1 2	3 2 1
$\text{exc}(w)$	0	1	1	2	1	1

and thus  $f_{\text{rec}}(x) = 1 + 4x + x^2 = f_{\text{des}}(x)$ .

**Theorem 3.13.**  $f_{\text{exc}}(x) = f_{\text{des}}(x)$  for every  $n \in \mathbb{N}$ , so  $\text{exc}$  is Eulerian.

*Proof.* Define *anti-excedance* of  $w$  as  $i \in [n]$  such that  $i > w_i$ . Observe, that anti-excedances of  $w$  are precisely excedances of  $w^{-1}$ , so anti-excedances and excedances are equidistributed.

Now, we claim that the map  $f$  from the proof of Thm 3.10 takes anti-excedances of  $w$  to descents of  $w' = f(w)$ . □

**Exercise.** Complete the proof of the theorem.

## 4 Posets and lattices

### 4.1 Definitions

**Definition 4.1.** A *partially ordered set*, or *poset*,  $P$  is a set with a binary relation  $\leq$  (or  $\leq_P$  if there is an ambiguity) satisfying the following axioms:

- $a \leq a$  (reflexivity);

- if  $a \leq b$  and  $b \leq c$  then  $a \leq c$  (transitivity);
- if  $a \leq b$  and  $b \leq a$  then  $a = b$  (symmetry).

We say  $a < b$  if  $a \leq b$  and  $a \neq b$ , and the notation  $a \geq b$  means that  $b \leq a$ .  
Two elements are *incomparable* if  $a \not\leq b$  and  $b \not\leq a$ .

**Example 4.2.** Any ordered field (e.g.  $\mathbb{R}$ ) is a poset, where one considers usual order relation. (Such posets are *totally ordered*, i.e. they do not contain incomparable elements).

The power set of any set (i.e., the set of all subsets), with the order being inclusion, is also a poset (see below). Two subsets are incomparable if their symmetric difference is not empty.

**Definition 4.3.** A *covering relation*  $a < \cdot b$  in a poset  $P$  is defined as follows:  $a < \cdot b$  if  $a < b$  and there is no element  $c \in P$  such that  $a < c < b$ .

**Definition 4.4.** A *Hasse diagram* of a poset  $P$  is an oriented graph, where vertices are elements of  $P$ , and there is an edge  $a \rightarrow b$  if  $a < \cdot b$ . Hasse diagram is usually drawn as an undirected graph with  $b$  positioned above  $a$  if  $a < \cdot b$ .

Note that any finite poset is uniquely defined by its Hasse diagram.

**Example 4.5.** The Hasse diagram shown in Fig. 4.1 defines a poset. We have  $x \leq y$  if and only if there is a path from  $x$  to  $y$  every step of which goes up. For example,  $a \leq c, d, e, f$  and is not comparable to  $b$ . Overall, there are five pairs of incomparable elements.




Figure 4.1: Hasse diagram of a poset.

**Definition 4.6.** A *chain* in a poset  $P$  is a sequence of elements  $a_1 < a_2 < \dots < a_k$ . A chain is *saturated* if  $a_i < \cdot a_{i+1}$  for every  $i = 1, \dots, k - 1$ . An *antichain* is a subset of  $P$  consisting of mutually incomparable elements.

**Example.** In the poset defined in Fig. 4.1,  $(c, f)$  is an antichain, and  $(a, d, f)$  is a chain as well as  $(b, e)$ .

**Definition 4.7.** A *Boolean lattice*  $B_n$  is the poset of subsets of  $[n]$  ordered by inclusion:  $A \leq B$  if  $A \subset B$ . The set of subsets (or the power set) of  $[n]$  is denoted by  $2^{[n]}$ .

**Example 4.8.** Hasse diagrams for  $B_2$  and  $B_3$  are shown in Fig. 4.2. Note that for every  $n$  the Hasse diagram for  $B_n$  is a 1-skeleton of a cube.

We are interested in the following questions: what are the maximal sizes of chains and antichains in posets? For example, in the poset defined in Fig. 4.1, these are 3 and 2 respectively. In  $B_n$ , the maximal size of chain is  $n + 1$ , but what is the maximal size of an antichain?



Figure 4.2: Hasse diagram of  $B_2$  and  $B_3$ .

## 4.2 Sperner's Theorem

In this section we will answer the question above by proving the following result.

**Theorem 4.9** (Sperner's Theorem). *Let  $S_1, \dots, S_N$  be different subsets of  $[n]$  such that  $S_i \not\subseteq S_j$  and  $S_j \not\subseteq S_i$  for any  $i \neq j$ . Then  $N \leq \binom{n}{\lfloor n/2 \rfloor}$ , where  $\lfloor x \rfloor$  is the maximal integer not exceeding  $x$ .*

**Remark.** Note that the bound in the theorem is sharp: the number of  $\lfloor n/2 \rfloor$ -subsets of  $[n]$  is precisely  $\binom{n}{\lfloor n/2 \rfloor}$ , and they are all incomparable.

**Definition 4.10.** A poset  $P$  is *ranked* if there is a function  $\rho : P \rightarrow \mathbb{Z}$  such that  $a < \cdot b$  implies  $\rho(b) = \rho(a) + 1$ . If  $P$  is finite, we assume that the minimal rank is zero. If  $P$  is finite and ranked, the *rank numbers*  $r_i$  are defined as the numbers of elements of  $p$  of rank  $i$ .

In other words, a poset is ranked if its Hasse diagram can be drawn "on levels".

**Example 4.11.** The left poset shown in Fig. 4.3 is ranked, and the right is not.



Figure 4.3: Ranked (left) and not ranked (right) posets.

A finite ranked poset  $P$  with non-zero rank numbers  $r_0, \dots, r_l$  is *rank symmetric* if  $r_i = r_{l-i}$ , *unimodal* if there is  $k$  such that  $r_0 \leq r_1 \leq r_k \geq r_{k+1} \geq \dots \geq r_l$ , and *Sperner* if the maximal size of antichain is the maximum of  $r_i$  (i.e., the maximal size of antichain is minimal possible).

**Example 4.12.** The Hasse diagram in Fig. 4.4 defines a non-Sperner poset: both rank numbers are four, but there is an antichain of size five.

**Remark.** The Boolean lattice  $B_n$  is ranked, where  $\rho(A) = |A|$ , with ranking numbers  $r_i = r_i(B_n) = \binom{n}{i}$ . Since  $\binom{n}{i} = \binom{n}{n-i}$ ,  $B_n$  is rank-symmetric. It is also unimodal as

$$\binom{n}{0} \leq \binom{n}{1} \leq \dots \leq \binom{n}{\lfloor n/2 \rfloor} \geq \dots \geq \binom{n}{n}$$




Figure 4.4: Non-Sperner rank-symmetric poset. Colored vertices form an antichain of size 5.

Then Theorem 4.9 can be reformulated as:  $B_n$  is Sperner.

**Definition 4.13.** Let  $P$  be a finite ranked poset,  $\rho$  is the ranking function with values  $0, \dots, l$ . A *symmetric chain decomposition* (or SCD for short) of  $P$  is a decomposition of  $P$  as a disjoint union  $P = C_1 \sqcup C_2 \sqcup \dots \sqcup C_k$ , such that each  $C_i$  is a saturated chain, and if  $C_i = a_0^i < a_1^i < \dots < a_s^i$  then  $\rho(a_0^i) + \rho(a_s^i) = l$ .

**Example.**  $B_2$  has a SCD:  $C_1 = \emptyset < \{1\} < \{1, 2\}$  and  $C_2 = \{1\}$ .

**Lemma 4.14.** *If a finite ranked poset  $P$  has a SCD, then it is rank-symmetric, unimodal and Sperner.*

*Proof.*  $P$  is clearly rank-symmetric and unimodal, where the maximal ranking number is  $r_i$  for  $i = \lfloor l/2 \rfloor$ , as this property holds for every chain in the SCD. We need to show that  $P$  is Sperner.

Take any antichain  $A$ , then  $|A \cap C_i| \leq 1$ , so  $|A| \leq k$ . Since every chain  $C_i$  intersects the middle level, the number of chains  $k = r_{\lfloor l/2 \rfloor}$ , so the size of any antichain does not exceed the maximal ranking number, which is precisely the definition of Sperner poset.  $\square$

Due to Lemma 4.14, to prove Thm 4.9 we are left to show that  $B_n$  has a SCD.

**Definition 4.15.** Let  $P_1, P_2$  be posets, then  $P_1 \times P_2$  is also a poset: we can define the order by  $(a, b) \leq (c, d)$  if  $a \leq c$  and  $b \leq d$ . Similarly, one can define a product  $P_1 \times \dots \times P_n$  of any number of posets.

Observe that  $B_n = [2]^n = [2] \times \dots \times [2]$ . Indeed, we can assign to every element  $A$  of  $B_n$  a sequence of 0 and 1 of length  $n$ , where  $a_i = 0$  if  $i \notin A$  and  $a_i = 1$  if  $i \in A$ . Then  $A \subset B$  is equivalent to  $a_i \leq b_i$  for all  $i$ , and thus to  $A \leq B$ .

We are now left to prove the following statement.

**Theorem 4.16.** *Any product of chains has a SCD.*

*Proof.* We proceed by induction. First, a product of two chains has a SCD (prove this!). Now take  $P = P' \times C = (C_1 \sqcup C_2 \sqcup \dots \sqcup C_k) \times C$ . As a product of two chains has a SCD, every  $C_i \times C$  has a SCD. Note that the “middle rank” of all  $C_i \times C$  is the same – this is guaranteed by the fact  $P' = C_1 \sqcup C_2 \sqcup \dots \sqcup C_k$  is a SCD. Thus, taking the union of all SCD for  $C_i \times C$ ,  $i \in [k]$ , we obtain a SCD for  $P$ .  $\square$

### 4.3 Greene’s Theorem

Let  $P$  be a finite poset.

**Theorem 4.17** (Dilworth). *The maximal size of an antichain of  $P$  is equal to the minimal number of chains needed to cover  $P$ .*

**Remark.** Note that one inequality in the theorem is evident: an antichain cannot contain more elements than the number of chains as it intersect every chain at most once.

A dual version also holds.

**Theorem 4.18** (Minsky). *The maximal size of a chain of  $P$  is equal to the minimal number of antichains needed to cover  $P$ .*

More generally, define  $l_k$  to be the maximal size of a union of  $k$  chains of  $P$ , and  $m_k$  to be the maximal size of a union of  $k$  antichains of  $P$ . In particular,  $l_0 = m_0 = 0$ , and  $l_1$  is the maximal size of a chain in  $P$ .

**Theorem 4.19** (Greene). *Denote  $\lambda_i = l_i - l_{i-1}$  for  $i > 0$ , and let  $\lambda(p) = \lambda_1, \lambda_2, \dots$  (note that only finitely many  $\lambda_i$  are positive). Similarly, denote  $\mu_i = m_i - m_{i-1}$  for  $i > 0$ , and let  $\mu(p) = \mu_1, \mu_2, \dots$ . Then  $\lambda_i \geq \lambda_{i+1}$  and  $\mu_i \geq \mu_{i+1}$  for all  $i > 0$  (so  $\lambda$  and  $\mu$  can be considered as Young diagrams or partitions of  $|P|$ ), and Young diagrams  $\lambda$  and  $\mu$  are conjugate to each other, i.e. they are symmetric with respect to the main diagonal.*

**Example.** Let  $P$  be defined by the Hasse diagram shown in Fig. 4.5. Then  $l_1 = 3$ ,  $l_2 = 5$ , and  $l_i = 5$  for  $i \geq 2$ . Also,  $m_1 = 2$ ,  $m_2 = 4$ ,  $m_3 = 5$ , and  $m_i = 5$  for  $i \geq 3$ . Therefore,  $\lambda = (3, 2)$ , and  $\mu = (2, 2, 1)$  which are clearly conjugated.




Figure 4.5: Hasse diagram of a poset.

**Remark.** Theorems of Dilworth and Minsky are partial cases of Greene's Theorem. Indeed, the maximal size of antichain is  $m_1 = \mu_1$ , i.e. the length of the first row of  $\mu$ . The minimal number of chains needed to cover  $P$  is precisely the number of non-zero  $\lambda_i$ , i.e. the number of rows of  $\lambda$ . Greene's Theorem says that Young diagrams  $\lambda$  and  $\mu$  conjugate, which implies that the first row of  $\mu$  is equal to the first column of  $\lambda$ , and this is precisely Dilworth's Theorem. Minsky's Theorem follows similarly.

Given  $w \in S_n$ , define a poset  $P_w$  as follows: elements of  $P_w$  are elements of  $[n]$ , and  $w_i <_{P_w} w_j$  if  $w_i < w_j$  and  $i < j$ .

**Example.** Let  $w = 3261574 \in S_7$ . Hasse diagram of  $P_w$  is shown in Fig. 4.6.




Figure 4.6: Hasse diagram of the poset  $P_w$  for  $w = 3261574 \in S_7$ .

**Exercise.** Show that chains of  $P_w$  are precisely increasing subsequences of  $w_i$ , and antichains are decreasing subsequences of  $w_i$ .

Another corollary of Greene's Theorem is the following statement.

**Theorem 4.20** (Erdos–Szekeres). *Let  $m, n \geq 1$ . Then any permutation of size at least  $mn + 1$  contains either an increasing subsequence of length  $m + 1$  or a decreasing subsequence of length  $n + 1$ .*

*Proof.* According to Greene's Theorem, we can associate a Young diagram  $\lambda$  to  $P_w$ , in which the length of the first row is the maximal size of chain in  $P_w$ , i.e. the maximal length of an increasing subsequence in  $w$  (see the exercise above). Similarly, the length of the first row of the conjugate Young diagram (and thus the length of the first column of  $\lambda$ ) is the maximal size of antichain in  $P_w$ , i.e. the maximal length of a decreasing subsequence in  $w$ . The rest follows from the following elementary statement.

**Exercise.** Let  $\lambda$  be a Young diagram,  $|\lambda| \geq mn + 1$ . Then either  $\lambda_1 \geq n + 1$ , or  $\lambda'_1 \geq m + 1$ , where  $\lambda'_1$  is the length of the first column of  $\lambda$ .

□

**Exercise.** Prove Theorem 4.20 without using Greene's Theorem.

## 4.4 Lattices

**Definition 4.21.** Let  $P$  be a poset,  $x, y \in P$ .

·  $z \in P$  is a *join* of  $x$  and  $y$  (notation  $z = x \vee y$ ) if the following hold:

- (1)  $z \geq x, y$ ;
- (2) if  $v \in P$  and  $v \geq x, y$  then  $v \geq z$ .

Note that (2) implies that if join exists then it is unique.

·  $t \in P$  is a *meet* of  $x$  and  $y$  (notation  $t = x \wedge y$ ) if the following hold:

- (1)  $t \leq x, y$ ;
- (2) if  $v \in P$  and  $v \leq x, y$  then  $v \leq t$ .

Again, meet is unique if exists.

·  $P$  is a *lattice* if every two elements have a join and a meet.

**Example.** Out of the three posets whose Hasse diagrams shown in Fig. 4.7 only the left one is a lattice. In the other two posets elements  $x, y$  have no join.




Figure 4.7: The poset on the left is a lattice, the other two are not.

Note that if  $x \leq y$ , then  $x \vee y = y$  and  $x \wedge y = x$ .

**Exercise.** Every finite lattice has a unique *minimal* and a unique *maximal* elements (i.e., elements  $c$  and  $d$  such that  $c \leq x \leq d$  for all  $x$ ).

**Lemma 4.22.** *Boolean lattice  $B_n$  is indeed a lattice.*

*Proof.* Observe that if  $X, Y \in B_n$ , then  $X \vee Y = X \cup Y$  and  $X \wedge Y = X \cap Y$ . □

**Example 4.23.** *Young lattice  $\mathbb{Y}$  is the poset of all Young diagrams ordered by inclusion. The covering relation in  $\mathbb{Y}$  is defined as follows:  $\lambda < \cdot \mu$  if  $\mu$  has precisely one extra box.*

**Exercise.** Check that union and intersection of Young diagrams is again a Young diagram. Show that  $\mathbb{Y}$  is a lattice.

**Example.** *Partition lattice  $\Pi_n$  consists of all set partitions of  $[n]$  ordered by refinement:  $\lambda \leq \mu$  if  $\mu$  is obtained by joining together some blocks of  $\lambda$ . The covering relation is then following:  $\lambda < \cdot \mu$  if  $\mu$  is obtained by combining two blocks of  $\lambda$ .*

**Exercise.** Show that  $\Pi_n$  is a lattice.

**Definition 4.24.** Let  $P$  be a poset. An *order ideal* of  $P$  is a set  $I \subset P$  such that if  $x \in I$  and  $y \leq x$  then  $y \in I$ . Define an *order* on order ideals:  $I \leq J$  if  $I \subset J$ . This define a *poset of order ideals*  $J(P)$ .

**Example.** A poset and its poset of order ideals are shown in Fig. 4.8.




Figure 4.8: Hasse diagrams of a poset (left) and of its poset of order ideals (right).

**Lemma 4.25.**  *$J(P)$  is a lattice.*

*Proof.* Since the order is defined as in the Boolean lattice, meet and join are the intersection and the union (which are also order ideals – check this!). □

**Example 4.26.** · Consider a poset on  $\mathbb{Z}_{\geq 0}$  with usual order. Then  $J(\mathbb{Z}_{\geq 0}) \cong \mathbb{Z}_{\geq 0} \cup \{\infty\}$  (an order ideal corresponds to its maximal element).

- $J(\mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 0}) \cong \mathbb{Y}$  (Young lattice)
- We can also interpret  $B_n$  as  $J([n])$ , where all elements of  $[n]$  are incomparable.

**Definition 4.27.** A Lattice is *distributive* if

$$x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z) \quad \text{and} \quad x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$$

**Remark.** An ordered field is not a distributive lattice: only one of the distributivity laws holds.

**Exercise.**  $J(P)$  is a distributive lattice.

**Theorem 4.28** (Fundamental Theorem of Finite Distributive Lattices, Birkhoff). *Any finite distributive lattice is a poset of order ideals for some finite poset.*

*Proof.* Let  $P$  be any poset. We call  $z \in P$  *join-irreducible* if  $z \neq x \vee y$  for any incomparable  $x, y \in P$ , and  $z$  is not a minimal element of  $P$ .

Now, let  $L$  be a finite distributive lattice. Define  $P$  to be the poset of all join-irreducible elements of  $L$  (with the order inherited from  $L$ ). We will prove that  $L = J(P)$ .

Take  $x \in L$ , and consider  $I_x = \{y \in P \mid y \leq x\} \subset P$ . Clearly,  $I_x \in J(P)$ : if  $y \in I_x$ ,  $z \in P$  and  $z \leq y$  then  $z \leq y \leq x$  and thus  $z \in I_x$ .

Thus, we constructed a map from  $L$  to  $J(P)$  taking  $x \in L$  to  $I_x \in J(P)$ . We need to show that this map is injective and surjective. Injectivity follows from the following exercise:

**Exercise.** For every  $x \in L$ ,  $x = \vee\{t \mid t \in I_x\}$ .

Indeed, the exercise implies that if  $I_x = I_y$  and  $x \neq y$ , then the same set have two different joins, which is impossible in a lattice. To surjectivity, we will prove the following claim.

**Claim.** Let  $I \in J(P)$ , take  $x = \vee\{t \mid t \in I\}$ . Then  $I = I_x$ .

The claim explicitly states that the map  $x \mapsto I_x$  is surjective: for every order ideal of  $P$  we find a preimage. Therefore, we are left to prove the claim. □

*Proof of the claim.* First,  $I \subset I_x$ : since  $x = \vee\{t \mid t \in I\}$ ,  $t \leq x$  for all  $t \in I$ , and thus every  $t \in I$  also lies in  $I_x$  by the definition of  $I_x$ . We are left to prove that  $I_x \subset I$ .

Since  $x = \vee\{y \mid y \in I_x\} = \vee\{t \mid t \in I\}$ , we have

$$x = \vee\{t \mid t \in I\} = \vee\{y \mid y \in I_x\}$$

Take any  $s \in I_x$ , we want to show that  $s \in I$ . Take the meet  $s \wedge x$ , expanding both sides of the equality above by distributivity. Then we get

$$\vee\{t \wedge s \mid t \in I\} = \vee\{y \wedge s \mid y \in I_x\}$$

In the RHS, every  $y \wedge s \leq s$ , and since  $s \in I_x$  there is also a term  $s \wedge s = s$ , so the RHS is equal to  $s$ , and thus the LHS is equal to  $s$ . Since  $s \in I_x$ ,  $s$  is join-irreducible, which means that at least one of the elements of the set  $\{t \wedge s \mid t \in I\}$  must be equal to  $s$ . But  $s \wedge t = s$  implies  $s \leq t$ , and since  $I$  is an order ideal,  $s \leq t \in I$  implies  $s \in I$ . □

**Example.** Let  $\lambda$  be a Young diagram. Define a lattice  $\mathbb{Y}_\lambda$  consisting of all Young diagrams that fit inside  $\lambda$  (with the order inherited from  $\mathbb{Y}$ ). The  $\mathbb{Y}_\lambda$  is a distributive lattice (check this!). Join-irreducible elements of  $\mathbb{Y}_\lambda$  are Young diagrams with precisely one corner, i.e. rectangles. Define  $P_\lambda$  as a poset on boxes of  $\lambda$ , with  $(i, j) \leq (k, l)$  if  $i \leq k$  and  $j \leq l$ . Then  $J(P_\lambda) = \mathbb{Y}_\lambda$  (check this!).



## 4.5 Linear extensions of posets

Let  $P$  be a finite poset,  $|P| = n$ .

**Definition 4.29.** A *linear extension* of a poset  $P$  is a bijective map  $f : P \rightarrow [n]$  such that  $x \leq_P y$  implies  $f(x) \leq f(y)$ . Denote by  $\text{ext}(P)$  the number of linear extensions of  $P$ .

**Example.** Let  $P = P_\lambda$ , where  $\lambda = (2, 2) \vdash 4$ . Then linear extensions can be identified with SYT of shape  $\lambda$ . In particular,  $\text{ext}(P) = 2$ .

**Lemma 4.30.** Let  $P$  be a finite poset, and let  $J(P)$  be its poset of order ideals. Then  $\text{ext}(P)$  is equal to the number of maximal saturated chains in  $J(P)$ .

**Corollary 4.31.** The number of SYT of shape  $\lambda$ ,  $f_\lambda$ , is equal to the number of saturated chains from  $\emptyset$  to  $\lambda$  in  $\mathbb{Y}_\lambda$ .

*Proof of Lemma 4.30.* We want to construct a bijection between extensions of  $P$  and maximal saturated chains in  $J(P)$ . Let  $\varphi : P \rightarrow [n]$  be an extension,  $j \in [n]$ . Define an order ideal  $I_j = \varphi^{-1}([j])$ , then we get a saturated chain  $\emptyset < I_1 < \dots < I_n = \lambda$ .

**Exercise.** Show that  $\varphi$  is a bijection.

□

## 5 Robinson-Schensted correspondence

### 5.1 Algorithm

Let  $\lambda = (\lambda_1, \dots, \lambda_l) \vdash n$  be an integer partition. Recall that  $f_\lambda$  is the number of standard Young tableaux (SYT) of shape  $\lambda$ , which is also equal to the number of saturated chains in the Young lattice  $\mathbb{Y}$  from  $\emptyset$  to  $\lambda$ .

**Theorem 5.1.**

$$\sum_{\lambda \vdash n} (f_\lambda)^2 = n!$$

**Example 5.2.** For  $n = 3$ , there are precisely three Young diagrams. Two of these correspond to a unique SYT, and the other has two SYT. Then  $1^2 + 2^2 + 1^2 = 6 = 3!$ .

**Remark** (can be ignored by those not taking Representation Theory). Theorem 5.1 can be interpreted as a fact from the representation theory of symmetric groups. Irreducible representations of  $S_n$  are parametrized by integer partitions of  $n$ , i.e. by Young diagrams. Then  $f_\lambda$  is precisely the dimension of the irreducible representation  $V_\lambda$ ,  $n!$  is the order of  $S_n$ , and thus the theorem says that sum of squares of dimensions of irreducible representations is equal to the order of the group (which is true for any finite group and can be proved by considering regular representation).

We will prove Thm 5.1 combinatorially by constructing a bijection between permutations in  $S_n$  and pairs  $(P, Q)$  of SYT of the same shape  $\lambda \vdash n$ .

The algorithm is due to Robinson–Schensted, with a generalization due to Knuth, and is usually referred as RSK (Robinson–Schensted–Knuth).

Given  $w = w_1 w_2 \dots w_n \in S_n$ , we will construct an *insertion tableau*  $P$  and a *recording tableau*  $Q$  of the same shape step by step.

**Example 5.3.** Let  $w = 3261574 \in S_7$ . In the beginning,  $P = \emptyset$  and  $Q = \emptyset$ . We start adding  $w_i$  in turn. First, let us consider how  $P$  is changing. Every new  $w_i$  is inserted in the first row.

- At the first step, 3 is inserted in the box (1,1).
- At the second step, 2 is inserted. It cannot be inserted in the box (1,2) as  $2 > 3$ , so it is inserted in the box (1,1) and thus pushes down 3 into a new box (2,1).
- At the third step, 6 is inserted in the box (1,2).
- At the fourth step, 1 is inserted in the box (1,1), and thus pushes down 2 into the second row. 2, in its turn, pushes down 3 to the third row into a new box (3,1).
- At the fifth step, 5 is inserted in the box (1,2), and thus pushes down 6 into the second row, where 6 forms a new box (2,2).
- At the sixth step, 7 is inserted in the box (1,3).
- At the last step, 4 is inserted in the box (1,2), and thus pushes down 5 into the second row. 5, in its turn, pushes down 6 to the third row into a new box (3,2).

As a result, we get the following SYT of shape  $\lambda = (3, 2, 2)$ :

$$P = \begin{array}{|c|c|c|} \hline 1 & 4 & 7 \\ \hline 2 & 5 & \\ \hline 3 & 6 & \\ \hline \end{array}$$

Now,  $Q$  records the number of the step each box of  $P$  was introduced for the first time. For example, the box (1,3) was introduced at the sixth step, so there will be 6 in it, and the box (3,1) has shown up at the fourth step, so there will be 4 there. As a result, we have

$$Q = \begin{array}{|c|c|c|} \hline 1 & 3 & 6 \\ \hline 2 & 5 & \\ \hline 4 & 7 & \\ \hline \end{array}$$

Let us formalize the procedure. The step (inserting  $w_i$  to the tableau  $P_{i-1}$  obtained after inserting  $w_{i-1}$ ) consists of the following:

- if  $w_i$  is greater than all entries in the first row of  $P_{i-1}$  then  $w_i$  is added to the end of the first row;
- otherwise, take the minimal  $w_j$  in the first row which is greater than  $w_i$ , substitute  $w_j$  with  $w_i$ , and insert  $w_j$  in the tableau obtained by removing the first row from  $P_{i-1}$ ; i.e., if  $w_j$  is greater than all entries in the second row of  $P_{i-1}$ , then  $w_j$  is added to the end of the second row; otherwise, find the minimal  $w_k$  in the second row which is greater than  $w_j$ , substitute  $w_k$  with  $w_j$ , and push  $w_k$  into the third row, etc.

**Example.** Show that  $P$  and  $Q$  are SYT.

Both  $P$  and  $Q$  are SYT of the same shape  $\lambda$ , which is called the *Schensted shape* of  $w$ .

To verify that the procedure works, we need to check that the map  $w \mapsto (P, Q)$  is injective and surjective. For this, we can construct the inverse map.

**Example 5.4.** Let us take  $P = P_7$  and  $Q = Q_7$  we got in Example 5.3, and try to reconstruct  $w \in S_7$ . According to  $Q$ , the last box to appear was  $(3, 2)$  (with 6 in it). It is in row 3, so 6 was pushed down by some element from row 2. The maximal number in row 2 which is less than 6 is 5, which implies that 6 was pushed down by 5, and 5 was pushed down to row 2 by someone from the first row. The maximal number in the first row which is less than 5 is 4, which implies that 5 was pushed down by 4, and thus 4 is precisely the number inserted at the last step.

Therefore, we have found  $w_7 = 4$  and reconstructed  $P_6$  (and, obviously,  $Q_6$ ), where

$$P_6 = \begin{array}{|c|c|c|} \hline 1 & 5 & 7 \\ \hline 2 & 6 & \\ \hline 3 & & \\ \hline \end{array} \quad Q_6 = \begin{array}{|c|c|c|} \hline 1 & 3 & 6 \\ \hline 2 & 5 & \\ \hline 4 & & \\ \hline \end{array}$$

We now repeat this procedure for the box  $(1, 3)$  of  $(P_6, Q_6)$  which, according to  $Q_6$ , is the last one to appear, etc. After seven steps we will reconstruct the permutation  $w$ .

## 5.2 Properties of RSK

We will now look at some corollaries of the algorithm, as well as at some of its properties.

**Theorem 5.5.** *Let  $\lambda$  be the Schensted shape of  $w \in S_n$ , and let  $(P, Q)$  be the corresponding SYT. Then*

- (1)  $\lambda_1$  is the maximal size of an increasing subsequence in  $w$ ;
- (2)  $\lambda'_1$  (= the number of rows in  $\lambda$ ) is the maximal size of a decreasing subsequence in  $w$ .

**Remark.** Theorem 5.5 immediately implies Erdos-Szekeres Theorem.

**Remark.** The Schensted shape of  $w \in S_n$  is precisely the Young diagram  $\lambda$  constructed by the poset  $P_w$  in the Greene's Theorem. See Appendix 1 (written by Sergey Fomin) to Stanley's "Enumerative Combinatorics, vol. II".

**Exercise 5.6.** Denote by  $r_x(P)$  an insertion of  $x$  in a partial tableau  $P$  in the RSK algorithm. Suppose that during  $r_x(P)$  the elements  $x_1, \dots, x_k$  are pushed down from rows  $1, 2, \dots, k$  and columns  $j_1, j_2, \dots, j_k$  respectively. Then

- (a)  $x < x_1 < \dots < x_k$ ;
- (b)  $j_1 \geq \dots \geq j_k$ ;
- (c) if  $P' = r_x(P)$ , then  $P'_{i,j} \leq P_{i,j}$  for all  $i, j$ .

Statement (1) of Theorem 5.5 immediately follows from the following lemma.

**Lemma 5.7.** *Let  $w = w_1 w_2 \dots w_n \in S_n$ , denote by  $P_k$  a partial tableau obtained after insertion of  $w_1, \dots, w_k$ . Let  $w_k$  enter  $P_{k-1}$  in column  $j$ . Then the longest increasing subsequence ending in  $w_k$  has length  $j$ .*

*Proof.* We use induction on  $k$ . The base ( $k = 1$ ) is obvious.

Let us first prove the existence of an increasing subsequence of length  $j$  ending in  $w_k$ . Let  $w_i$  be the element of  $P_{k-1}$  in the box  $(1, j-1)$ . Since  $w_k$  is inserted in the box  $(1, j)$ ,  $w_i < w_k$ . By induction, there exists an increasing subsequence ending in  $w_i$  of length  $j-1$ . By adding  $w_k$ , we obtain an increasing subsequence ending in  $w_k$  of length  $j$ .

No, let us prove the maximality. Suppose there exists an increasing subsequence ending in  $w_k$  of length greater than  $j$ . Take this subsequence, and let  $w_i$  be the element preceding  $w_k$ ,  $i < k$ . By the induction assumption,  $w_i$  is inserted in box  $(1, m)$ ,  $m \geq j$  (as there exists an increasing subsequence ending in  $w_i$  of length at least  $j$ ). Therefore, for the element  $w_p$  in the box  $(1, j)$  of  $P_i$  we have  $w_p \leq w_i < w_k$ . Let  $w_q$  be the element in the box  $(1, j)$  of  $P_{k-1}$ . By Exercise 5.6(c),  $w_q \leq w_p$  (as  $k-1 \geq i$ ), and thus  $w_q < w_k$ . However, during the insertion of  $w_k$  it pushes down  $w_q$ , which implies that  $w_k < w_q$ , so we came to a contradiction. □

We are left to prove statement (2) of the theorem.

Recall that we denote by  $r_x(P)$  a step of the RSK algorithm consisting of inserting  $x$  in a partial tableau  $P$ . Define  $c_x(P)$  as a (purely formal) *row insertion* of  $x$  into  $P$ . This can be understood as transposing  $P$ , then doing  $r_x(P^t)$ , and then transposing the result again (here by the transpose  $P^t$  of  $P$  we mean the reflection of  $P$  with respect to the main diagonal).

**Lemma 5.8.** *Let  $P$  be a partial tableau,  $x, y \notin P$ . Then  $c_y r_x(P) = r_x c_y(P)$ .*

*Proof.* The proof is case-by-case, we will consider some and leave others as an exercise.

Assume first that  $y > x$  and  $y$  is greater than all elements of  $P$ . Then  $c_y$  places  $y$  at the end of the first column, and thus  $r_x c_y(P)$  is just  $r_x(P)$  with additional  $y$  attached to the bottom of the first column. It is clear that  $c_y r_x(P)$  does precisely the same.

If we assume that  $x$  is the maximal element, then the same proof works (just need to transpose the whole picture).

**Exercise.** Complete the proof of the lemma. □

**Lemma 5.9.** *Let  $w = w_1 w_2 \dots w_n \in S_n$ , and let  $P$  be the insertion tableau of  $w$ . Define  $w^r = w_n w_{n-1} \dots w_2 w_1$ . Then the insertion tableau of  $w^r$  is  $P^t$ .*

*Proof.* We can write  $P = P(w) = r_{w_n} r_{w_{n-1}} \dots r_{w_2} r_{w_1}(\emptyset)$ .

Following this, and by using Lemma 5.8, we have

$$P(w^r) = r_{w_1} r_{w_2} \dots r_{w_n}(\emptyset) = r_{w_1} r_{w_2} \dots r_{w_{n-1}} c_{w_n}(\emptyset) = c_{w_n} r_{w_1} r_{w_2} \dots r_{w_{n-1}}(\emptyset)$$

We can now continue, with

$$P(w^r) = c_{w_n} r_{w_1} r_{w_2} \dots r_{w_{n-1}}(\emptyset) = c_{w_n} r_{w_1} r_{w_2} \dots c_{w_{n-1}}(\emptyset) = c_{w_n} c_{w_{n-1}} r_{w_1} r_{w_2} \dots r_{w_{n-2}}(\emptyset)$$

Applying this transformation  $n$  times, we get

$$P(w^r) = r_{w_1} r_{w_2} \dots r_{w_n}(\emptyset) = c_{w_n} c_{w_{n-1}} \dots c_{w_2} c_{w_1}(\emptyset) = P^t$$

as required. □

We can now complete the proof of Theorem 5.5(2).

Consider  $w^r$ , its increasing subsequences are precisely decreasing subsequences of  $w$ . Therefore, the maximal length of a decreasing subsequence of  $w$  is equal to the maximal length of an increasing subsequence of  $w^r$ , which, by (1), is equal to  $(\lambda')_1$ , where  $\lambda'$  is the Schensted shape of  $w^r$ . By Lemma 5.9,  $\lambda'$  is conjugate to  $\lambda$ , so  $(\lambda')'_1$  is precisely the size of the first column of  $\lambda$ .

We will now explore more symmetries of RSK.

**Example 5.10.** Recall from Example 5.3 that for  $w = 3261574 \in S_7$  we have

$$P = \begin{array}{|c|c|c|} \hline 1 & 4 & 7 \\ \hline 2 & 5 & \\ \hline 3 & 6 & \\ \hline \end{array} \quad Q = \begin{array}{|c|c|c|} \hline 1 & 3 & 6 \\ \hline 2 & 5 & \\ \hline 4 & 7 & \\ \hline \end{array}$$


Let now take the insertion tableau  $P' = Q$  and the recording tableau  $Q' = P$ , which permutation does this pair correspond to? An application of the RSK algorithm leads to  $w' = 4217536 \in S_7$ . Now, observe that  $w = (13674)(2)(5)$  and  $w' = (14763)(2)(5)$ , so  $w' = w^{-1}$ .

**Theorem 5.11.** *Let the application of the RSK takes  $w$  to  $(P, Q)$ . Then  $w^{-1}$  is taken to  $(Q, P)$ .*

Recall that, given  $w \in S_n$ , one can define a poset  $P_w$  on  $[n]$  with order  $w_i <_{P_w} w_j$  if  $w_i < w_j$  and  $i < j$ . Then chains of  $P_w$  are increasing subsequences of  $w$ , and antichains are decreasing subsequences of  $w$ .

Denote by  $P_1$  the set of minimal elements of  $P_w$ , by  $P_2$  the set of minimal elements of  $P_w \setminus P_1$ , and then by  $P_i$  the set of minimal elements of  $P_w \setminus \bigcup_{j < i} P_j$ . Note that every  $P_i$  is an antichain of  $P_w$ .

**Example.** Let  $w = 3261574 \in S_7$ , we have already seen that the Hasse diagram of  $P_w$  is



Then  $P_1 = \{1, 2, 3\}$ ,  $P_2 = \{4, 5, 6\}$ , and  $P_3 = \{7\}$  (cf columns of the the insertion tableau).

Write  $w$  in 2-line notation:  $w = \begin{array}{cccccc} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 3 & 2 & 6 & 1 & 5 & 7 & 4 \end{array}$ . Then we can interpret elements of  $P_w$  as  $\begin{matrix} i \\ w_i \end{matrix}$ . In particular, we can write every  $P_i$  ordering its elements by the top number:

$$P_1 = \left\{ \begin{matrix} 1 & 2 & 4 \\ 3 & 2 & 1 \end{matrix} \right\} = \left\{ \begin{matrix} u_{11} & u_{12} & u_{13} \\ w_{11} & w_{12} & w_{13} \end{matrix} \right\}$$

$$P_2 = \left\{ \begin{matrix} 3 & 5 & 7 \\ 6 & 5 & 4 \end{matrix} \right\} = \left\{ \begin{matrix} u_{21} & u_{22} & u_{23} \\ w_{21} & w_{22} & w_{23} \end{matrix} \right\}$$

$$P_3 = \left\{ \begin{matrix} 6 \\ 7 \end{matrix} \right\} = \left\{ \begin{matrix} u_{31} \\ w_{31} \end{matrix} \right\}$$

where  $\begin{matrix} u_{ij} \\ w_{ij} \end{matrix}$  denotes  $j$ -th element of  $P_i$ .

Note that the first row of the insertion tableau  $P$  is  $147 = w_{13} w_{23} w_{31}$ , and the first row of the recording tableau  $Q$  is  $136 = u_{11} u_{21} u_{31}$ .

**Remark.** In general, if  $P_j = \left\{ \begin{matrix} u_{j1} & u_{j2} & \dots & u_{jn_j} \\ w_{j1} & w_{j2} & \dots & w_{jn_j} \end{matrix} \right\}$  (where we always assume that  $u_{j1} < u_{j2} < \dots < u_{jn_j}$ ), then  $w_{j1} > w_{j2} > \dots > w_{jn_j}$  as all these elements compose an antichain of  $P_w$ .

**Remark.** While applying RSK to a permutation  $w = \begin{matrix} 1 & 2 & \dots & n \\ w_1 & w_2 & \dots & w_n \end{matrix}$  we insert the elements of the second row into tableau  $P$  and the elements of the first row into tableau  $Q$ . Note that the algorithm is completely defined by the relations  $w_i < w_j$  for every pair  $(i, j)$ . Thus, if we substitute the first row by any increasing sequence, and the second row by any  $n$  distinct numbers with the same relations, then we will get tableaux  $P'$  and  $Q'$  of the same shape as of  $P$  and  $Q$  (these are not SYT anymore, although the numbers will still be increasing to the right and to the bottom). Moreover, we can apply RSK to any two-row array of numbers (such that numbers in every row are distinct): sort the columns so that the top row is increasing, and then apply the procedure described above. We will use this modification of RSK to complete the proof of the theorem.

**Lemma 5.12.** *The first row of the insertion tableau  $P$  is  $w_{1n_1} w_{2n_2} w_{3n_3} \dots$ , and the first row of the recording tableau  $Q$  is  $u_{11} u_{21} u_{31} \dots$ . In other words, the first rows of  $P$  and  $Q$  consist of minimal  $w$  and  $u$  from every  $P_j$ .*

*Proof.* Let  $w = w_1 w_2 \dots w_n \in S_n$ , we use induction on  $n$ . If  $n = 1$  then the statement is trivial.

Denote  $\tilde{w} = w_1 w_2 \dots w_{n-1}$ , and let  $\tilde{P}$  and  $\tilde{Q}$  be the corresponding insertion and recording tableaux (see the remark above: here the first row of  $\tilde{w}$  is still  $[n-1]$ , but the second row may miss any one element of  $[n]$ ). Let  $\tilde{P}_j$  be the corresponding antichains of  $P_{\tilde{w}}$ ,  $\tilde{P}_j = \left\{ \begin{matrix} \tilde{u}_{j1} & \tilde{u}_{j2} & \dots & \tilde{u}_{jm_j} \\ \tilde{w}_{j1} & \tilde{w}_{j2} & \dots & \tilde{w}_{jm_j} \end{matrix} \right\}$ , where  $j = 1, \dots, l$ . By the induction assumption, the first row of  $\tilde{P}$  is  $\tilde{w}_{1m_1} \tilde{w}_{2m_2} \dots \tilde{w}_{lm_l}$ , and the first row of  $\tilde{Q}$  is  $\tilde{u}_{11} \tilde{u}_{21} \dots \tilde{u}_{l1}$ .

Insert  $w_n$  in  $\tilde{P}$ . If  $w_n > \tilde{w}_{jm_j}$  for all  $j$ , then we insert  $w_n$  in the  $(l+1)$ -st column. At the same time, this means that  $\begin{matrix} n \\ w_n \end{matrix}$  cannot be added to any of  $l$  antichains  $P_j$ , so it forms a new antichain  $P_{l+1}$  in  $P_w$ , and thus we can write  $\begin{matrix} n \\ w_n \end{matrix} = \begin{matrix} u_{l+11} \\ w_{l+11} \end{matrix} = \begin{matrix} u_{l+1m_{l+1}} \\ w_{l+1m_{l+1}} \end{matrix}$ , so the first row of  $P$  is  $w_{1m_1} w_{2m_2} \dots w_{lm_l} w_{l+1m_{l+1}}$ , and the first row of  $Q$  is  $u_{11} u_{21} \dots u_{l1}, u_{l+11}$ , as desired.

Assume now that  $w_n < \tilde{w}_{jm_j}$  for some  $j$ . Then  $\tilde{P}_j \cup \left\{ \begin{matrix} n \\ w_n \end{matrix} \right\}$  is an antichain of  $P_w$  (note that there may be many such  $j$ ). One can easily see that  $\begin{matrix} n \\ w_n \end{matrix}$  belongs to  $P_j$  with minimal  $j$  amongst those with  $w_n < \tilde{w}_{jm_j}$ . For example, if  $w_n < \tilde{w}_{1m_1}$ , then  $\begin{matrix} n \\ w_n \end{matrix}$  is incomparable with all minimal elements of  $P_w$ , and thus it is a minimal element itself. Therefore, the element  $\begin{matrix} n \\ w_n \end{matrix}$  is inserted in column  $j$  if and only if  $\begin{matrix} n \\ w_n \end{matrix} = \begin{matrix} u_{jm_j} \\ w_{jm_j} \end{matrix}$ , so again the first row of  $P$  is of the form required in the lemma. Since we do not start the new column,  $Q$  stays intact, so it also has the required form. □

**Exercise 5.13.** The poset  $P_w$  is isomorphic to the poset  $P_{w^{-1}}$ , with the isomorphism given by  $\begin{matrix} i \\ w_i \end{matrix} \mapsto \begin{matrix} w_i \\ i \end{matrix}$ .

*Proof of Theorem 5.11.* Let the insertion and recording tableaux of  $w^{-1}$  be  $P'$  and  $Q'$ . Denote by  $P'_j$  the corresponding antichains of the poset  $P_{w^{-1}}$ . According to Exercise 5.13,  $P'_j = \left\{ \begin{matrix} w_{j1} & w_{j2} & \dots & w_{jn_j} \\ u_{j1} & u_{j2} & \dots & u_{jn_j} \end{matrix} \right\}$ , so the minimal elements in the second row are  $u_{j1}$ , and the minimal elements in the first row are  $w_{jn_j}$ . According to Lemma 5.12, the first row of  $P'$  is now  $u_{11} u_{21} u_{31} \dots$ , and the first row of  $Q'$  is  $w_{1n_1} w_{2n_2} w_{3n_3} \dots$ , which are precisely first rows of  $Q$  and  $P$  respectively.

Now the plan is the following: denote by  $\bar{P}$  and  $\bar{Q}$  the tableaux  $P$  and  $Q$  with their first rows removed, and find a two-rows array  $\bar{w}$  (i.e. a “permutation”) which results in  $(\bar{P}, \bar{Q})$  under RSK. We already know the set of elements in both rows, the question is how to match them.

Consider two elements  $\begin{smallmatrix} u_{jq} \\ w_{jq} \end{smallmatrix}$ ,  $q < m_j$ , and  $\begin{smallmatrix} u_{rs} \\ w_{rs} \end{smallmatrix}$ ,  $s < m_r$ , then both  $w_{jq}$  and  $w_{rs}$  are in  $\bar{P}$ . Thus, there are two elements which pushed these two down.

**Exercise.**  $\begin{smallmatrix} u_{jq} \\ w_{jq} \end{smallmatrix}$  is pushed down from the first row of  $P$  by  $\begin{smallmatrix} u_{jq+1} \\ w_{jq+1} \end{smallmatrix}$ .

Therefore,  $w_{jq}$  enters  $\bar{P}$  before  $w_{rs}$  if and only if  $u_{jq+1} < u_{rs+1}$ . This implies that the following array  $\bar{w}$  is taken to  $(\bar{P}, \bar{Q})$  (check this!):

$$\bar{w} = \begin{array}{ccccccc} u_{12} \dots & u_{1m_1} & u_{22} \dots & u_{2m_2} & \dots & u_{l2} \dots & u_{lm_l} \\ w_{11} \dots & w_{1m_1-1} & w_{21} \dots & w_{2m_2-1} & \dots & w_{l1} \dots & w_{lm_l-1} \end{array}$$

for some  $l$ , where the columns should be permuted for the first row to be increasing.

Now, performing a similar exercise for  $w^{-1}$ , we see that

$$\overline{w^{-1}} = \begin{array}{ccccccc} w_{1m_1-1} \dots & w_{11} & w_{2m_2-1} \dots & w_{21} & \dots & w_{lm_l-1} \dots & w_{l1} \\ u_{1m_1} & \dots & u_{12} & u_{2m_2} & \dots & u_{22} & \dots & u_{lm_l} & \dots & u_{l2} \end{array}$$

Observe that  $\overline{w^{-1}} = (\bar{w})^{-1}$ , and thus, by the induction assumption,  $\bar{P}' = \bar{Q}$  and  $\bar{Q}' = \bar{P}$ . As we have already proved that the first rows also coincide, this implies that  $P' = Q$  and  $Q' = P$ . □

## 6 Games on graphs

Let  $G = (V, E)$  be a graph with the set of vertices  $V$  (where  $|V| = n$  and the vertices are identified with elements of  $[n]$ ), and the set of edges  $E$ , where  $E \subset V \times V$  (we remove the diagonal from  $V \times V$  and identify  $(i, j)$  with  $(j, i)$ ).

Denote by  $N(i)$  the set of *neighbors* of  $i$ , i.e. vertices connected to  $i$  by an edge. A *configuration* is a non-negative integer vector  $\mathbf{c} = (c_1, \dots, c_n)$  – this can be understood as we put  $c_i$  chips in a vertex  $i$ .

### 6.1 Reflection game

We call a vertex  $i$  *unstable* if  $2c_i < \sum_{j \in N(i)} c_j$ , and *stable* otherwise.

The *initial configuration* is a vector with all  $c_j = 0$  except for a single  $c_i = 1$ .

A *move* consists of choosing any unstable vertex  $i$  and changing the configuration as follows:  $c_i \mapsto -c_i + \sum_{j \in N(i)} c_j$ , while all other  $c_j$  stay intact. The *goal* of the game is to make every vertex stable.

**Example.** For  $G$  being a path with three vertices, there are three possible initial configurations, they all lead to a final configuration  $(1, 1, 1)$ .

We can ask several questions:

- For which graphs the game can stop?
- For which graphs the game stops for any sequence of moves?
- What are possible final configurations?

**Example.** For  $G$  being a cycle of length three, the game never stops.

**Exercise.** • A *subgraph* of  $G = (V, E)$  consists of a subset  $V'$  of  $V$  and all edges from  $E$  joining elements of  $V'$ . Show that if the game never stops on some subgraph of  $G$ , then it never stops on  $G$ .

- Show that if  $G$  has a cycle, then the game never stops.
- The *valence* of a vertex of  $G$  is the number of edges incident to it. Show that if  $G$  has a vertex of valence at least four, then the game never stops.
- Show that if  $G$  has at least two vertices of valence three, then the game never stops.
- Let  $G$  be a tree with a unique vertex  $v_0$  of valence three. Denote the lengths of the “legs” (including  $v_0$ ) by  $p, q, r$ , see Fig. 6.1. Show that if  $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} \leq 1$  then the game never stops.




Figure 6.1: Tree with precisely one vertex of valence at least three

The remaining graphs are shown in Fig. 6.2.

**Exercise.** Show that for the graphs in Fig. 6.2 any sequence of moves terminates, and the final configuration is always the same (such graphs are called *graphs of finite type*).




Figure 6.2: Graphs of finite type

## 6.2 Cartan firing

We now call a vertex  $i$  *unstable* if  $c_i > 1$ , and *stable* otherwise.

We can start with any initial configuration.

A *move* consists of choosing any unstable vertex  $i$  and changing the configuration as follows:  $c_i \mapsto c_i - 2$ ,  $c_j \mapsto c_j + 1$  if  $j \in N(i)$ , and  $c_j$  stays intact otherwise. The goal of the game is to make every vertex stable.

**Example.** For  $G$  being a path with three vertices and the initial configuration  $(2, 3, 2)$ , there is a sequence of moves which terminates.

Given a graph  $G$ , define its *Cartan matrix*  $A_G = (a_{ij})$  as follows:  $a_{ii} = 2$ ,  $a_{ij} = -1$  if  $i \in N(j)$ , and  $a_{ij} = 0$  otherwise. Then the move at vertex  $i$  can be understood as  $\mathbf{c} \mapsto \mathbf{c} - A_i$ , where  $A_i$  is the  $i$ -th row of  $A_G$ .



**Theorem 6.1.** *Let  $G$  be a graph, and let  $A = A_G$  be its Cartan matrix. Then the following are equivalent:*

- (1) For any initial configuration and any sequence of moves the game stops.*
- (2) There exists a positive vector  $v = (v_1, \dots, v_n)$ ,  $v_i > 0$ , such that  $Av$  has also all coordinates positive.*
- (3)  $A$  is positive definite.*