

## Riemannian Geometry IV, Homework 7 (Week 18)

Due date for starred problems: **Tuesday, March 12.**

**7.1.** (a) Let  $c(t)$  be a geodesic, and  $c(t_0)$  is conjugate to  $c(t_1)$ . Let  $J$  be any Jacobi field along  $c$  vanishing at  $t_0$  and  $t_1$ . Show that  $J$  is orthogonal, i.e.  $\langle J(t), c'(t) \rangle \equiv 0$ .

(b) Show that the dimension of the space  $J_c^\perp$  of orthogonal vector fields along  $c$  is  $2n - 2$ .

**7.2.** ( $\star$ ) Let  $c : [0, 1] \rightarrow M$  be a geodesic, and  $J$  be a Jacobi field along  $c$ . Denote  $c(0) = p, c'(0) = v$ . Define a curve  $\gamma(s)$ ,

$$\gamma : (-\varepsilon, \varepsilon) \rightarrow M, \quad \gamma(0) = p, \gamma'(0) = J(0)$$

Define also a vector field  $V(s) \in \Gamma(\gamma^{-1}TM)$ , such that

$$V(0) = v, \quad \nabla_s V(0) = \nabla_t J(0),$$

and a variation  $F(s, t) = \exp_{\gamma(s)} tV(s)$ .

(a) Show that  $F(s, t)$  is a geodesic variation of  $c(t)$ .

(b) Show that  $\frac{\partial F}{\partial s}(0, 0) = \gamma'(0) = J(0)$ , and  $\nabla_t \frac{\partial F}{\partial s}(0, 0) = \nabla_s V(0) = \nabla_t J(0)$ .

(c) Deduce from (a) and (b) that every Jacobi field along a geodesic  $c(t)$  is a variational vector field of some geodesic variation of  $c$ .

**7.3.** Let  $c : [a, b] \rightarrow M$  be a geodesic. For  $t \in [a, b]$ , denote by  $J_c^t$  the space of Jacobi fields along  $c$  vanishing at  $t$ . Let  $a \leq t_0 < t_1 \leq b$ , and define a map  $\psi : J_c^{t_0} \rightarrow M_{c(t_1)}$  by  $\psi(J) = J(t_1)$ .

(a) Show that  $\psi$  is linear.

Now assume that  $c(t_0)$  is not conjugate to  $c(t_1)$ .

(b) Show that  $\psi$  is an isomorphism.

(c) Show that for any  $u_1 \in M_{c(t_1)}$  there exists a unique  $J \in J_c^{t_0}$  such that  $J(t_1) = u_1$ .

(d) Using the arguments similar to ones from (a)–(c), show that for any  $u_0 \in M_{c(t_0)}$  there exists a unique  $J \in J_c^{t_1}$  such that  $J(t_0) = u_0$ .

(e) Deduce from (c) and (d) that if  $c(t_0)$  is not conjugate to  $c(t_1)$ , then for any  $u_0 \in M_{c(t_0)}$  and  $u_1 \in M_{c(t_1)}$  there exists a unique Jacobi field  $J$  along  $C$  such that  $J(t_0) = u_0$  and  $J(t_1) = u_1$ .