

Riemannian Geometry IV, Solutions 6 (Week 16)

- 6.1.** (a) Let $c(t)$ be a geodesic, and let $c(t_0)$ be conjugate to $c(t_1)$. Let J be any Jacobi field along c vanishing at t_0 and t_1 . Show that J is orthogonal, i.e. $\langle J(t), c'(t) \rangle \equiv 0$.
- (b) Show that the dimension of the space J_c^\perp of orthogonal vector fields along c is $2n - 2$.

Solution:

- (a) We have proved in class that the function $t \mapsto \langle J(t), c'(t) \rangle$ is linear. Since it is equal to zero at two points t_0 and t_1 , it vanishes everywhere, so $J(t)$ is orthogonal.
- (b) Recall that $J(t)$ is orthogonal if and only if both $\langle J(0), c'(0) \rangle$ and $\langle \frac{D}{dt} J(0), c'(0) \rangle$ vanish. Each of these equations defines a codimension one subspace in $T_p M$, so the dimension of $J_c^\perp = (n - 1) + (n - 1) = 2n - 2$.
- 6.2.** (★) Let $c : [0, 1] \rightarrow M$ be a geodesic, and let J be a Jacobi field along c . Denote $c(0) = p, c'(0) = v$.

Define a curve $\gamma(s)$,

$$\gamma : (-\varepsilon, \varepsilon) \rightarrow M, \quad \gamma(0) = p, \gamma'(0) = J(0)$$

Define also a vector field $V(s) \in \mathfrak{X}_\gamma(M)$, such that

$$V(0) = v, \quad \frac{D}{ds} V(0) = \frac{D}{dt} J(0),$$

and a variation $F(s, t) = \exp_{\gamma(s)} tV(s)$.

- (a) Show that $F(s, t)$ is a geodesic variation of $c(t)$.
- (b) Show that $\frac{\partial F}{\partial s}(0, 0) = \gamma'(0) = J(0)$, and $\frac{D}{dt} \frac{\partial F}{\partial s}(0, 0) = \frac{D}{ds} V(0) = \frac{D}{dt} J(0)$.
- (c) Deduce from (a) and (b) that every Jacobi field along a geodesic $c(t)$ is a variational vector field of some geodesic variation of c .

Solution:

- (a) By the definition of the exponential map, for given s the curve $t \mapsto \exp_{\gamma(s)} tV(s)$ is a geodesic.
- (b) We have

$$\frac{\partial F}{\partial s}(0, 0) = \frac{\partial}{\partial s} \Big|_{s=0} \exp_{\gamma(s)} tV(s) \Big|_{t=0} = \frac{\partial}{\partial s} \Big|_{s=0} \exp_{\gamma(s)}(0) = \frac{\partial}{\partial s} \Big|_{s=0} \gamma(s) = \gamma'(0) = J(0),$$

and

$$\frac{D}{dt} \frac{\partial F}{\partial s}(0, 0) = \frac{D}{ds} \Big|_{s=0} \frac{\partial F}{\partial t} \Big|_{t=0}(s, 0) = \frac{D}{ds} \Big|_{s=0} V(s) = \frac{D}{ds} V(0) = \frac{D}{dt} J(0)$$

- (c) According to (a), the variation F is geodesic, thus its variational vector field $\frac{\partial F}{\partial s}(0, t)$ is Jacobi. By (b), $\frac{\partial F}{\partial s}(0, 0) = J(0)$, and $\frac{D}{dt} \frac{\partial F}{\partial s}(0, 0) = \frac{D}{dt} J(0)$, which means that $\frac{\partial F}{\partial s}(0, t) = J(t)$ due to the uniqueness theorem.

6.3. Jacobi fields and conjugate points on locally symmetric spaces

A Riemannian manifold (M, g) is called a *locally symmetric space* if $\nabla R = 0$ (see Exercise 9.3). Let (M, g) be an n -dimensional locally symmetric space and $c : [0, \infty) \rightarrow M$ be a geodesic with $p = c(0)$ and $v = c'(0) \in T_p M$. Prove the following facts:

- (a) Let X, Y, Z be parallel vector fields along c . Show that $R(X, Y)Z$ is also parallel.

(b) Let $K_v \in \text{Hom}(T_pM, T_pM)$ be the curvature operator defined by

$$K_v(w) = R(w, v)v.$$

Show that K_v is self-adjoint, i.e.,

$$\langle K_v(w_1), w_2 \rangle = \langle w_1, K_v(w_2) \rangle$$

for every pair of vectors $w_1, w_2 \in T_pM$.

(c) Choose an orthonormal basis $w_1, \dots, w_n \in T_pM$ that diagonalizes K_v , i.e.,

$$K_v(w_i) = \lambda_i w_i$$

(such a basis exists since K_v is self-adjoint). Let W_1, \dots, W_n be the parallel vector fields along c with $W_i(0) = w_i$ (i.e., $\{W_i\}$ form a parallel orthonormal basis along c). Show that for all $t \in [0, \infty)$

$$K_{c'(t)}(W_i(t)) = \lambda_i W_i(t).$$

(d) Let $J(t) = \sum_i J_i(t)W_i(t)$ be a Jacobi field along c . Show that Jacobi equation translates into

$$J_i''(t) + \lambda_i J_i(t) = 0, \quad \text{for } i = 1, \dots, n.$$

(e) Show that the conjugate points of p along c are given by $c(\pi k / \sqrt{\lambda_i})$, where k is any positive integer and λ_i is a positive eigenvalue of K_v .

Solution:

(a) We know that $\nabla R = 0$. Let $\frac{D}{dt}$ denote covariant derivative along c . Then we have, for parallel vector fields X, Y, Z along c that

$$\begin{aligned} 0 = \nabla R(X, Y, Z, c')(t) &= \frac{D}{dt} R(X(t), Y(t))Z(t) - \\ &\quad - \underbrace{R\left(\frac{D}{dt} X(t), Y(t)\right)Z(t)}_{=0} - R(X(t), \underbrace{\frac{D}{dt} Y(t)}_{=0})Z(t) - R(X(t), Y(t)) \underbrace{\frac{D}{dt} Z(t)}_{=0} = \\ &= \frac{D}{dt} R(X(t), Y(t))Z(t). \end{aligned}$$

This shows that $R(X, Y)Z$ is parallel.

(b) The symmetries of R yield

$$\langle K_v(w_1), w_2 \rangle = \langle R(w_1, v)v, w_2 \rangle = \langle R(w_2, v)v, w_1 \rangle = \langle K_v(w_2), w_1 \rangle = \langle w_1, K_v(w_2) \rangle.$$

(c) Since K_v is self-adjoint, we can find an orthonormal basis $w_1, \dots, w_n \in T_pM$ with $K_v(w_i) = \lambda_i w_i$. We know, by (a), that $K_{c'(t)}(W_i(t)) = R(c'(t), W_i(t))W_i(t)$ is parallel and, since $K_{c'(0)}(W_i(0)) = K_v(w_i) = \lambda_i w_i$, we must have

$$K_{c'(t)}(W_i(t)) = \lambda_i W_i(t),$$

since parallel vector fields V along c are uniquely determined by their initial values $V(0) \in T_pM$.

(d) Let J be a Jacobi field along c . Then J satisfies the Jacobi equation

$$\frac{D^2}{dt^2} J + R(J, c')c' = 0.$$

Since W_1, \dots, W_n form a parallel basis along c , we obtain, by taking inner product with W_i :

$$\begin{aligned} 0 &= \left\langle \frac{D^2}{dt^2} J, W_i \right\rangle + \langle R(J, c')c', W_i \rangle = \\ &= \frac{d^2}{dt^2} \sum_j J_j \langle W_j, W_i \rangle + \sum_j J_j \langle R(W_j, c')c', W_i \rangle = \\ &= J_i'' + \sum_j J_j \lambda_j \langle W_j, W_i \rangle = J_i'' + \lambda_i J_i. \end{aligned}$$

(e) The unique solution of $J_i''(t) + \lambda_i J_i(t) = 0$, $J_i(0) = 0$ (up to scalar multiples) is given by

$$J_i(t) = \begin{cases} \sin(t\sqrt{\lambda_i}) & \text{if } \lambda_i > 0, \\ t & \text{if } \lambda_i = 0, \\ \sinh(t\sqrt{-\lambda_i}) & \text{if } \lambda_i < 0. \end{cases}$$

So J_i has zeros for positive t only if $\lambda_i > 0$, and these are precisely at $t = \pi k / \sqrt{\lambda_i}$. The corresponding Jacobi fields with $J(0) = 0$ and $\frac{D}{dt}J(0) = w_i$ produce the conjugate points $c(\pi k / \sqrt{\lambda_i})$.