

Solitons III
(2022-23)

Exercises

September 23, 2022

0 Introduction

Ex 1 Numerical results seen in the lectures suggest that the KdV equation

$$u_t + 6uu_x + u_{xxx} = 0 \quad (0.1)$$

has an exact solution of the form

$$u(x, t) = \frac{2}{\cosh^2(x - vt)} \quad (0.2)$$

for some constant velocity v . Verify this by direct substitution into the KdV equation and determine the value of v .

- Ex 2**
1. Show that if $u(x, t) = g(x, t)$ solves the KdV equation (0.1), then so does $u(x, t) = Ag(Bx, Ct)$, provided that the constants B and C are related to A in a specific way (which you should determine).
 2. Apply this transformation to the basic KdV solution found in exercise 1 to construct a one-parameter family of one-soliton solutions of the KdV equation.
 3. Find a formula relating the velocities to the heights for solitons in this one-parameter family. How does the width of a soliton in this family change if its velocity is rescaled by a factor of 4?

Ex 3 Show that if $u(x, t)$ solves the KdV equation and ϵ is a constant, then $v(x, t) := \frac{1}{\epsilon}u(x, t)$ solves the rescaled KdV equation

$$v_t + 6\epsilon v v_x + v_{xxx} = 0, \quad (0.3)$$

while $w(x, t) := \epsilon u(x, \epsilon t)$ solves the differently-rescaled KdV equation

$$w_t + 6w w_x + \epsilon w_{xxx} = 0. \quad (0.4)$$

- Ex 4** Consider a pair of solitons with velocities m and n in the ball and box model, with $m > n$ and the faster soliton to the left of the slower one, with separation $l \geq n$ (*i.e.* there are $l \geq n$ empty boxes between the two solitons). Evolve various such initial conditions forward in time using the ball and box rule, for different values of m , n and l . Prove that the system always evolves into an oppositely-ordered pair of the same two solitons, and find a general formula for the phase shifts¹ of the solitons in terms of m and n .
[**Optional:** What can go wrong if $l < n$? [**Hint:** Evolve the system backwards...]]

- Ex 5** In the two-colour (blue and red) ball and box model, we'll call a row of n consecutive balls a soliton if it keeps its form over time, so that after each time-step its only change is a possible (fixed) translation. There's no need for both colours to be represented, so a row of n blue balls, or a row of n red balls, is also a potential soliton. How many solitons of length n are there? What are their speeds?

- Ex 6*** The ball and box model can be further generalised to the M -colour ball and box model. The balls now come in M colours, $1, 2, \dots, M$, and the time-evolution rule is generalised to say that first all balls of colour 1 are moved, then all of colour 2, and so on, with a single time-step being completed once all balls of all colours have been moved. How many solitons of length n are there in this model? Again, there is no need for every colour to be present in a given soliton. You might start by classifying the 'top-speed' solitons of length n , that is, those that move at speed n .

- Ex 7*** Investigate the scattering of solitons in the two-colour ball and box model. You should find that the lengths of top-speed solitons are preserved under collisions, but their forms can change. Try to formulate a general rule for this behaviour. Can you generalise it to the M -colour model?

¹The phase shift of a soliton is defined to be the shift of its position, at a time in the far future, relative to the position it would have had at the same time if the other soliton hadn't been there.

1 Waves, dispersion and dissipation

- Ex 8**
- Express d'Alembert's general solution of the wave equation $u_{tt} - u_{xx} = 0$ in terms of the initial conditions $u(x, 0) = p(x)$ and $u_t(x, 0) = q(x)$.
 - Find a relation between $p(x)$ and $q(x)$ which produces a single wave travelling to the right.

Ex 9 The wave profile

$$\phi(x, t) = \cos(k_1x - \omega(k_1)t) + \cos(k_2x - \omega(k_2)t) \quad (1.1)$$

is a superposition of two plane waves. Rewrite ϕ as a product of cosines, and use this to sketch the wave profile when $|k_1 - k_2| \ll |k_1|$. Find the velocity at which the envelope of the wave profile moves (the **group velocity**), again for $k_1 \approx k_2$; in the limit $k_1 \rightarrow k_2$ verify that this reduces to $d\omega/dk$, consistent with the result obtained in lectures.

Ex 10

- Completing the square, derive the formula

$$\int_{-\infty}^{+\infty} dk e^{-A(k-\bar{k})^2} e^{ikB} = \sqrt{\frac{\pi}{A}} e^{i\bar{k}B} e^{-B^2/(4A)}. \quad (1.2)$$

(You can quote the result $\int_{-\infty}^{+\infty} dk e^{-Ak^2} = \sqrt{\pi/A}$ for $A > 0$.)

- For the Gaussian wavepacket (where Re denotes the real part)

$$u(x, t) = \text{Re} \int_{-\infty}^{+\infty} dk e^{-a^2(k-\bar{k})^2} e^{i(kx - \omega(k)t)}, \quad (1.3)$$

expand $\omega(k)$ to second order in $k - \bar{k}$, and then use the result of part 1 to derive a better approximation for $u(x, t)$ than that obtained in lectures.

- Given that a function of the form $e^{-(x-x_0)^2/C}$ describes a profile centred at x_0 with width⁻² equal to the real part of C^{-1} , show that the result of part 2 is a wave profile moving at velocity $\omega'(\bar{k})$, with width² increasing with time as $4a^2 + \omega''(\bar{k})^2 t^2 / a^2$. (Hence, for $\omega'' \neq 0$, the wave disperses.)

Ex 11 Find the dispersion relation and the phase and group velocities for:

(a) $u_t + u_x + \alpha u_{xxx} = 0$;

$$(b) \quad u_{tt} - \alpha^2 u_{xx} = \beta^2 u_{ttxx}.$$

Ex 12 For which values of n does the equation

$$u_t + u_x + u_{xxx} + \frac{\partial^n u}{\partial x^n} = 0 \tag{1.4}$$

admit "physical" dissipation? (A wave is said to have physical dissipation if the amplitude of plane waves decreases with time.)

2 Travelling waves

Ex 13 Find (if possible) real non-singular travelling wave solutions of the following equations, satisfying the given boundary conditions:

1. Modified KdV (mKdV) equation:

$$\begin{aligned} u_t + 6u^2u_x + u_{xxx} &= 0 \\ u \rightarrow 0, \quad u_x \rightarrow 0, \quad u_{xx} \rightarrow 0 \quad \text{as } x \rightarrow \pm\infty. \end{aligned} \quad (2.1)$$

2. ‘Wrong sign’ mKdV equation:

$$\begin{aligned} u_t - 6u^2u_x + u_{xxx} &= 0 \\ u \rightarrow 0, \quad u_x \rightarrow 0, \quad u_{xx} \rightarrow 0 \quad \text{as } x \rightarrow \pm\infty. \end{aligned} \quad (2.2)$$

3. ϕ^4 theory:

$$\begin{aligned} u_{tt} - u_{xx} + 2u(u^2 - 1) &= 0 \\ u_t \rightarrow 0, \quad u_x \rightarrow 0, \quad u \rightarrow -1 \quad \text{as } x \rightarrow -\infty \\ u_t \rightarrow 0, \quad u_x \rightarrow 0, \quad u \rightarrow +1 \quad \text{as } x \rightarrow +\infty. \end{aligned} \quad (2.3)$$

4. ϕ^6 theory:

$$\begin{aligned} u_{tt} - u_{xx} + u(u^2 - 1)(3u^2 - 1) &= 0 \\ u_t \rightarrow 0, \quad u_x \rightarrow 0, \quad u \rightarrow 0 \quad \text{as } x \rightarrow -\infty \\ u_t \rightarrow 0, \quad u_x \rightarrow 0, \quad u \rightarrow 1 \quad \text{as } x \rightarrow +\infty. \end{aligned} \quad (2.4)$$

5. Burgers equation:

$$\begin{aligned} u_t + uu_x - u_{xx} &= 0 \\ u \rightarrow u_0, \quad u_x \rightarrow 0 \quad \text{as } x \rightarrow -\infty \\ u \rightarrow u_1, \quad u_x \rightarrow 0 \quad \text{as } x \rightarrow +\infty, \end{aligned} \quad (2.5)$$

where u_0 and u_1 are real constants with $u_0 > u_1 > 0$.

[Hint: Start by showing that the boundary conditions relate the velocity v of the travelling wave to the sum of the constants u_0 and u_1 .]

6. * Generalised KdV equation with $n = 1, 2, 3, \dots$:

$$\begin{aligned} u_t + (n+1)(n+2)u^n u_x + u_{xxx} &= 0 \\ u \rightarrow 0, \quad u_x \rightarrow 0, \quad u_{xx} \rightarrow 0 \quad \text{as } x \rightarrow \pm\infty. \end{aligned} \quad (2.6)$$

Ex 14* Using the analogy with the classical mechanics of a point particle moving in one spatial dimension, determine the qualitative behaviour of travelling wave solutions of the KdV equation on a circle, for which the integration constants A and B are non-zero.

3 Topological lumps and the Bogomol'nyi bound

Ex 15 This exercise involves the infinite chain of identical coupled pendula of section 2.3, whose equations of motion reduce to the sine-Gordon equation in the continuum limit $a \rightarrow 0$. We will simplify expression by setting $g = L = \frac{M}{a} = 1$. Let $\theta_n(t)$ be the angle to the vertical of the n -th pendulum ($n \in \mathbb{Z}$), which is hung at the position $x = na$ along the chain, at time t . The configuration of the system at time t is then specified by the collection of angles $\{\theta_n(t)\}_{n \in \mathbb{Z}}$.

1. Starting from the force (note: m is a dummy variable)

$$F_n(\{\theta_m\}) = -a \sin \theta_n + \frac{1}{a}(\theta_{n+1} - \theta_n) + \frac{1}{a}(\theta_{n-1} - \theta_n) \quad (3.1)$$

acting on the n -th pendulum, deduce the potential energy

$$V(\{\theta_m\}) = \sum_{n=-\infty}^{+\infty} (\dots) \quad (3.2)$$

such that $F_n = -\frac{\partial V}{\partial \theta_n}$ for all $n \in \mathbb{Z}$, and fix the integration constant by requiring that the potential energy be zero when all pendula point down: $V(\{0\}) = 0$.

2. Show that in the continuum limit $a \rightarrow 0$, the potential energy computed above becomes

$$V = \int_{-\infty}^{+\infty} dx \left[(1 - \cos \theta) + \frac{1}{2} \theta_x^2 \right], \quad (3.3)$$

and the kinetic energy

$$T(\{\theta_m\}) = \frac{a}{2} \sum_{n=-\infty}^{+\infty} \dot{\theta}_n^2 \quad (3.4)$$

becomes

$$T = \int_{-\infty}^{+\infty} dx \frac{1}{2} \dot{\theta}_t^2, \quad (3.5)$$

where the function $\theta(x, t)$ is the continuum limit of $\{\theta_n(t)\}_{n \in \mathbb{Z}}$.

[**Hint:** in the continuum limit, $a \sum_{n=-\infty}^{+\infty} \rightarrow \int_{-\infty}^{+\infty} dx$.]

Ex 16 A field $u(x, t)$ has kinetic energy T and potential energy V , where

$$\begin{aligned} T &= \int_{-\infty}^{+\infty} dx \frac{1}{2} u_t^2, \\ V &= \int_{-\infty}^{+\infty} dx \left[\frac{1}{2} u_x^2 + \frac{\lambda}{2} (u^2 - a^2)^2 \right], \end{aligned} \quad (3.6)$$

and a and $\lambda > 0$ are (real) constants. (This is a version of the ‘ ϕ^4 ’ theory. It’s called like that because the scalar potential is quartic, and the field u is usually called ϕ .) The equation of motion for u is

$$u_{tt} - u_{xx} + 2\lambda u(u^2 - a^2) = 0. \quad (3.7)$$

1. If u is to have finite energy, what boundary conditions must be imposed on u , u_x and u_t at $x = \pm\infty$?
2. Find the general travelling-wave solution(s) to the equation of motion, consistent with the boundary conditions found in part 1. Compute the total energy $E = T + V$ for these solutions. For which velocity do the solutions have the lowest energy?
3. One of the possible boundary conditions for part 1 implies that u is a kink, with $[u(x)]_{x=-\infty}^{x=+\infty} = 2a$. Use the Bogomol’nyi argument to show that the total energy $E = T + V$ of that configuration is bounded from below by $C\sqrt{\lambda}a^3$, where C is a constant that you should determine, and find the solution u which saturates this bound. Verify that this solution agrees with the lowest-energy solution of part 2.

Ex 17 1. Explain why the Bogomol’nyi argument given in the lectures fails to provide a useful bound on the energy of a two-kink solution of the sine-Gordon equation (a two-kink solution is one with topological charge $n - m$ equal to 2). What is the most that can be said about the energy of a k -kink?

2. For a sine-Gordon field u , generalise the Bogomol’nyi argument to show that

$$\int_A^B dx \left[\frac{1}{2} u_t^2 + \frac{1}{2} u_x^2 + (1 - \cos u) \right] \geq \pm 4 \left[\cos \frac{u}{2} \right]_A^B. \quad (3.8)$$

3. * Use this result and the intermediate value theorem (look it up if necessary!) to show that if the field u has the boundary conditions of a k -kink, then its energy is at least k times that of a single kink. Can this bound be saturated?

Ex 18 A system on the finite interval $-\pi/2 \leq x \leq \pi/2$ is defined by the following expressions for the kinetic energy T and the potential energy V :

$$\begin{aligned} T &= \int_{-\pi/2}^{\pi/2} dx \frac{1}{2} u_t^2 \\ V &= \int_{-\pi/2}^{\pi/2} dx \frac{1}{2} (u_x^2 + 1 - u^2) . \end{aligned} \tag{3.9}$$

The function $u(x, t)$ satisfies the boundary condition $|u(\pm\pi/2, t)| = 1$ and is required to satisfy $|u(x, t)| \leq 1$ everywhere. Show that with "kink" boundary conditions, the total energy E is bounded below by a positive constant, and find a solution for which the bound is saturated.

4 Conservation laws

Ex 19 Check explicitly that the energy

$$E = \int_{-\infty}^{+\infty} dx \left[\frac{1}{2} u_t^2 + \frac{1}{2} u_x^2 + \mathbb{V}(u) \right] \quad (4.1)$$

and the momentum

$$P = - \int_{-\infty}^{+\infty} dx u_t u_x \quad (4.2)$$

of a relativistic field $u(x, t)$ in 1 space and 1 time dimensions are conserved when the equation of motion

$$u_{tt} - u_{xx} = -\mathbb{V}'(u) \quad (4.3)$$

and the boundary conditions

$$u_t, u_x, \mathbb{V}(u), \mathbb{V}'(u) \xrightarrow{x \rightarrow \pm\infty} 0 \quad \forall t \quad (4.4)$$

are satisfied.

Ex 20 1. Compute the conserved topological charge, energy and momentum of a sine-Gordon kink moving with velocity v , and check that the results do not depend on time. [**Hint:** The integral (A.9) might be useful. For the scalar potential term in the energy, write $1 - \cos(u) = 2 \sin^2(u/2)$, plug in the kink solution and manipulate the result using trigonometric formulae until (A.9) becomes useful.] Confirm that for $|v| \ll 1$ the energy and the momentum take the forms

$$E = M + \frac{1}{2} M v^2 + \mathcal{O}(v^4), \quad P = M v + \mathcal{O}(v^3) \quad (4.5)$$

where the ‘mass’ M is the energy of the static kink, which appears in the Bogomol’nyi bound.

2. * If you are fearless and have time on your hands, try also to compute the conserved spin 3 charge

$$Q_3 = \int_{-\infty}^{+\infty} dx \left[u_{++}^2 - \frac{1}{4} u_+^4 + u_+^2 \cos u \right] \quad (4.6)$$

for the sine-Gordon kink. The integrals are not at all straightforward, but can be evaluated using appropriate changes of variables. (Did I write fearless?)

Ex 21 Find three conserved charges for the mKdV equation (2.1) of Ex 13.1, which involve u , u^2 and u^4 respectively. The boundary conditions on $u(x, t)$ are u , u_x and $u_{xx} \rightarrow 0$ as $|x| \rightarrow \infty$. Evaluate these quantities for the travelling-wave solution found in Ex 13.1. The list of definite integrals at the end of the problems sheet might help.

Ex 22 Show that u is a conserved density for Burgers' equation (2.5). Why is this result of no use in analysing the travelling wave solution of Ex 13.5?

Ex 23 Consider the KdV equation $u_t + 6uu_x + u_{xxx} = 0$ for the field $u(x, t)$.

1. Show that $\rho_1 \equiv u$, $\rho_2 \equiv u^2$ and $\rho_* \equiv xu - 3tu^2$ are all conserved densities, so that

$$Q_1 = \int_{-\infty}^{+\infty} dx u, \quad Q_2 = \int_{-\infty}^{+\infty} dx u^2, \quad Q_* = \int_{-\infty}^{+\infty} dx (xu - 3tu^2) \quad (4.7)$$

are all conserved charges.

2. Evaluate the conserved charges Q_1 , Q_2 and Q_* for the one-soliton solution centred at x_0 and moving with velocity $v = 4\mu^2$:

$$u_{\mu, x_0}(x, t) = 2\mu^2 \operatorname{sech}^2 [\mu(x - x_0 - 4\mu^2 t)] \quad (4.8)$$

3. According to the KdV equation, the initial condition $u(x, 0) = 6 \operatorname{sech}^2(x)$ is known to evolve into the sum of two well-separated solitons with different velocities $v_1 = 4\mu_1^2$ and $v_2 = 4\mu_2^2$ at late times. Use the conservation of Q_1 and Q_2 to determine v_1 and v_2 .
4. A two-soliton solution separates as $t \rightarrow -\infty$ into two one-solitons u_{μ_1, x_1} and u_{μ_2, x_2} . As $t \rightarrow +\infty$, two one-solitons are again found, with μ_1 and μ_2 unchanged but with x_1, x_2 replaced by y_1, y_2 . Use the conservation of Q_* to find a formula relating the *phase shifts* $y_1 - x_1$ and $y_2 - x_2$ of the two solitons.

Ex 24 1. Show that if $u(x, t)$ satisfies the KdV equation $u_t + 6uu_x + u_{xxx} = 0$, and $u = \lambda - v^2 - v_x$ where λ is a constant and $v(x, t)$ some other function, then v satisfies

$$\left(2v + \frac{\partial}{\partial x}\right)(v_t + 6\lambda v_x - 6v^2 v_x + v_{xxx}) = 0.$$

2. Compute the Gardner transform expansion

$$w(x, t) = \sum_{n=0}^{\infty} w_n(x, t) \varepsilon^n \quad (4.9)$$

up to order ε^4 . Use the results to find the conserved charges \tilde{Q}_3 and \tilde{Q}_4 , where

$$\tilde{Q}_n = \int_{-\infty}^{+\infty} dx w_n . \quad (4.10)$$

Show that \tilde{Q}_3 is the integral of a total x -derivative (and hence is zero), while $\tilde{Q}_4 = \alpha Q_3$, where

$$Q_3 = \int_{-\infty}^{+\infty} dx \left(u^3 - \frac{1}{2} u_x^2 \right) \quad (4.11)$$

is the third KdV conserved charge (the ‘energy’) and α a constant that you should determine. * If you’re feeling energetic, try to compute \tilde{Q}_5 and \tilde{Q}_6 as well.

Ex 25 This question is also about the KdV equation $u_t + 6uu_x + u_{xxx} = 0$.

1. Evaluate the first three KdV conserved charges

$$Q_1 = \int_{-\infty}^{+\infty} dx u , \quad Q_2 = \int_{-\infty}^{+\infty} dx u^2 , \quad Q_3 = \int_{-\infty}^{+\infty} dx \left(u^3 - \frac{1}{2} u_x^2 \right) \quad (4.12)$$

for the initial state $u(x, 0) = A \operatorname{sech}^2(Bx)$, where A and B are constants.

2. The initial state

$$u(x, 0) = N(N + 1) \operatorname{sech}^2(x) , \quad (4.13)$$

where N is an integer, is known to evolve at late times into N well-separated solitons, with velocities $4k^2$, $k = 1 \dots N$. So for $t \rightarrow +\infty$, this solution approaches the sum of N single well-separated solitons

$$u(x, t) \approx \sum_{k=1}^N 2\mu_k^2 \operatorname{sech}^2[\mu_k(x - x_k - 4\mu_k^2 t)] , \quad (4.14)$$

where μ_1, \dots, μ_N are N different constants. Since Q_1 , Q_2 and Q_3 are conserved, their values at $t = 0$ and $t \rightarrow +\infty$ must be equal. Use this fact to deduce formulae for the sums of the first N integers, the first N cubes, and the first N fifth powers.

3. * Use Q_4 and Q_5 and the method just described to find the sum of the first N seventh and ninth powers, $\sum_{k=1}^N k^7$ and $\sum_{k=1}^N k^9$.

5 The Bäcklund transform

Ex 26 1. Show that the pair of equations

$$\begin{aligned}(u - v)_+ &= \sqrt{2} e^{(u+v)/2} \\ (u + v)_- &= \sqrt{2} e^{(u-v)/2}\end{aligned}\tag{5.1}$$

provides a Bäcklund transformation linking solutions of $v_{+-} = 0$ (the wave equation in light-cone coordinates) to those of $u_{+-} = e^u$ (the Liouville equation).

2. Starting from d'Alembert's general solution $v = f(x^+) + g(x^-)$ of the wave equation, use the Bäcklund transform (5.1) to obtain the corresponding solutions of the Liouville equation for u . [**Hint:** Set $u(x^+, x^-) = 2U(x^+, x^-) + f(x^+) - g(x^-)$. You might simplify the notation by setting $f(x^+) = \log(F'(x^+))$ and $g(x^-) = -\log(G'(x^-))$, where prime means first derivative.]

Ex 27 Consider the Bäcklund transform

$$v_x + \frac{1}{2}uv = 0\tag{5.2}$$

$$v_t + \frac{1}{2}u_xv - \frac{1}{4}u^2v = 0.\tag{5.3}$$

1. Show that (5.2) and (5.3) together imply that v satisfies the linear heat equation $v_t = v_{xx}$, while u satisfies Burgers' equation $u_t + uu_x - u_{xx} = 0$. [**Hint:** for the first, solve (5.2) for u and substitute in (5.3); for the second, start by cross-differentiating.]
2. Find the *general* travelling-wave solution for $v(x, t)$ and, via the Bäcklund transform, re-obtain the travelling-wave for Burgers' equation found in question (2.5).
3. * The linear equation satisfied by $v(x, t)$ allows for the linear superposition of solutions. Use this fact, and your answers to part 2, to construct solutions for v and then u which describe the interaction of *two* travelling waves.
4. * Sketch your solutions functions of x at fixed times both before and after the interaction, and also draw their trajectories in the (x, t) plane, perhaps starting with the help of a computer. Are the travelling waves of Burgers' equation true solitons, in the sense given in lectures?

[**Hints:** Examine the asymptotics of the solution viewed from frames moving at various velocities V (that is, set $X_V = x - Vt$ and consider $t \rightarrow \pm\infty$ keeping X_V finite). This should allow you to isolate various travelling waves in these limits, and to decide whether they preserve their form under interactions. For definiteness, consider the case $c_1 > c_2 > 0$, where c_1 and c_2 are the velocities of the two separate travelling waves before they were superimposed. A further hint: as well as the 'expected' special values for V , namely c_1 and c_2 , be careful about what happens when $V = c_1 + c_2$.]

Ex 28 1. Show that the two equations

$$\begin{aligned} v_x &= -u - v^2 \\ v_t &= 2u^2 + 2uv^2 + u_{xx} - 2u_x v \end{aligned} \quad (5.4)$$

are a Bäcklund transform relating solutions of the KdV equation

$$u_t + 6uu_x + u_{xxx} = 0 \quad (5.5)$$

and the wrong sign modified KdV (mKdV) equation

$$v_t - 6v^2 v_x + v_{xxx} = 0. \quad (5.6)$$

(Note the appearance of the Miura transform in (5.4).)

2. Taking $u = c^2$, where c is a constant, as a seed solution of the KdV equation, find the corresponding solution of the wrong sign mKdV equation.

Ex 29 The 2-soliton solution of the sine-Gordon equation with Bäcklund parameters a_1 and a_2 is

$$u(x, t) = 4 \arctan \left(\mu \frac{e^{\theta_1} - e^{\theta_2}}{1 + e^{\theta_1 + \theta_2}} \right), \quad \theta_i = \varepsilon_i \gamma_i (x - v_i t - \bar{x}_i) \quad (5.7)$$

where $\mu = (a_2 + a_1)/(a_2 - a_1)$, $v_i = (a_i^2 - 1)/(a_i^2 + 1)$, $\gamma_i = 1/\sqrt{1 - v_i^2}$, $\varepsilon_i = \text{sign}(a_i)$, and \bar{x}_1 and \bar{x}_2 are constants, as in the lectures. Rewriting u as a function of $X_V \equiv x - Vt$ and t , show that, for $V \neq v_1, v_2$ (and $v_1 \neq v_2$)

$$\lim_{\substack{t \rightarrow \pm\infty \\ X_V \text{ finite}}} u = 2n\pi,$$

where n is an integer. If $v_2 > v_1 > 0$ and $\varepsilon_i = 1$, how does the parity of n (whether it is even or odd) depend on the value of v relative to v_1 and v_2 ?

[Hints: First show that $|\theta_i| \rightarrow +\infty$ as $t \rightarrow \pm\infty$; then consider each of the four possible options $(\theta_1, \theta_2) \rightarrow (+\infty, +\infty)$, $(-\infty, -\infty)$, $(+\infty, -\infty)$, $(-\infty, +\infty)$. Remember that $\arctan(0) = m\pi$ and $\arctan(\pm\infty) = \pm\pi/2 + m\pi$, where the ambiguities of $m\pi$, $m \in \mathbb{Z}$, encode the multivalued nature of the arctan function.]

Ex 30 Find the asymptotics of the 2-soliton sine-Gordon solution defined in equation (5.7), in the case $a_2 > a_1 > 0$, as $t \rightarrow \pm\infty$ with $X_{v_2} \equiv x - v_2 t$ held finite.

Ex 31 Show by direct analysis (as in the lectures) that taking a_1 and a_2 of opposite signs in (5.7) results in a two-kink, or two-antikink, solution to the sine-Gordon equation.

Ex 32

1. The argument of the arctangent in the sine-Gordon 2-soliton solution (5.7) is a continuous function of x for all $x \in \mathbb{R}$. Show that, in particular, it is never infinite. What does this imply about the range of u ? [**Hint:** consider the graph of $\tan u/4$.]
2. By taking the limits of this function as $x \rightarrow \pm\infty$ (with $t = \bar{x}_1 = \bar{x}_2 = 0$ for simplicity), show that the topological charge of the two-soliton solution (5.7) is 0 if $\text{sign}(a_1) = \text{sign}(a_2)$, and ± 2 if $\text{sign}(a_1) = -\text{sign}(a_2)$, in units where the topological charge of a kink is 1.

Ex 33 Consider the two-soliton solution of the sine-Gordon equation (5.7) with complex Bäcklund parameters $a_1 = \bar{a}_2 := a \in \mathbb{C}$ and with vanishing integration constants, as is appropriate to find the breather solution. Show that

$$\begin{aligned} \text{Re}(\theta_1) &= +\text{Re}(\theta_2) = \gamma(x - vt) \cos \varphi , \\ \text{Im}(\theta_1) &= -\text{Im}(\theta_2) = \gamma(vx - t) \sin \varphi , \end{aligned} \tag{5.8}$$

where $\varphi = \arg(a)$ and

$$\begin{aligned} v &= \frac{|a|^2 - 1}{|a|^2 + 1} \\ \gamma &= \frac{1}{\sqrt{1 - v^2}} = \frac{1 + |a|^2}{2|a|} . \end{aligned} \tag{5.9}$$

Ex 34 The *stationary* breather solution of the sine-Gordon equation (that is the breather solution with $v = 0$) has the form

$$\tan \frac{u}{4} = \frac{\cos \varphi}{\sin \varphi} \cdot \frac{\sin(t \sin \varphi)}{\cosh(x \cos \varphi)} . \tag{5.10}$$

Show that in the limit $\varphi \rightarrow 0$, in which the kink and antikink that form the breather are very loosely bound, the time period τ of a single oscillation of the breather scales like $\tau \sim |\varphi|^{-1}$, and the spatial size x_{\max} of the breather scales like $x_{\max} \sim -\log \varphi$.

[**Hint:** You could define x_{\max} as the value of x at which $\tan(u/4) = 1$ when the oscillatory factor in the numerator is at its maximum. Focus only on the parametric dependence on φ , ignoring all numerical factors.]

6 The Hirota method

Ex 35 We have seen in lectures that the KdV equation $u_t + 6uu_x + u_{xxx} = 0$ for the field $u(x, t)$ that describes the profile of a wave translates into the following equation for the new variable $w(x, t) = \int dx u$:

$$w_t + 3w_x^2 + w_{xxx} = 0 . \quad (6.1)$$

Let $w = 2\frac{\partial}{\partial x} \log f = 2\frac{f_x}{f}$ where $f(x, t)$ is a nowhere vanishing function of x and t , so that $u = 2\frac{\partial^2}{\partial x^2} \log f$. The aim of this exercise is to rewrite (6.1) as an equation for f .

1. Express w_t , w_x , w_{xx} and w_{xxx} in terms of f and its derivatives.
2. Show that the equation (6.1) can be rewritten as

$$ff_{xt} - f_x f_t + 3f_{xx}^2 - 4f_x f_{xxx} + ff_{xxxx} = 0 , \quad (6.2)$$

which is known as the *quadratic form* of the KdV equation.

Ex 36 The Hirota bilinear differential operator $D_t^m D_x^n$ is defined for any pair of natural numbers (m, n) by

$$D_t^m D_x^n (f, g) = \left(\frac{\partial}{\partial t} - \frac{\partial}{\partial t'} \right)^m \left(\frac{\partial}{\partial x} - \frac{\partial}{\partial x'} \right)^n f(x, t) g(x', t') \Bigg|_{\substack{x'=x \\ t'=t}} \quad (6.3)$$

and maps a pair of functions $(f(x, t), g(x, t))$ into a single function.

1. Prove that the Hirota operators $B_{m,n} := D_t^m D_x^n$ are bilinear, *i.e.* for all constants a_1, a_2

$$\begin{aligned} B_{m,n}(a_1 f_1 + a_2 f_2, g) &= a_1 B_{m,n}(f_1, g) + a_2 B_{m,n}(f_2, g) , \\ B_{m,n}(f, a_1 g_1 + a_2 g_2) &= a_1 B_{m,n}(f, g_1) + a_2 B_{m,n}(f, g_2) . \end{aligned} \quad (6.4)$$

2. Prove the symmetry property

$$B_{m,n}(f, g) = (-1)^{m+n} B_{m,n}(g, f) . \quad (6.5)$$

3. Compute the Hirota derivatives $D_t^2(f, g)$ and $D_x^4(f, g)$, and verify that your expression for the latter is consistent with the result for $D_x^4(f, f)$ given in lectures.

Ex 37 Define a “non-Hirota” bilinear differential operator $\tilde{D}_t^m \tilde{D}_x^n$ by

$$\tilde{D}_t^m \tilde{D}_x^n(f, g) = \left(\frac{\partial}{\partial t} + \frac{\partial}{\partial t'} \right)^m \left(\frac{\partial}{\partial x} + \frac{\partial}{\partial x'} \right)^n f(x, t)g(x', t') \Big|_{\substack{x'=x \\ t'=t}} \quad (6.6)$$

(note the plus signs!).

1. Compute $\tilde{D}_x(f, g)$ and $\tilde{D}_t(f, g)$, verifying that in both cases the answer is given by the corresponding ‘ordinary’ derivative of the product $f(x, t)g(x, t)$.
2. How does this result generalise for arbitrary non-Hirota differential operators (6.6)? Prove your claim.
3. Compare your answer with the Hirota operators defined above.

Ex 38 1. If $\theta_i = a_i x + b_i t + c_i$, prove that

$$D_t D_x(e^{\theta_1}, e^{\theta_2}) = (b_1 - b_2)(a_1 - a_2)e^{\theta_1 + \theta_2}.$$

2. Prove the corresponding result for $D_t^m D_x^n(e^{\theta_1}, e^{\theta_2})$, as quoted in lectures.

Ex 39 Prove that

$$D_t^m D_x^n(f, 1) = \frac{\partial^m}{\partial t^m} \frac{\partial^n}{\partial x^n} f. \quad (6.7)$$

Ex 40 Consider the function f , such that $u = 2 \frac{\partial^2}{\partial x^2} \log f$ is the KdV field, which corresponds to a 2-soliton solution:

$$f = 1 + \epsilon f_1 + \epsilon^2 f_2 = 1 + \epsilon (e^{\theta_1} + e^{\theta_2}) + \epsilon^2 \left(\frac{a_1 - a_2}{a_1 + a_2} \right)^2 e^{\theta_1 + \theta_2}, \quad (6.8)$$

where $\theta_i = a_i x - a_i^3 t + c_i$, with a_i and c_i constants. Check that $B(f_1, f_2) = 0$ and $B(f_2, f_2) = 0$, where $B = D_x(D_t + D_x^3)$, and show that this implies that the expansion (6.8), which is truncated at order ϵ^2 , is a solution of the bilinear form of the KdV equation.

Ex 41* Derive the solution of the bilinear form of the KdV equation $D_x(D_t + D_x^3)(f, f) = 0$ which represents the 3-soliton solution, in the form

$$f = 1 + \epsilon f_1 + \epsilon^2 f_2 + \epsilon^3 f_3 \quad (6.9)$$

where $f_1 = \sum_{i=1}^3 e^{\theta_i}$. [This includes proving that the higher order terms in the ϵ expansion can be consistently set to zero, as in Ex 40.]

Ex 42 Show that the Boussinesq equation

$$u_{tt} - u_{xx} - 3(u^2)_{xx} - u_{xxxx} = 0 \quad (6.10)$$

can be written in the bilinear form

$$(D_t^2 - D_x^2 - D_x^4)(f, f) = 0 \quad (6.11)$$

where $u = 2 \frac{\partial^2}{\partial x^2} \log f$.

Ex 43 Show that the following higher-dimensional version of the KdV equation,

$$(u_t + 6uu_x + u_{xxx})_x + 3\sigma^2 u_{yy} = 0 \quad (6.12)$$

for the field $u(x, y, t)$, also known as the Kadomtsev-Petviashvili (KP) equation, can be written in the bilinear form

$$(D_t D_x + D_x^4 + 3\sigma^2 D_y^2)(f, f) = 0 \quad (6.13)$$

where $u(x, y, t) = 2 \frac{\partial^2}{\partial x^2} \log f(x, y, t)$.

7 Exam-style problem

Ex 44 The complex field $u(x, t)$ obeys the equation

$$iu_t + \frac{1}{2}u_{xx} + |u|^2u = 0, \quad (7.1)$$

where $i = \sqrt{-1}$, and the boundary conditions

$$u, u_x, u_{xx} \rightarrow 0 \quad \text{as } x \rightarrow \pm\infty. \quad (7.2)$$

1. Show that the quantities

$$\begin{aligned} Q_1 &= \int_{-\infty}^{+\infty} dx |u|^2 \\ Q_2 &= \int_{-\infty}^{+\infty} dx \operatorname{Im}(\bar{u}u_x) \\ Q_3 &= \int_{-\infty}^{+\infty} dx \left(\frac{1}{2}|u_x|^2 + C|u|^4 \right) \end{aligned} \quad (7.3)$$

are conserved provided that the constant C takes a value that you should find. (Here Im denotes the imaginary part and a bar denotes complex conjugation.)

2. Show that given a ‘seed’ solution $u(x, t)$ of equation (7.1),

$$u^{(v)}(x, t) := u(x - vt, t)e^{i(Ax+Bt)} \quad (7.4)$$

is also a solution for all $v \in \mathbb{R}$, provided that the constants A and B depend on v in a way that you should find.

3. Determine the functional dependence of the conserved charges Q_1, Q_2, Q_3 in (7.3) on the parameter v that labels the one-parameter family of solution (7.4).
4. Find all solutions of the form

$$u(x, t) = \rho(x)e^{i\varphi(t)} \quad (7.5)$$

of equation (7.1) with boundary conditions (7.2), where ρ and φ are real and $u(x, 0)$ is a real even function of x . [You can use the integrals at the end of the problem sheet.] Apply the method of part 2 to this seed solution to find the associated one-parameter family of solutions $u^{(v)}(x, t)$.

A Useful integrals

You can freely quote the following formulae, although deriving them may be instructive:

- **Indefinite integrals:** [**Note:** the integration constant is in principle complex]

$$\int \frac{dx}{x\sqrt{1-x}} = -2\operatorname{arcsech}(\sqrt{x}) \quad (\text{A.1})$$

$$\int \frac{dx}{x\sqrt{1-x^2}} = -\operatorname{arcsech}(x) \quad (\text{A.2})$$

$$\int \frac{dx}{x\sqrt{1+x^2}} = -\operatorname{arccosech}(x) \quad (\text{A.3})$$

$$\int \frac{dx}{\sin(x/2)} = 2 \log \tan(x/4) \quad (\text{A.4})$$

$$\int \frac{dx}{\cosh(x)} = 2 \arctan(e^x) \quad (\text{A.5})$$

$$\int \frac{dx}{1-x^2} = \operatorname{arctanh}(x) \quad (\text{A.6})$$

$$\int dx \sqrt{1-x^2} = \frac{1}{2} [x\sqrt{1-x^2} + \arcsin(x)] \quad (\text{A.7})$$

$$\int \frac{dx}{\cos^2(x)} = \tan(x) \quad (\text{A.8})$$

$$\int \frac{dx}{\cosh^2(x)} = \tanh(x) \quad (\text{A.9})$$

- **Definite integrals:**

$$\int_{-\infty}^{+\infty} dx e^{-Ax^2} = \sqrt{\frac{\pi}{A}} \quad (A > 0) \quad (\text{A.10})$$

$$\int_{-\infty}^{+\infty} dx \operatorname{sech}^{2n}(x) = \frac{2^{2n-1}((n-1)!)^2}{(2n-1)!} \quad (\text{A.11})$$

Note: the result of the Gaussian integral (A.10) does not change if the integration variable x is shifted by a finite imaginary amount c , namely if you replace $x \rightarrow x + ic$.